

Coffee Ingestion Enhances One-Mile Running Race Performance

Clarke, N, Richardson, DL, Thie, J & Taylor, R

Author post-print (accepted) deposited by Coventry University's Repository

Original citation & hyperlink:

Clarke, N, Richardson, DL, Thie, J & Taylor, R 2017, 'Coffee Ingestion Enhances One-Mile Running Race Performance' *International Journal of Sports Physiology and Performance*, vol (in press), pp. (in press)

<https://dx.doi.org/10.1123/ijsp.2017-0456>

DOI 10.1123/ijsp.2017-0456

ISSN 1555-0265

Publisher: Human Kinetics

Accepted author manuscript version reprinted, by permission, from *International Journal of Sports Physiology and Performance*, 2017,

<https://doi.org/10.1123/ijsp.2017-0456>

Copyright © and Moral Rights are retained by the author(s) and/ or other copyright owners. A copy can be downloaded for personal non-commercial research or study, without prior permission or charge. This item cannot be reproduced or quoted extensively from without first obtaining permission in writing from the copyright holder(s). The content must not be changed in any way or sold commercially in any format or medium without the formal permission of the copyright holders.

This document is the author's post-print version, incorporating any revisions agreed during the peer-review process. Some differences between the published version and this version may remain and you are advised to consult the published version if you wish to cite from it.

Article Title: Coffee Ingestion Enhances One-Mile Running Race Performance.

Submission type: Original investigation

Preferred Running Title: Coffee Ingestion and Mile Race Performance

Authors: Neil D. Clarke¹, Darren L. Richardson¹, James Thie², and Richard Taylor³.

Author Affiliations:

¹ School of Life Sciences, Faculty of Health and Life Sciences, Coventry University, Coventry, UK.

² Cardiff School of Sport, Cardiff Metropolitan University, Cardiff, UK.

³ Sport, Exercise and Health Research Centre, Newman University, Birmingham, UK.

Corresponding author:

Neil D. Clarke, School of Life Sciences, Faculty of Health Health and Life Sciences, Coventry University, Priory Street, CV1 5FB, UK.

Telephone: (44) 24776 65 8559

Fax: (44) 2476 88 7688

Email: Neil.clarke@coventry.ac.uk

Abstract word count: 245

Text-only word count: 3064

Number of figures and tables: 2 figures and 1 table

Abstract

Purpose: Caffeine, often in the form of coffee, is frequently supplemented by athletes in an attempt to facilitate improved performance during exercise. The aim of the present study was to investigate the effectiveness of coffee ingestion as an ergogenic aid prior to a one-mile (1609 m) race. **Methods:** In a double-blind, randomised, crossover, placebo-controlled design 13 trained male runners completed a one-mile race 60 minutes following the ingestion of 0.09 g·kg⁻¹ coffee (COF), 0.09 g·kg⁻¹ decaffeinated coffee (DEC), or a placebo (PLA). All trials were dissolved in 300 ml of hot water. **Results:** The race completion time was 1.3% faster following the ingestion of COF (04:35:37 ± 00:10:51 mm:ss) compared with DEC (04:39:14 ± 00:11:21 mm:ss; $P=0.018$; 95%CI: -0.11, -0.01; $d=0.32$) and 1.9% faster compared with PLA (04:41:00 ± 00:09:57 mm:ss; $P=0.006$; 95%CI: -0.15, -0.03; $d=0.51$). A large trial and time interaction for salivary caffeine concentration was observed ($P<0.001$; $\eta_p^2=0.69$) with a very large increase ($6.40 \pm 1.57 \mu\text{g}\cdot\text{ml}^{-1}$, 95%CI: 5.5, 7.3; $d=3.86$) following the ingestion of COF. However, only a trivial difference between DEC and PLA was observed ($P=0.602$; 95%CI: -0.09, 0.03; $d=0.17$). Furthermore, only trivial differences for blood glucose ($P=0.839$; $\eta_p^2=0.02$) and lactate ($P=0.096$; $\eta_p^2=0.18$), and maximal heart rate ($P=0.286$; $\eta_p^2=0.13$) were observed between trials. **Conclusions:** The results of the present study show that 60 minutes after ingesting 0.09 g·kg⁻¹ of caffeinated coffee one-mile race performance was enhanced by 1.9% and 1.3% compared with placebo and decaffeinated coffee respectively, in trained male runners.

Keywords: Caffeine; Ergogenic aid; Competition; Middle distance running.

Introduction

Caffeine, often in the form of coffee¹, is frequently supplemented by athletes in an attempt to facilitate improved performance during exercise. However, the available research typically focuses on the ingestion of 3-8 mg·kg⁻¹ of anhydrous caffeine rather than coffee¹. Caffeine non-selectively blocks both adenosine receptors and competitively inhibits the action of adenosine, which during exercise, can lower pain perception, increase neuro-excitability and sustain motor unit firing². Therefore, adenosine receptor antagonism is the leading hypothesis as to how caffeine could have an ergogenic effect on exercise performance². Despite this, Astorino and Roberson³ concluded that the mechanism by which caffeine provides an ergogenic effect in high intensity exercise is likely to be multifactorial, with central factors such as adenosine antagonism being the most probable mechanism, but with reductions in perceived exertion, and increased reaction time, cognition, and mood also having an influence on performance.

Recently, there has been debate as to whether the same ergogenic benefits observed from caffeine ingestion can be obtained from consuming caffeine in the form of coffee, especially as Liguori et al.⁴ reported that peak salivary caffeine concentration was faster and higher following coffee ingestion, compared with a caffeine capsule. However, Graham et al.⁵ concluded that only 4.5 mg·kg⁻¹ of anhydrous caffeine increased exercise distance by 2-3 km when running at 85% $\dot{V}O_{2max}$ until voluntary exhaustion and proposed that chlorogenic acids, and possibly other ingredients within coffee, nullified the ergogenic benefits of caffeine. In contrast, Hodgson et al.⁶ compared 5 mg·kg⁻¹ of anhydrous caffeine to coffee during a cycling time trial lasting approximately 75 min and observed that both coffee and caffeine ingestion were beneficial to exercise performance by

similar magnitudes (~5%). Furthermore, Trexler et al.⁷ concluded that providing 3-5 mg·kg⁻¹ of caffeine in the form of coffee or anhydrous caffeine improved repeated sprint performance and Wiles et al.⁸ observed that 1500 m running performance improved by 4.2 s following the ingestion of 3 g of coffee (150-200 mg caffeine). In addition, Richardson and Clarke⁹ reported that during resistance exercise, the total weight lifted during back squats was 22% higher following the ingestion of coffee, providing 5 mg·kg⁻¹ of caffeine, when compared with a placebo. The inconsistency in performance outcomes associated with coffee ingestion may be due to the blend of coffee ingested and consequently substantial variation in chlorogenic acid content and caffeine concentration of the coffee¹⁰, exercise modality¹¹, and training status¹². However, Higgins et al.¹ concluded that coffee providing 3 mg·kg⁻¹ of caffeine may be used as an alternative to anhydrous caffeine to improve endurance performance.

While sporting performance is more than just a physiological construct, many of the interventions that are researched tend to be physiological in nature, such as nutritional or training. Furthermore, to be a valid simulation of competitive performance, the protocol should provide similar physiological (and psychological) responses to actual performance¹³ but actual competitive events are seldom investigated. Consequently, with studies suggesting that coffee may improve exercise performance^{8,14}, it would be beneficial to investigate the effect of coffee ingestion on actual race performance, as no previous study has investigated the effect of coffee ingestion on performance in ecologically valid situations such as a competitive race. Therefore, the aim of the present study was to provide an insight into the effectiveness of coffee ingestion as an ergogenic aid prior to a one-mile (1609 m) race in trained males.

Methods

Participants

Fifteen trained middle distance male runners began the study, although only 13 (Mean \pm SD age: 24 ± 6 years, height: 1.81 ± 0.07 m, body mass, 69.3 ± 4.7 kg, 800 m personal best performance time: $01:54:43 \pm 00:33:19$ mm:ss, 1500 m personal best performance time: $04:07:56 \pm 00:15:18$ ms:ss; one mile personal best performance time: $04:37:33 \pm 00:23:25$ mm:ss) completed all the races and are therefore included in the data analysis. In a double-blind, Latin-square randomisation, crossover, placebo-controlled design, participants completed a one-mile (1609 m) race following the ingestion of $0.09 \text{ g}\cdot\text{kg}^{-1}$ coffee (COF), $0.09 \text{ g}\cdot\text{kg}^{-1}$ decaffeinated coffee (DEC), or a placebo (PLA). All procedures were undertaken in accordance with the Declaration of Helsinki and approved by the institutional ethics committee. Participants were made fully aware of the exact procedures, including any risks and benefits of participation in the study before providing written informed consent.

On separate days, trials were performed on an indoor 200 m running track with banked curves (temperature: $19.2 \pm 0.9^\circ\text{C}$ and relative humidity: $45 \pm 3\%$) at the same time (17:30) to minimise performance variation due to circadian variation¹⁵. Participants were instructed to abstain from caffeine, alcohol, and strenuous activity for 12 h prior to each race. Caffeine was not withheld for the 24-h period prior because it has been demonstrated that a $3 \text{ mg}\cdot\text{kg}^{-1}$ body mass dose of caffeine improves performance irrespective of whether a withdrawal period is imposed on habitual caffeine users¹⁶. In addition, the participants had not been taking any additional supplements, such as beta

alanine, that may influence running performance for at least three months prior to data collection. Trials were separated by a period of seven days to ensure complete recovery. Habitual caffeine consumption was assessed using an adapted version of the Landrum et al.¹⁷ caffeine consumption questionnaire. Many participants were regular consumers of caffeine (mean \pm SD: 171 ± 250 mg \cdot day⁻¹ and range: 0-705 mg \cdot day⁻¹) with low habitual caffeine consumption defined as <300 mg \cdot day⁻¹ and >300 mg \cdot day⁻¹ was defined as high¹⁸. In addition, a 24-hour dietary record was completed by each participant prior to the first race; it was then photocopied and handed back to the participants so that the same diet could be repeated for subsequent trials (Daily energy: 2738 ± 650 Kcal; Carbohydrate: 394 ± 95 g; Protein: 95 ± 32 g; Fat: 98 ± 29 g; Water: 2409 ± 2132 ml).

Coffee Treatments

Each participant competed in three races following the ingestion of the following: 0.09 g \cdot kg⁻¹ coffee (Nescafé Original, Nestlé, UK; COF), 0.09 g \cdot kg⁻¹ decaffeinated coffee (Nescafé Original Decaffeinated, Nestlé, UK; DEC), and a placebo (PLA). Nescafé original coffee (from the same batch) was used in the coffee trials as it has been reported by Hodgson et al.⁶ to contain 3.4 g of caffeine per 100 g of coffee meaning that each participant consumed 0.09 g \cdot kg⁻¹ of coffee to achieve the 3 mg \cdot kg⁻¹ of caffeine required. Nescafé decaffeinated coffee was used and provides ~ 0.17 mg \cdot kg⁻¹ of caffeine⁶ and was prepared in identical fashion and concentration to the coffee trial. All trials were dissolved in 300 ml of hot water ($65.8 \pm 3.1^\circ\text{C}$) and served in lidded cups. The placebo trial was hot water of the same volume and temperature as the other trials with coffee flavour (Espresso Coffee Flavouring Compound, MSK Ingredients, UK) and colour (Brown Food

Colouring, Lakeland, UK) added to maintain treatment blinding and to ensure all treatments tasted similar. Participants had a maximum of five minutes to fully consume the treatment beverage.

Procedures

On arrival at the athletics track the body mass of each participant was recorded and saliva and capillary blood samples were collected before ingesting the test drink (baseline). The participants then performed their usual pre-race standardised warm-up lasting 45 min, which consisted of a 2000 m self-paced jog followed by an individual static and dynamic stretching programme and ending with 3 × 80 m strides. Then, 60 minutes after the ingestion of the fluid, further saliva and capillary blood samples were collected (post-warm-up) and the one-mile race was run. Collection of saliva and blood samples were repeated post-race. No encouragement was given and watches were removed. The one mile was timed using electronic timing lights (ALGE Timing GmbH, Lustenau, Austria). Time to completion was not given to the participants until after the end of the study. To ensure maximum effort, the races were handicapped based on the participants' personal best time and adjusted based on their performance for the following race to ensure each trial was a competitive race e.g. a participant with a one-mile personal best of 04:22 mm:ss started 3 s after a participant with a personal best of 04:25 mm:ss. Furthermore, the participants received £2 for each place in every race i.e. £2 for 13th and £26 for 1st and an additional £50 gift voucher for the overall winner based on accumulated prize money.

Measurements

Heart rate (HR) was measured throughout each race using a short-range telemetry HR transmitter strap recording at 1 s intervals (Polar Team 2 System, Polar Electro Oy, Kempele, Finland). Saliva samples (minimum 0.5 ml by the passive drool technique) were obtained immediately before fluid ingestion to establish compliance with the washout period, one-hour post-ingestion and following the race. Participants were instructed to *expectorate* into a 20-ml plastic cup, and the sample was then transferred to a capped glass vial that was immediately placed in a portable -20°C freezer before being transferred to a -80°C freezer for subsequent analysis of caffeine concentration using a standard enzyme-linked immunoassay kit (Caffeine ELISA Kit; Creative Diagnostics, Shirley, USA). At the same time points, a capillary blood sample was drawn from the index finger for determination of blood glucose and lactate concentrations (*Biosen C-line*, EKF-diagnostic GmbH, Germany).

Data Analysis

An *a priori* power calculation based for race time revealed that a sample size of ten participants were necessary to detect a statistical difference given an estimated effect size of 0.5, a $1-\beta$ error probability of 0.8 and a *P* value of less than 0.05. Data are reported as the mean \pm the standard deviation (SD). The Shapiro-Wilk test was applied to the data to assess for a normal distribution. All variables, except for salivary caffeine and blood glucose and lactate were assessed using a one-way analysis of variance (ANOVA) with repeated measures. Salivary caffeine concentration and blood glucose and lactate were analysed with a two-way ANOVA with repeated measures.

Sphericity was analysed by Mauchly's test of sphericity followed by the Greenhouse-Geisser adjustment where required. Where any differences were identified, pairwise comparisons with Bonferroni correction were used to show where they lay. The data was analysed using IBM SPSS Statistics for Windows, Version 24.0 (Armonk, NY: IBM Corp.). Furthermore, 95% confidence intervals (95%CI) and effect sizes using partial eta squared (η_p^2), defined as trivial (<0.10), small (0.10-0.24), moderate (0.25-0.39) or large (≥ 0.40), and Cohen's d , defined as trivial (<0.20), small (0.20-0.49), moderate (0.50-0.79) or large (≥ 0.80) according to the cut-offs suggested by Cohen¹⁹, were also calculated. In addition, the changes in race performance were standardized and expressed as a factor of the smallest worthwhile change, based on Cohen's effect size principle.

Results

The results show a large main effect for trial ($F_{2,24}=9.524$; $P=0.001$; $\eta_p^2=0.44$; Figure 1) with race completion time being faster following the ingestion of COF (04:35:37 \pm 00:10:51 mm:ss) compared with DEC (04:39:14 \pm 00:11:21 mm:ss; $P=0.018$; 95%CI: -0.11, -0.01; $d=0.32$) and PLA (04:41:00 \pm 00:09:57 mm:ss; $P=0.006$; 95%CI: -0.15, -0.03; $d=0.51$). However, only a trivial difference between DEC and PLA was observed ($P=0.602$; 95%CI: -0.09, 0.03; $d=0.17$). Overall, race performance was improved by $1.9 \pm 1.7\%$ (95%CI: 1, 2.8) and $1.3 \pm 1.4\%$ (95%CI: 0.5, 2.0) following the ingestion of COF, compared with PLA and DEC respectively with a $0.6 \pm 1.6\%$ (95%CI: -0.2, 1.5) improvement following DEC compared with PLA. Furthermore, no order effect was observed ($F_{2,24}=0.041$; $P=0.959$; $\eta_p^2=0.00$).

A large trial and time interaction for salivary caffeine concentration was observed ($F_{2,21}=24.260$; $P<0.001$; $\eta_p^2=0.69$; Figure 2) with a very large increase ($6.40 \pm 1.57 \mu\text{g}\cdot\text{ml}^{-1}$, 95%CI: 5.5, 7.3, $d=3.86$) in salivary caffeine concentration between pre-race and post-warm-up observed following the ingestion of COF. In contrast, only small changes were observed during DEC ($0.87 \pm 1.68 \mu\text{g}\cdot\text{ml}^{-1}$, 95%CI: -0.1, 1.8, $d=0.43$) and PLA ($0.83 \pm 2.06 \mu\text{g}\cdot\text{ml}^{-1}$, 95%CI: -0.3, 2.0, $d=0.41$). Only trivial differences between trials for blood glucose ($F_{2,24}=0.177$; $P=0.839$; $\eta_p^2=0.02$; Table 1) and lactate ($F_{2,24}=2.584$; $P=0.096$; $\eta_p^2=0.18$; Table 1), and maximum heart rate ($F_{1,10}=1.345$; $P=0.286$; $\eta_p^2=0.13$; Table 1) were observed.

Discussion

The aim of the present study was to determine the effect of ingesting caffeinated coffee on one-mile (1609 m) race performance. The main findings are that the ingestion of COF markedly increased salivary caffeine concentration and improved one-mile race performance compared with DEC and PLA. In addition, blood glucose and lactate concentration, and maximal heart rate were unaffected by the ingestion of COF or DEC.

Following the ingestion of caffeinated coffee race performance was improved by 1.9% and 1.3% compared with PLA and DEC respectively. The ergogenic benefits of caffeine ingestion are well documented (e.g.^{3,21}). However, fewer studies have documented the effects of coffee ingestion. In support of the findings of the present study, Wiles et al.⁸ reported that the ingestion of 3 g of caffeinated coffee, containing approximately 150-200 mg of caffeine, improved 1500 m treadmill running performance by 4.2 s (1.4%) when compared with decaffeinated coffee. Similarly,

Hodgson et al.⁶ demonstrated that coffee and caffeine ingestion improved time trial performance by approximately 5%. Richardson and Clarke⁹ also reported that resistance exercise performance was improved by 22% following coffee ingestion, when compared with a placebo. Furthermore, Trexler et al.⁷ reported that caffeine and coffee ingestion improved total work performed during repeated sprints (95%CI: 40, 219 J). In contrast, Clarke et al.²⁰ observed that the ingestion of coffee had little effect on repeated sprint cycling performance in relatively untrained males. Graham et al.⁵ previously suggested that the bioactive compounds in coffee, such as chlorogenic acids, may attenuate the ergogenic effect of caffeine. However, more recent studies^{6,7,14} have reported both coffee and caffeine ingestion yield similar benefits for exercise performance, most likely through the antagonistic effect on adenosine receptors reducing symptoms of central fatigue¹. One potential reason for the inconsistent findings is training status. It is apparent that caffeine and coffee is less effective for non-trained individuals participating in high-intensity exercise¹², possibly due to trained athletes, like the participants in the present study, having more muscle mass than recreational athletes and the concentration of adenosine receptors (the hypothesised primary target of caffeine²) appearing to be higher in trained compared to untrained individuals²¹. In addition, the high variability in performance that is typical for untrained subjects may also contribute.

Although the purpose of this study was not to determine the mechanism of coffee action, adenosine antagonism, enhanced motor unit recruitment and reduced perception of pain and exertion have been proposed to explain the effects of caffeine supplementation on sport performance². However, since caffeine interacts with many tissues, it is difficult to independently investigate its effects on the central and peripheral nervous systems, and metabolism²². When specifically examining exercise of the nature in the present study, the primary mechanisms by which caffeine exerts its

ergogenic effects are considered to arise from the antagonism of adenosine receptors leading to an increase in neurotransmitter release and motor unit firing rates, pain suppression, reduced fatigue and improved neuromuscular performance². Another possible mechanism through which caffeine may improve performance is by increasing the secretion of β -endorphins²³, which may, at least partially, explain the mechanism by which caffeine attenuates pain sensation²⁴ and rating of perceived exertion²⁵ during exercise, thereby decreasing perceptions of effort and improving performance, as observed in the present study.

In addition to its impact on the central nervous system, caffeine can affect substrate utilization during exercise with a decrease in glycogen utilization and an increase in free fatty acid mobilization²⁶. However, alterations in substrate utilization is unlikely to mediate performance changes in the present study due to fat metabolism representing a minor substrate and muscle glycogen not being limiting². In addition, de Paulis, et al.²⁷ demonstrated that coffee ingestion caused a blunted response of adrenaline which was attributed to chlorogenic acids antagonising the binding of caffeine to adenosine receptors. Furthermore, Hodgson, et al.⁶ reported that the increase in glucose, fatty acids and glycerol observed when ingesting caffeine had an attenuated response following coffee ingestion, which was attributed to other compounds in coffee inducing subtle effects on the antagonism of A₁ and A_{2A} adenosine receptors. Despite these findings, the performance improvements observed by Hodgson, et al.⁶ were similar for both caffeine and coffee leading the authors to conclude coffee ingestion may alter the metabolic effects, but not the ergogenic effects on performance. In support of this observation, McLellan and Bell¹⁴ and Trexler et al.⁷ reported similar performance benefits from ingesting coffee and caffeine. However, in the present study no differences in blood glucose and lactate concentrations, and heart rate were

observed, although these values were comparable with previous studies^{5,8,14}. Overall, it is probable that coffee and caffeine supplementation enhances physical performance via a combined effect on both the central and peripheral systems.

The present study is not without limitations. The use of a 24-hour dietary record to ensure pre-trial standardisation, whilst being an acceptable method and a good reflection of standard nutritional practice, there may be some concern about the accuracy. In addition, the baseline salivary caffeine concentrations suggest that caffeine was not abstained from prior to each trial. However, both factors are likely to provide a more ecologically valid condition and represent typical pre-race preparation. A further potential limitation is the wide range of the participants' habitual caffeine intake, although Gonçalves et al.²⁸ recently demonstrated that during ~30 min cycling time trials, performance was not influenced by habitual caffeine consumption.

Practical Implications

The coefficient of variation (CV) for running performance at distances between 1500 m and 10,000 m for elite males has been reported to be 1.4% (90%CI: 1.2-1.6)²⁹ and the smallest worthwhile enhancement in performance of similar distances to be 0.3-0.5 of the CV²⁹. In the present study, based on the placebo condition, this corresponds to a 1.01-2.25 s improvement in the time taken to complete the mile. Therefore, the performance improvements observed [*vs.* PLA: 5.38 ± 4.75 s (95%CI: 2.80, 7.97); *vs.* DEC: 3.62 ± 3.84 s (95%CI: 1.53, 5.70)] following the ingestion of 300 ml of caffeinated coffee providing 3 mg·kg⁻¹ of caffeine 60 minutes before a race could be considered practically meaningful during actual competitions. However, athletes need to be aware

that the type of coffee and brewing method alters the caffeine and chlorogenic acids content¹⁰ and inter-individual variation in pharmacodynamic and pharmacokinetic polymorphisms³⁰ may cause variation in the responses to caffeine and its ergogenic effects.

Conclusion

In conclusion, the ingestion of caffeinated coffee 60 minutes before a one-mile race markedly increased salivary caffeine concentration and improved race performance compared with decaffeinated coffee and a placebo solution. These findings suggest that the ingestion of caffeinated coffee is a suitable source of caffeine prior to a one-mile running race.

Acknowledgements

The authors express appreciation to Professor Derek Renshaw for his input on the salivary caffeine analysis. Funding for this research was received from The Institute for Scientific Information on Coffee. The result of the present study does not constitute endorsement of the product by the authors or the journal.

References

1. Higgins S, Straight CR, Lewis RD. The Effects of Pre-Exercise Caffeinated-Coffee Ingestion on Endurance Performance: An Evidence-Based Review. *Int J Sports Nutr Exerc Metab.* 2016; 26: 221-239.

2. Davis JK, Green JM. Caffeine and Anaerobic Performance: Ergogenic Value and Mechanisms of Action. *Sports Med.* 2009; 39: 813-832.
3. Astorino TA, Roberson DW. Efficacy of Acute Caffeine Ingestion for Short-Term High-Intensity Exercise Performance: A Systematic Review. *J Strength Cond Res.* 2010; 24: 257-265.
4. Liguori A, Hughes JR, Grass JA Absorption and Subjective Effects of Caffeine from Coffee, Cola and Capsules. *Pharmacol Biochem Behav.* 1997; 58: 721-726.
5. Graham TE, Hibbert E, Sathasivam P. Metabolic and exercise endurance effects of coffee and caffeine ingestion. *J Appl Physiol.* 1998; 85: 883–9.
6. Hodgson AB, Randell RK, Jeukendrup AE The Metabolic and Performance Effects of Caffeine Compared to Coffee during Endurance Exercise. *PLoS One.* 2013; 8: e59561.
7. Trexler ET, Smith-Ryan AE, Roelofs EJ, Hirsch KR, Mock MG. Effects of coffee and caffeine anhydrous on strength and sprint performance. *Eur J Sport Sci.* 2016; 16: 702-707.
8. Wiles JD, Bird SR, Hopkins J, Riley M. Effect of Caffeinated Coffee on Running Speed, Respiratory Factors, Blood Lactate and Perceived Exertion during 1500-m Treadmill Running. *Br J Sports Med.* 1992; 26: 116-120.

9. Richardson DL, Clarke ND. Effect of Coffee and Caffeine Ingestion on Resistance Exercise Performance. *J Strength Cond Res.* 2016; 30: 2892-2900.
10. Fujioka K, Shibamoto T. Chlorogenic Acid and Caffeine Contents in Various Commercial Brewed Coffees. *Food Chemistry.* 2008; 106: 217-221.
11. Hoffman JR, Kang J, Ratamess NA, Jennings PF, Mangine GT, Faigenbaum, AD. Effect of nutritionally enriched coffee consumption on aerobic and anaerobic exercise performance. *J Strength Cond Res.* 2007; 21: 456-459.
12. Goldstein ER, Ziegenfuss T, Kalman D, et al. International Society of Sports Nutrition Position Stand: Caffeine and Performance. *J Int Soc Sports Nutr.* 2010; 7: 5.
13. Currell K, Jeukendrup A. Validity, reliability and sensitivity of measures of sporting performance. *Sports Med.* 2008; 34: 297-316.
14. McLellan TM, Bell DG. The Impact of Prior Coffee Consumption on the Subsequent Ergogenic Effect of Anhydrous Caffeine. *Int J Sports Nutr Exerc Metab.* 2004; 14: 698-708.
15. Drust B, Waterhouse J, Atkinson G, Edwards B, Reilly T. Circadian rhythms in sports performance - an update. *Chronobiol Int.* 2005; 22: 21-44.

16. Irwin C, Desbrow B, Ellis A, O'Keeffe B, Grant G, Leveritt M. Caffeine withdrawal and high-intensity endurance cycling performance. *J Sports Sci.* 2011; 29: 509-515.
17. Landrum RE, Meliska CJ, Loke WH. Effects of Caffeine and Task Experience on Task Performance. *Psychologia.*1988; 31: 91-97.
18. Westerterp-Plantenga MS, Lejeune MP, Kovacs EM. Body Weight Loss and Weight Maintenance in Relation to Habitual Caffeine Intake and Green Tea Supplementation. *Obes Res.* 2005; 13: 1195-1204.
19. Cohen J. A power primer. *Psychological Bulletin.* 1992; 112: 155-159.
20. Clarke ND, Baxter H, Fajemilua E, Jones V, Oxford S, Richardson D, Wyatt C, Mundy P. Coffee and Caffeine Ingestion Have Little Effect on Repeated Sprint Cycling in Relatively Untrained Males. *Sports.* 2016; 4: 45.
21. Mizuno M, Kimura Y, Tokizawa K, Ishii K, Oda K, Sasaki T, Nakamura Y, Muraoka I, Ishiwata K: Greater adenosine A(2A) receptor densities in cardiac and skeletal muscle in endurance-trained men: a [11C]TMSX PET study. *Nucl Med Biol* 2005, 32:831-836.
22. Spriet LL. Exercise and sport performance with low doses of caffeine. *Sports Med.* 2014; 44: S175-S184.

23. Laurent D, Schneider KE, Prusaczyk WK, Franklin C, Vogel SM, Krssak M, Petersen KF, Goforth HW, Shulman GI. Effects of caffeine on muscle glycogen utilization and the neuroendocrine axis during exercise. *J Clin Endocrinol Metab.* 2000; 85: 2170-2175.
24. Gliottoni RC, Motl RW. Effect of caffeine on leg muscle pain during intense cycling exercise: Possible role of anxiety sensitivity. *Int J Sports Nutr Exerc Metab.* 2008; 18: 103-115.
25. Doherty M, Smith PM. Effects of caffeine ingestion on rating of perceived exertion during and after exercise: a meta-analysis. *Scand J Med Sci Sports.* 2005; 15: 69–78.
26. Spriet LL, MacLean DA, Dyck DJ, Hultman E, Cederblad G, Graham TE. Caffeine ingestion and muscle metabolism during prolonged exercise in humans. *Am J Physiol.* 1992; 262: E891-898.
27. de Paulis T, Schmidt DE, Bruchey AK, Kirby MT, McDonald MP, Commers P, Lovinger DM, Martin PR. Dicinnamoylquinides in roasted coffee inhibit the human adenosine transporter. *Eur J Pharmacol.* 2002; 442: 215-223.
28. Gonçalves LDS, Painelli VDS, Yamaguchi G, de Oliveira LF, Saunders B, Silva RPD, Maciel E, Artioli GG, Roschel H, Gualano B. Dispelling the myth that habitual caffeine consumption influences the performance response to acute caffeine supplementation. *J Appl Physiol.* 2017; Epub ahead of print.

29. Hopkins WG. Competitive performance of elite track-and-field athletes: Variability and smallest worthwhile enhancements. *Sportscience*. 2005; 9: 17-20.

30. Yang A. Palmer, A.A., and de Wit, H. (2010) Genetics of caffeine consumption and responses to caffeine. *Psychopharmacology* .211, 245-257.




Figure 1: Mean and individual race completion times for each condition.

* COF time faster than PLA ($d=0.32$) and DEC ($d=0.51$).




Figure 2: Mean (\pm SD) salivary caffeine concentration during each condition.

* COF higher than PLA ($d=2.34$) and DEC ($d=2.47$).

COF higher than PLA ($d=1.99$) and DEC ($d=1.83$).

Table 1: Blood glucose and lactate at baseline, post-warm-up and post-race, and heart rate post-warm-up and maximum value achieved during each race.

	COF	DEC	PLA
<i>Blood Glucose (mmol·l⁻¹)</i>			
Baseline	4.64 ± 0.71	4.33 ± 0.83	4.72 ± 0.54
Post-Warm-Up	4.86 ± 0.52	4.78 ± 0.48	4.85 ± 0.79
Post-Race	7.47 ± 1.17	7.54 ± 1.15	7.14 ± 0.92
<i>Blood Lactate (mmol·l⁻¹)</i>			
Baseline	1.94 ± 0.52	1.91 ± 0.64	1.66 ± 0.45
Post-Warm-Up	4.07 ± 0.99	3.99 ± 1.11	3.99 ± 1.08
Post-Race	14.35 ± 3.52	13.42 ± 3.26	12.40 ± 3.02
<i>Heart Rate (beats·min⁻¹)</i>			
Post-Warm-Up	104 ± 11	104 ± 13	106 ± 5
Maximal	187 ± 10	182 ± 17	190 ± 7