

ELN and FBN2 gene variants as risk factors for two sports-related musculoskeletal injuries

Khoury, LE, Posthumus, M, Collins, M, van der Merwe, W, Handley, C, Cook, J & Raleigh, SM

Author post-print (accepted) deposited by Coventry University's Repository

Original citation & hyperlink:

Khoury, LE, Posthumus, M, Collins, M, van der Merwe, W, Handley, C, Cook, J & Raleigh, SM 2015, 'ELN and FBN2 gene variants as risk factors for two sports-related musculoskeletal injuries' *International Journal of Sports Medicine*, vol 36, no. 4, pp. 333-337

<https://dx.doi.org/10.1055/s-0034-1390492>

DOI [10.1055/s-0034-1390492](https://dx.doi.org/10.1055/s-0034-1390492)

ISSN 0172-4622

ESSN 1439-3964

Publisher: Georg Thieme Verlag

© 2016 Georg Thieme Verlag KG

Copyright © and Moral Rights are retained by the author(s) and/ or other copyright owners. A copy can be downloaded for personal non-commercial research or study, without prior permission or charge. This item cannot be reproduced or quoted extensively from without first obtaining permission in writing from the copyright holder(s). The content must not be changed in any way or sold commercially in any format or medium without the formal permission of the copyright holders.

This document is the author's post-print version, incorporating any revisions agreed during the peer-review process. Some differences between the published version and this version may remain and you are advised to consult the published version if you wish to cite from it.

1 ***ELN* and *FBN2* gene variants as risk factors for two sports related musculoskeletal**
2 **injuries**

3 **Abstract**

4 The *ELN* and *FBN2* proteins are important in extracellular matrix function. The *ELN*
5 rs2071307 and *FBN2* rs331079 gene variants have been associated with soft tissue
6 pathologies. We aimed to determine whether these variants were predisposing factors for
7 both Achilles tendinopathy (AT) and anterior cruciate ligament (ACL) ruptures.

8 For the AT study, 135 cases (TEN group) and 239 asymptomatic controls were recruited.

9 For the ACL rupture study our cohort consisted of 141 cases (ACL group) and 219 controls.

10 Samples were genotyped for both the *ELN* rs2071307 and *FBN2* rs331079 variants using
11 TaqMan assays. Analysis of variance and chi-squared tests were used to determine whether
12 either variant was associated with AT or ACL rupture with significance set at $p < 0.05$.

13 The GG genotype of the *FBN2* variant was significantly over-represented within the TEN
14 group ($p = 0.035$; OR=1.83; 95% CI 1.04–3.25) compared to the CON group. We also found
15 that the frequency of the G allele was significantly different between the TEN ($p = 0.017$;
16 OR=1.90; 95% CI 1.11–3.27) and ACL groups ($p = 0.047$; OR=1.76; 95% CI 1.00–3.10)
17 compared to controls. The *ELN* rs207137 variant was not associated with either AT or ACL
18 rupture. In conclusion, DNA sequence variation within the *FBN2* gene is associated with both
19 AT and ACL rupture.

20 **Keywords** Gene; Achilles Tendinopathy; Tendon rupture; Ligament rupture; Injury
21 prevention.

22

23 **Introduction**

24 Injury to the Achilles tendon and the anterior cruciate ligament are severe traumas typically
25 sustained during sports activities. Achilles tendon injuries, including chronic Achilles
26 tendinopathy (AT) and acute Achilles tendon rupture, are prevalent within athletic populations
27 [26]. Indeed, the lifetime incidence of AT is approximately 10% in the general population and
28 as high as 50% within competitive runners [12]. Chronic AT may be due, in part, to excessive
29 exposure of the Achilles tendon to acute or repetitive mechanical loading forces experienced
30 during exercise [15]. Anterior cruciate ligament (ACL) ruptures have low lifetime prevalence
31 in the general population but have been reported to be nearly 80% among netball players [7].
32 The most common mechanism of ACL rupture involves a sudden change in an athlete's
33 direction or rapid deceleration [19]. Both AT and ACL ruptures are complex multifactorial
34 phenotypes with several intrinsic and extrinsic risk factors. However, the exact aetiology is
35 not yet fully understood [3].

36 Among the intrinsic risk factors, several genetic sequence variants have been shown to
37 increase the risk (predispose individuals) to AT and ACL ruptures. Variants within the *TNC*
38 [3], *MMP3* [29], *GDF5* [23] and *TIMP2* genes [5] are associated with the risk of AT.
39 Furthermore, variants within the *COL1A1* [22] and *COL12A1* genes [24] have also been
40 associated with ACL ruptures. Interestingly, a variant within the *COL5A1* gene was found to
41 be associated with both AT [16] and ACL ruptures [25]. These findings show that both
42 chronic AT and ACL ruptures have a partial polygenic basis where complex interactions
43 between genes and the environment are likely to exacerbate the risk of both types of injuries
44 [3]. All the genes described above encode proteins with either a structural or regulatory role
45 in maintaining the homeostasis of the soft tissue extracellular matrix (ECM). Therefore, it is
46 fair to assume that other genes, which code for additional regulatory components of the ECM
47 might also be candidates for AT and ACL rupture.

48 Elastin (ELN) is an insoluble polymer composed of several tropoelastin molecules covalently
49 bound to each other by cross-links [31]. ELN proteins contribute to tendon and ligament
50 elasticity by allowing them to stretch and return to their original state. These proteins have an
51 important load-bearing role in musculoskeletal tissues and are expressed in places where
52 mechanical energy is stored [8]. The *ELN* rs2071307 gene variant has been shown to be
53 associated with other multifactorial conditions of the extracellular matrix, such as aortic
54 stenosis [6] and aortic aneurysm [33]. Interestingly, the *ELN* rs2071307 variant is located
55 within exon 20 of the gene and is a non-synonymous SNP. It is predicted to be deleterious
56 (Queen's University. <http://compbio.cs.queensu.ca/F-SNP/>) since it substitutes a hydrophobic
57 amino acid glycine with a hydrophilic serine residue (National Center for Biotechnology
58 Information. <http://www.ncbi.nlm.nih.gov/projects/SNP/>). This substitution may disrupt the
59 integrity of the microfibrils rendering them more prone to damage [18] and therefore this
60 variant may predispose to soft tissue damage during sports performance.

61 Fibrillins are large glycoproteins present in the extracellular matrix of tendons and ligaments
62 [2]. Both fibrillin-1 (FBN-1) and fibrillin-2 (FBN-2) share high amino acid homology and are
63 involved in providing strength and flexibility to various soft tissues. FBN-2 is abundant in
64 elastic tissues, such as tendons and ligaments [35] where it plays an important role in the
65 assembly of elastic fibres [2]. Mutations within the *FBN2* gene are known to associate with
66 musculoskeletal pathologies such as congenital contractural arachnodactyly [9].
67 Furthermore, the rs331079 variant located within intron 7 of the gene (University of Florida.
68 www.snpper.chip.org) has previously been associated with intracranial aneurysms [32].

69 As both the *FBN2* rs331079 and the *ELN* rs2071307 variants associate with other conditions
70 related to the extracellular matrix we considered them as possible risk determinants for both
71 AT and ACL rupture. Accordingly, the aim of this study was to test that hypothesis.

72

73 **Material and Methods**

74 One hundred and thirty five (60 Australian (AUS) and 75 South African (SA)) Caucasian
75 participants with Achilles tendinopathy (TEN group) were recruited to this study from the
76 Musculoskeletal Research Centre at La Trobe University in Melbourne, and from the Medical
77 Practice at the Sports Science Institute of South Africa. Furthermore, 239 (143 AUS and 96
78 SA) asymptomatic Caucasian controls (CON groups) were recruited to this study from
79 recreational sports clubs within the Melbourne area in Australia, and within the Cape Town
80 area in South Africa. Chronic AT was clinically diagnosed as described by Mokone et al.[17]
81 in the first manuscript describing the South African AT cohort. The Australian cohort used the
82 same clinical diagnosis described by Mokone et al. In addition, diagnosis was confirmed with
83 soft tissue ultrasound examination in all the AUS and 40 of 75 SA participants. In addition,
84 141 South African Caucasian participants with surgically diagnosed ACL ruptures (ACL
85 group) and 219 apparently healthy (CON group), unrelated, physically active, gender
86 matched South African Caucasian participants without any self-reported history of ligament
87 or tendon injury were recruited for this study as previously described [22]. Seventy four
88 participants sustained the injury through a non-contact mechanism and were analysed as a
89 separate subgroup (NON subgroup).

90 Previous injury data was used as inclusion criteria in the various cohorts analysed. In the
91 AUS Achilles cohort, the CON group had no history of any tendon injury, whereas in the SA
92 Achilles cohort, the CON group merely had no previous history of Achilles tendon injuries. In
93 the case of ACL rupture, the first ACL rupture was documented as the specific inclusion
94 injury. Therefore, by definition, no participant in the ACL group had a previous ACL rupture

95 None of the participants included in this study had symptoms or signs –of Ehlers-Danlos
96 syndrome (EDS), hypermobility or benign hypermobility joint syndrome or other monogenic
97 connective tissue disorders when their medical examinations were reviewed by the medical
98 practitioner [16,17,34].

99 Physical activity data was recorded for the South African Achilles tendinopathy cohort (SA
100 CON and SA TEN), but not for the Australian Achilles tendinopathy cohort (AUS CON and
101 AUS TEN). In addition physical activity data was also recorded for the South African ACL
102 cohort (SA ACL and SA CON). The data recorded for the SA CON and SA TEN groups
103 included total years participated in running and high impact sports, as well as hours per week
104 of participation in the last 2 years. The data reported for the ACL cohort included years of
105 participation in contact sports, non-contact jumping sports, non-contact non-jumping sports
106 and skiing sports. Data were collected as previously described [16, 25].

107 Based on our earlier work, this study had a large enough sample size to detect associations
108 with an OR of 2.0 at $p < 0.05$ with 80% power [28]. All participants gave informed written
109 consent, in accordance with the journal's recommendations [10,11], and all completed a
110 medical and injury history questionnaire. Ethical approval was obtained from the Research
111 Ethics Committees at the University of Cape Town, South Africa, La Trobe University,
112 Australia, Monash University, Australia and the University of Northampton, United Kingdom
113 prior to initiating this work.

114 For the Australian cohort, DNA was extracted from whole blood using Qiagen DNA extraction
115 kits (Flexigene DNA kit, Qiagen P/L, Valencia, California, USA) as per the manufacturer's
116 recommendations. DNA from the South African individuals was extracted from blood using
117 the method described by Lahiri and Nurnberg [14] and modified by Mokone et. al. [16,17].
118 Upon extraction, DNA was frozen at $-20\text{ }^{\circ}\text{C}$ for long-term storage, and smaller aliquots were
119 stored at $4\text{ }^{\circ}\text{C}$ for short term usage.

120 DNA from all participants was genotyped for the *FBN2* rs331079 and *ELN* rs2071307 gene
121 variants using fluorescence-based TaqMan assays (Applied Biosystems, Foster City, CA,
122 USA). PCR reactions contained allele-specific probes and primers in a PCR mastermix
123 containing AmpliTaq DNA Polymerase Gold (Applied Biosystems, Foster City, CA, USA) in a
124 total reaction volume of 12 μL . PCR was performed on an Applied Biosystems

125 StepOnePlus™ real-time PCR system (Applied Biosystems, Foster City, CA, USA).
126 Genotypes were called according to output clustering profiles using Applied Biosystems
127 StepOnePlus™ real-time PCR software Version 2.1 (Applied Biosystems, Foster City, CA,
128 USA). Rox was used as a passive reference to normalise fluorescence signal intensity
129 relative to the amount of sample used.

130 The statistical power of the study was determined using Quanto v1.2
131 (<http://hydra.usc.edu/gxe>). The initial calculations were done using a recessive model and a
132 disease population prevalence of 10%. Assuming a risk allele frequency of 60%, a matched
133 case-control population of 136 individuals per group was adequate to detect an allelic OR of
134 2.0 at a power of 80% and a significance level of 5%.

135 Data were analysed using SPSS Version 20 (SPSS Science Inc, Chicago, Ill, USA) statistical
136 program. A one-way analysis of variance was used to establish if any significant difference
137 existed between the characteristics of the TEN and CON groups within the Australian and
138 South African cohorts as well as between the ACL rupture and CON groups. A chi-squared
139 (χ^2) analysis or Fisher's exact test was used to determine if significance differences existed
140 between genotype and/or allele frequencies, as well as other categorical data between the
141 groups. In all analysis significance was accepted when $p < 0.05$. Adjustments for multiple
142 testing were not conducted as it has been previously described [21] that no appropriate
143 method exists. Furthermore, the Bonferroni adjustment was considered too conservative [21]
144 and inappropriate for a situation like this where there is prior evidence that the gene of
145 interest is associated with a trait [20]. Hardy-Weinberg equilibrium was determined using the
146 program Genepop web version 3.4 (Curtin University. <http://genepop.curtin.edu.au/>).

147

148 Results

149 Running was the predominant sporting activity resulting in Achilles tendon injuries (63.1%,
150 N=65) in the SA cohort. The SA groups were matched for the mean number of years
151 participating in running (CON, 8.7 ± 8.2 yrs, n=95; TEN, 10.0 ± 11.0 yrs, n=62; p=0.402).
152 However, there was a significant difference in hours of training between the two groups
153 (CON, 3.6 ± 3.0 hrs/week, n=91; TEN, 2.4 ± 2.7 hrs/week, n=55; p=0.011), where the SA
154 CON group trained for more hours per week. The SA TEN participants participated in more
155 years of high impact sports compared to the SA CON group in the past (CON, 9.4 ± 8.4 yrs,
156 n=95; TEN, 13.1 ± 11.1 yrs, n=62; p=0.018), however, the SA CON group performed a
157 greater amount of high impact sports during the last 2 years (CON, 3.6 ± 3.1 yrs, n=95; TEN,
158 2.5 ± 12.9 yrs, N=62; p=0.029. Although all AUS participants were physically active
159 individuals, the type of sporting activity involved in, the hours of training and the frequency of
160 activity were not recorded.

161 The SA ACL and SA CON groups were matched for years of participation in contact sports
162 (SA CON, 11.7 ± 7.1 yrs, n=219; SA ACL, 11.5 ± 8.0 yrs, n=141; p=0.892), non-contact
163 jumping sports (SA CON, 27.8 ± 19.9 yrs, n=190; SA ACL, 25.7 ± 22.6 yrs, n=141; p=0.398),
164 non-contact non-jumping sports (SA CON, 11.5 ± 7.1 yrs, n=219; SA ACL, 10.5 ± 8.5 yrs,
165 n=141; p=0.575), and skiing sports (SA CON, 19.1 ± 16.9 yrs, n=219; SA ACL, 8.6 ± 8.5 yrs,
166 n=141; p=0.094).

167 Since the *ELN* rs2071307 and *FBN2* rs331079 allele and genotype frequencies in both of the
168 South African (SA) and Australian (AUS) TEN and CON groups were similar (Supplementary
169 table 1), the data was collectively analysed. The CON and TEN groups were similarly
170 matched for age and gender (Table 1). When co-varied for sex, the two groups were
171 similarly matched for height. Furthermore, when co-varied for sex and age at recruitment, the
172 TEN group was found to be significantly heavier ($p < 0.001$) with larger BMIs ($p < 0.001$) (Table
173 1). The TEN group was recruited on average 5.1 years after the initial injury.

174 Participants in the AUS TEN group carrying the *ELN* rs2071307 AA (53.1 ± 11.6 , n=10)
175 genotype were significantly ($p=0.005$) older when they reported their initial Achilles tendon
176 injury when compared to those with a GG (37.2 ± 12.6 , n=16) or GA (37.8 ± 13.6 , n=32)
177 genotype. There were, however, no significant differences in the mean ages of the three
178 genotype groups in the CON AUS group (GG: 40.7 ± 11.8 , n=48; GA: 37.4 ± 12.2 , n=68; AA:
179 40.1 ± 12.1 , n=24; $p=0.323$). There were no other significant genotype effects of either
180 variants with respect to height, weight, BMI, or sex in the AT group (data not shown).
181 Furthermore, the investigated variants did not show any interaction with age, height, weight,
182 BMI and sex in the ACL population (data not shown).

183 The genotype frequency distributions of the *FBN2* rs331079 and the *ELN* rs2071307 variants
184 within the AT and the ACL rupture groups are shown in table 2. In the combined TEN cohort,
185 the *FBN2* rs331079 genotype frequency was significantly different ($p=0.035$) between the
186 CON (GG, 76.9%; GC + CC, 23.1%) and TEN (GG, 85.9%; GC + CC, 14.1%) groups (Table
187 2). The GG genotype was significantly over-represented within the TEN group ($p=0.035$;
188 OR=1.83; 95% CI 1.04 – 3.25). We also found a significant ($p=0.017$; OR=1.90; 95% CI 1.11
189 – 3.27) allele frequency distribution difference for the *FBN2* rs331079 variant between the
190 CON (G, 87.4%; C, 12.6%) and TEN (G, 93.0%; C, 7.0%) groups (Table 2). Similarly, we
191 also found a significant ($p=0.047$; OR=1.76; 95% CI 1.00 – 3.10) allele frequency distribution
192 difference of the rs331079 locus between the CON (G, 89.3%; C, 10.7) and ACL (G, 93.6%;
193 C, 6.4%) groups. Also, in the AT population, there were no significant *ELN* rs2071307
194 genotype ($p=0.795$) or allelic ($p=0.741$) frequency differences between the CON and TEN
195 groups (Table 2).

196 Although not significant, we found a tendency towards an allelic ($p=0.064$) association for the
197 *ELN* rs2071307 variant and a tendency towards a genotypic ($p=0.075$; $p=0.112$) association
198 between the CON and ACL groups for the *FBN2* rs331079 and *ELN* rs2071307 variants
199 respectively. There were no genotypic or allelic associations between the CON and NON

200 subgroup. Furthermore, these gene variants did not show any significant distribution
201 difference when participants were grouped into genders (data not shown).

202 **Discussion**

203 We have shown that the *FBN2* rs331079 variant is significantly associated with the risk of
204 both AT and ACL rupture. Specifically, the GG genotype was over-represented in
205 participants with chronic AT and the G allele was over-represented in both pathologies.
206 Therefore, it appears that individuals carrying the G allele or the GG genotype are
207 approximately twice as likely to develop either of the two injuries. Interestingly this same
208 variant has recently been shown to associate with intracranial aneurysms in a Dutch
209 population [32]. However, in the Dutch study it was the C allele that was found to be the risk
210 factor as opposed to the G allele. It is noteworthy that *FBN2* mRNA levels have been shown
211 to be elevated in rat Achilles tendon undergoing repair with expression of *FBN2* reported to
212 be increased for ten days post injury [13]. Similarly, an increase in the expression of *FBN2*
213 has been found in other pathologies such as mitral valve prolapse [27].

214 ELN and FBN-2 are known to form a network of microfibrils that maintains the tendon
215 architecture [31]. An increase in FBN-2 levels might be expected to increase the density of
216 the tendon and lead to an increase in tendon stiffness and rigidity possibly affecting the
217 compliance of the tendon to muscle movement [4]. On the other hand, a decrease in FBN-2
218 levels could result in weaker tendons caused by structural deficiencies in the microfibril
219 network [30]. Impairment of the function of FBN-2 is believed to be a major determinant of
220 microfibrilopathy [30] which is speculated to precede a tendinopathy. Furthermore, the
221 increase in *FBN2* expression levels observed during tendon repair [13] is consistent with an
222 important role for FBN-2 in maintaining the tendon's architectural integrity.

223 Mutations such as the G3532T and G3590A substitutions have been found within the *FBN2*
224 gene that lead to the development of connective tissue disorders such as congenital
225 contractural arachnodactyly [9]. The rs331079 variant that we investigated in this study

226 resides within an intronic region of the *FBN2* gene (University of Florida.
227 www.snpper.chip.org). Although intronic variants do not determine the primary sequence of
228 a protein molecule [1], they may have other, hitherto, undiscovered roles that are necessary
229 for appropriate expression of protein molecules. However, at present the functionality of this
230 variant has not been described and therefore we do not know why it predisposes individuals
231 to AT and ACL rupture. The rs331079 variant is known to be part of a linkage block in
232 Caucasians and is in high linkage disequilibrium ($D'=1$) with the *FBN2* rs331081, rs331082,
233 and rs331085 variants (Wellcome Trust Sanger Institute. www.ensembl.com). All three of
234 these additional variants are also located within intron 7 of the *FBN2* gene (University of
235 Florida. www.snpper.chip.org). The linkage disequilibrium between the rs331079 variant that
236 we investigated and rs331081, rs331082, and rs331085 means that it is conceivable that one
237 of these linked variants may also have a role in predisposing to AT or ACL.

238 Our data do not support an association between the *ELN* rs2071307 variant and either AT or
239 ACL ruptures. It is interesting to note however, that although we found no relationship
240 between this variant and either pathology; the rs2071307 SNP is a non-synonymous and
241 possibly deleterious polymorphism (Queen's University. [http://compbio.cs.queensu.ca/F-](http://compbio.cs.queensu.ca/F-SNP/)
242 [SNP/](http://compbio.cs.queensu.ca/F-SNP/)) which results in a change of amino acid from hydrophobic glycine to hydrophilic serine
243 (University of Florida. www.snpper.chip.org). It is possible of course, that other variants
244 within this gene may be associated with either AT or ACL ruptures.

245 Although our study found a significant association between the *FBN2* rs331079 G allele and
246 the risk of AT and ACL rupture, the work has some limitations. Firstly, although our SA
247 cohorts (both TEN and ACL rupture groups) were matched for some aspects of physical
248 activity there were some differences in training behaviour and previous exposure to high
249 impact sports for the TEN cohort.. Secondly, we did not have detailed information on sports
250 history for the Australian cohort. Levels of physical activity should be accurately documented
251 in future studies. Furthermore, although the study was sufficiently powered to detect
252 associations with relatively large effects it should be repeated in bigger cohorts. Likewise,

253 additional association studies should be carried out in populations of different ethnicities
254 showing different minor allele frequencies for the rs331079 (African, 3%; European, 10%; ad-
255 mixed American, 28%; East Asian, 7%) and the rs2071307 (African, 26%; European, 39%;
256 ad-mixed American, 30%; East Asian, 14%) variants (1000 Genomes Project,
257 www.1000genomes.org).

258 Finally, the findings from this study advance our understanding of the polygenic basis of
259 musculoskeletal injuries. We suggest that the *FBN2* rs331079 variant should be considered
260 as an additional genetic locus to include in an injury risk assessment model that might be
261 used to identify athletes who are predisposed to AT and ACL ruptures.

262

263 **References**

- 264 1. *Berget S, Moore C, Sharp P.* Spliced Segments at 5' Terminus of Adenovirus 2 Late
265 Messenger-RNA. *Proc Natl Acad Sci U S A* 1977; 74: 3171-3175.
- 266 2. *Charbonneau NL, Ono RN, Corson GM, Keene DR, Sakai LY.* Fine tuning of growth factor
267 signals depends on fibrillin microfibril networks. *Birth Defects Res C Embryo Today* 2004; 72:
268 37-50.
- 269 3. *Collins M, Raleigh SM.* Genetic Risk Factors for Musculoskeletal Soft Tissue Injuries. *Med*
270 *Sport Sci* 2009; 54: 136-149.
- 271 4. *Cook C, McDonagh M.* Measurement of muscle and tendon stiffness in man. *Eur J Appl*
272 *Physiol Occup Physiol* 1996; 72: 380-382.
- 273 5. *El Khoury L, Posthumus M, Collins M, Handley CJ, Cook J, Raleigh SM.* Polymorphic
274 variation within the *ADAMTS2, ADAMTS14, ADAMTS5, ADAM12* and *TIMP2* genes and the
275 risk of Achilles tendon pathology: A genetic association study. *J Sci Med Sport* 2013; 16:
276 493-498.
- 277 6. *Ellis SG, Dushman-Ellis S, Luke MM, Murugesan G, Kottke-Marchant K, Ellis GM, Griffin*
278 *B, Tuzcu EM, Hazen S.* Pilot Candidate Gene Analysis of Patients ≥ 60 Years Old With Aortic
279 Stenosis Involving a Tricuspid Aortic Valve. *Am J Cardiol* 2012; 110: 88-92.
- 280 7. *Gianotti SM, Marshall SW, Hume PA, Bunt L.* Incidence of anterior cruciate ligament injury
281 and other knee ligament injuries: A national population-based study. *J Sci Med Sport* 2009;
282 12: 622-627.
- 283 8. *Gosline J, Lillie M, Carrington E, Guerette P, Ortlepp C, Savage K.* Elastic proteins:
284 biological roles and mechanical properties. *Philos Trans R Soc Lond B Biol Sci* 2002; 357:
285 121-132.

- 286 9. *Gupta PA, Putnam EA, Carmical SG, Kaitila I, Steinmann B, Child A, Danesino C,*
287 *Metcalf K, Berry SA, Chen E, Delorme CV, Thong M, Adès LC, Milewicz DM.* Ten novel
288 *FBN2* mutations in congenital contractural arachnodactyly: Delineation of the molecular
289 pathogenesis and clinical phenotype. *Hum Mutat* 2002; 19: 39-48.
- 290 10. *Harriss D, Atkinson G.* Ethical Standards in Sport and Exercise Science Research: 2014
291 Update. *Int J Sports Med* 2013; 34: 1025-1028.
- 292 11. *Harriss D, Atkinson G.* Update-ethical standards in sport and exercise science research.
293 *Int J Sports Med* 2011; 32: 819-821.
- 294 12. *Jarvinen TAH, Kannus P, Maffulli N, Khan KM.* Achilles tendon disorders: etiology and
295 epidemiology. *Foot Ankle Clin* 2005; 10: 255-66.
- 296 13. *Jelinsky SA, Li L, Ellis D, Archambault J, Li J, St Andre M, Morris C, Seeherman H.*
297 Treatment with rhBMP12 or rhBMP13 increase the rate and the quality of rat Achilles tendon
298 repair. *J Orthop Res* 2011; 29: 1604-1612.
- 299 14. *Lahiri D, Nurnberger J.* A Rapid Nonenzymatic Method for the Preparation of HMW DNA
300 from Blood for RFLP Studies. *Nucleic Acids Res* 1991; 19: 5444-5444.
- 301 15. *Ljungqvist A, Schwellnus MP, Bachl N, Collins M, Cook J, Khan KM, Maffulli N, Pitsiladis*
302 *Y, Riley G, Golspink G, Venter D, Derman EW, Engebretsen L, Volpi P.* International
303 Olympic Committee Consensus Statement: Molecular Basis of Connective Tissue and
304 Muscle Injuries in Sport. *Clin Sports Med* 2008; 27: 231-239.
- 305 16. *Mokone GG, Schwellnus MP, Noakes TD, Collins M.* The *COL5A1* gene and Achilles
306 tendon pathology. *Scand J Med Sci Sports* 2006; 16: 19-26.

- 307 17. Mokone GG, Gajjar M, September AV, Schwellnus MP, Greenberg J, Noakes TD, Collins
308 M. The Guanine-Thymine Dinucleotide Repeat Polymorphism Within the Tenascin-C Gene Is
309 Associated With Achilles Tendon Injuries. *Am J Sports Med* 2005; 33: 1016-1021.
- 310 18. Ng PC, Henikoff S. SIFT: predicting amino acid changes that affect protein function.
311 *Nucleic Acids Res* 2003; 31: 3812-3814.
- 312 19. Parkkari J, Pasanen K, Mattila VM, Kannus P, Rimpelä A. The risk for a cruciate ligament
313 injury of the knee in adolescents and young adults: a population-based cohort study of
314 46 500 people with a 9 year follow-up. *Brit J Sport Med* 2008; 42: 422-426.
- 315 20. Perneger TV. What's wrong with Bonferroni adjustments. *BMJ* 1998; 316: 1236.
- 316 21. Posthumus M, Collins M, van der Merwe L, O'Cuinneagain D, van der Merwe W, Ribbans
317 WJ, Schwellnus MP, Raleigh SM. Matrix metalloproteinase genes on chromosome 11q22
318 and the risk of anterior cruciate ligament (ACL) rupture. *Scand J Med Sci Sports* 2011: 321.
- 319 22. Posthumus M, September AV, Keegan M, Cuinneagain D, Van der Merwe W,
320 Schwellnus MP, Collins M. Genetic risk factors for anterior cruciate ligament ruptures:
321 *COL1A1* gene variant. *Brit J Sport Med* 2009; 43: 352-356.
- 322 23. Posthumus M, Collins M, Cook J, Handley CJ, Ribbans WJ, Smith RKW, Schwellnus MP,
323 Raleigh SM. Components of the transforming growth factor- β family and the pathogenesis of
324 human Achilles tendon pathology—a genetic association study. *Rheumatology* 2010; 49:
325 2090-2097.
- 326 24. Posthumus M, September AV, O'Cuinneagain D, van der Merwe W, Schwellnus MP,
327 Collins M. The association between the *COL12A1* gene and anterior cruciate ligament
328 ruptures. *Brit J Sport Med* 2009; 44: 1160-1165.

- 329 25. *Posthumus M, September AV, O'Cuinneagain D, van der Merwe W, Schwellnus MP,*
330 *Collins M.* The COL5A1 Gene Is Associated With Increased Risk of Anterior Cruciate
331 Ligament Ruptures in Female Participants. *Am J Sport Med* 2009; 37: 2234-2240.
- 332 26. *Puddu G, Ippolito E, Postacchini F.* A classification of Achilles tendon disease. *Am J*
333 *Sport Med* 1976; 4: 145-50.
- 334 27. *Radermecker MA, Limet R, Lapiere CM, Nusgens B.* Increased mRNA expression of
335 decorin in the prolapsing posterior leaflet of the mitral valve. *Interact CardioVasc Thorac*
336 *Surg* 2003; 2: 389-394.
- 337 28. *Raleigh SM, Posthumus M, O'Cuinneagain D, van der Merwe W, Collins M.* The GDF5
338 Gene and Anterior Cruciate Ligament Rupture. *Int J Sports Med* 2013; 34: 364-367.
- 339 29. *Raleigh SM, van der Merwe L, Ribbans WJ, Smith RKW, Schwellnus MP, Collins M.*
340 Variants within the MMP3 gene are associated with Achilles tendinopathy: possible
341 interaction with the COL5A1 gene. *Br J Sports Med* 2009; 43: 514-520.
- 342 30. *Robinson PN, Godfrey M.* The molecular genetics of Marfan syndrome and related
343 microfibrillopathies. *J Med Genet* 2000; 37: 9-25.
- 344 31. *Rosenbloom J, Abrams WR, Mecham R.* Extracellular matrix 4: the elastic fiber. *FASEB J*
345 1993; 7: 1208-1218.
- 346 32. *Ruigrok YM, Rinkel GJE, van't Slot R, Wolfs M, Tang S, Wijmenga C.* Evidence in favor
347 of the contribution of genes involved in the maintenance of the extracellular matrix of the
348 arterial wall to the development of intracranial aneurysms. *Hum Mol Genet* 2006; 15: 3361-
349 3368.

350 33. Saracini C, Bolli P, Sticchi E, Pratesi G, Pulli R, Sofi F, Pratesi C, Gensini GF, Abbate R,
351 Giusti B. Polymorphisms of genes involved in extracellular matrix remodeling and abdominal
352 aortic aneurysm. J Vasc Surg 2012; 55: 171-179.e2.

353 34. September AV, Cook J, Handley CJ, van der Merwe L, Schwellnus MP, Collins M.
354 Variants within the COL5A1 gene are associated with Achilles tendinopathy in two
355 populations. Br J Sports Med 2009; 43: 357-365.

356 35. Zhang H, Hu W, Ramirez F. Developmental expression of fibrillin genes suggests
357 heterogeneity of extracellular microfibrils. J Cell Biol 1995; 129: 1165-1176.

358

359 **List of legends**

360 **Table 1:** General characteristics of the Achilles tendinopathy group (TEN), the anterior
361 cruciate ligament rupture group (ACL), and the ACL subgroup with the non-contact (NON)
362 mechanism of injury as well as their respective control groups.

363 **Table 2:** The genotype and allele frequency distribution of the two selected candidate
364 variants within the Achilles tendinopathy (TEN), ACL ruptures (ACL) and their respective
365 asymptomatic control (CON) groups.

366

367

368

369