Investigating A Dose Response Relationship between High Fat Diet Consumption and the Contractile Performance of Isolated Mouse Soleus, EDL and Diaphragm Muscles

Hurst, J., James, R., Cox, V., Hill, C. & Tallis, J.

Author post-print (accepted) deposited by Coventry University's Repository

Original citation & hyperlink:

Hurst, J, James, R, Cox, V, Hill, C & Tallis, J 2018, 'Investigating A Dose Response Relationship between High Fat Diet Consumption and the Contractile Performance of Isolated Mouse Soleus, EDL and Diaphragm Muscles' European Journal of Applied Physiology, vol. 119, pp. 213-226. https://dx.doi.org/10.1007/s00421-018-4017-6

DOI 10.1007/s00421-018-4017-6 ISSN 1439-6319 ESSN 1439-6327

Publisher: Springer

The final publication is available at Springer via http://dx.doi.org/10.1007/s00421-018-4017-6

Copyright © and Moral Rights are retained by the author(s) and/ or other copyright owners. A copy can be downloaded for personal non-commercial research or study, without prior permission or charge. This item cannot be reproduced or quoted extensively from without first obtaining permission in writing from the copyright holder(s). The content must not be changed in any way or sold commercially in any format or medium without the formal permission of the copyright holders.

This document is the author's post-print version, incorporating any revisions agreed during the peer-review process. Some differences between the published version and this version may remain and you are advised to consult the published version if you wish to cite from it.

1	INVESTIGATING A DOSE-RESPONSE RELATIONSHIP BETWEEN HIGH-FAT DIET
2	CONSUMPTION AND THE CONTRACTILE PERFORMANCE OF ISOLATED MOUSE SOLEUS,
3	EDL AND DIAPHRAGM MUSCLES
4	
5	
6	Short title: EFFECT OF FATNESS ON MOUSE MUSCLE FUNCTION
7	
8	
9	Josh Hurst ¹ (\boxtimes), Rob S. James ¹ , Val M. Cox ¹ , Cameron Hill ¹ , Jason Tallis ¹
10	
11	¹ Centre for Sport, Exercise and Life Sciences, Science and Health Building, Coventry University, Priory
12	Street, Coventry CV1 5FB, UK
13	
14	
15	⊠: Hurstj5@uni.coventry.ac.uk; Phone Number: 024 7765 8562
16	
17	
18	
19	
20	
21	
22	
23	
24	
25	
26	
27	
28	
29	
30 31	
32	
33	
34	
35	
36	
37	
38	
39	
40	

- 41 Abstract

44	performance. The extent and magnitude of these changes in relation to total dose high fat diet consumption
45	remains unclear. This study aimed to examine the dose response relationship between a high fat diet and isolated
46	skeletal muscle contractility.
47	Methods: 120 female CD1 mice were randomly assigned to either control group or groups receiving 2, 4, 8 or
48	12-weeks of a high calorie diet (N=24). At 20 weeks soleus, EDL or diaphragm muscle was isolated (n=8 in each
49	case) and isometric force, work loop power output and fatigue resistance were measured.
50	Results: When analysed with respect to feeding duration, there was no effect of diet on the measured parameters
51	prior to 8 weeks of feeding. Compared to controls, 8-weeks feeding caused a reduction in normalised power of
52	the soleus and 8 and 12 weeks feeding caused reduced normalised isometric force, power and fatigue resistance
53	of the EDL. Diaphragm from the 12-week group produced lower normalised power, whereas 8 and 12-week
54	groups produced significantly lower normalised isometric force. Correlation statistics indicated that body fat
55	accumulation and decline in contractility will be specific to the individual and independent of the feeding duration.
56	Conclusion: The data indicate that a high fat diet causes a decline in muscle quality with specific contractile
57	parameters being affected in each muscle. We also uniquely demonstrate that the amount of fat gain, irrespective
58	of feeding duration, may be the main factor in reducing contractile performance.
59	
60	Key Words: Force; Muscle Quality; Muscular Lipid; Lipid Accumulation; Power.
61	
62	Abbreviations: ANOVA-Analysis of variance; CF- Cycle frequency; EDL-Extensor digitorum longus; HFD-High
63	fat diet; PO- power output; WL – work loop.
64	
65	
66	
67	
68	
69	
70	
71	
72	
73	
74	
75	
76	
77	
/8 70	
/9 80	
00	

Purpose: Recent evidence has demonstrated an obesity induced, skeletal muscle specific, reduction in contractile

81 Introduction

- Recent data has indicated that 1.4 billion adults (approximately 19% of the world's population) over the age of
- 18 are either overweight or obese (WHO. 2017), which is disproportionally higher in adults residing in westernised
- nations (WOF. 2017). The increased prevalence of obesity has been closely linked with an increase in diabetes,
- cardiovascular disease, cancer and other potentially life-threatening illnesses and diseases (Kopelman. 2007).
- Furthermore, there is an emerging body of evidence suggesting that obesity significantly affects the contractile performance of skeletal muscle (Ciapaite *et al.* 2015; Tallis *et al.* 2017b; Tomlinson *et al.* 2016; Tuttle *et al.* 2012).
- 688 Given that skeletal muscle is the primary site of substrate metabolism in the body (Zurlo *et al.* 1990), and that
- 89 contractile function is needed to fulfil activities of daily living and increase calorie expenditure, it has been
- proposed that the effects of obesity on skeletal muscle may be a catalyst for obesity related disease (Tallis *et al.*
- 91 2014).
- 92

93 Human in vivo studies that have examined the effects of obesity on muscle performance have demonstrated an 94 improvement in the absolute force producing capacity of postural and locomotor muscles (Abdelmoula et al. 2012; 95 Maffiuletti et al. 2007; Szymura et al. 2011), but little effect on the contractile performance of other muscles 96 (Capodaglio et al. 2009; Hulens et al. 2001). The improved performance in postural and locomotor muscles has 97 been proposed to relate to a positive training adaptation induced by the elevated body mass of obese individuals. 98 When contractile function is reported as a function of body mass, in humans, muscle function is significantly 99 reduced in obese individuals in both sexes and across a range of ages (Aucouturier et al. 2007; Miyatake et al. 100 2000; Rolland et al. 2004; Szymura et al. 2011; Ward et al. 1997).

101

102 A recent review examining the effect of obesity on skeletal muscle function (Tallis et al, 2018) outlines the value 103 of using isolated muscle to assess the effects of obesity on contractility. Although in vivo work has offered a 104 valuable insight into the effects of obesity on skeletal muscle function, a more comprehensive understanding can 105 be gained by also undertaking studies using isolated skeletal muscle. It has been suggested that human studies do 106 not make accurate assessments of obesity associated changes in muscle quality (muscle performance normalised 107 to muscle size) (Tallis et al. 2017b). Understanding obesity induced changes in muscle quality is important, as 108 producing the highest contractile performance per quantity of tissue reduces the muscle mass required, hence 109 lowering body mass, and decreases the cost of maintaining that muscle mass (Tallis et al. 2017b). It is common 110 for in vivo research to report absolute changes in contractile performance, or that normalised to body mass (Abdelmoula et al. 2012; Miyatake et al. 2000; Paolillo et al. 2012; Ward et al. 1997) which provide little insight 111 112 into obesity induced changes in muscle quality. Only a very small number of in vivo studies have used techniques 113 such as Magnetic Resonance Imaging or Computed Tomography scans in an attempt to assess muslce quality 114 (Blimkie et al. 1989), however these only provide estimates of muscle size which has led to ambiguity with respect 115 to the effect of obesity on the intrinsic force producing capacity of the contractile proteins. Isolated skeletal muscle 116 studies allow for a more accurate assessment of muscle quality as the whole muscle mass can be determined, and 117 the performance of one particular muscle is measured. Furthermore, an isolated skeletal muscle approach allows 118 assessment of the fibre type and muscle specific effect of obesity on contractile performance without the 119 confounding effects of the central nervous system, including central inhibition. Previous work has stated that the 120 effect of obesity on skeletal muscle fatigue cannot be accurately measured in vivo as the muscle of the obese

individual will be working at a greater intensity to overcome the increased body inertia (Tallis *et al.* 2017b). As
such, it has been proposed that assessment of the fatigue resistance of isolated skeletal muscle allows a true

- 123 examination of the effects of obesity on skeletal muscle performance.
- 124

125 To date, only a small number of studies have assessed the effect of obesity on isolated skeletal muscle contractility 126 with the majority of these studies using rodents (Bott et al. 2017; Ciapaite et al. 2015; Eshima et al. 2017). Results 127 demonstrate that the effect of obesity is muscle and contractile parameter specific. There is evidence to suggest 128 that the absolute force and power producing capacity of skeletal muscle is either increased or maintained, whereas 129 there is an obesity induced reduction in muscle quality, meaning that obesity may result in a reduction in the 130 intrinsic force producing capacity of the contractile proteins (Tallis et al. 2017b). Tallis et al (2018), suggested 131 that the ambiguity in findings is likely related to methodological approach (i.e. mechanical assessment method 132 and differences in the temperature at which the experiments were performed), or feeding duration, with previous 133 work administering HFD for durations between 3 and 16 weeks (Ciapaite et al. 2015; Tallis et al. 2017b; Thomas 134 et al. 2014). Although this has not been thoroughly investigated, several proposed mechanisms, such as a reduction 135 in protein synthesis (Akhmedov and Berdeaux. 2013; Anderson et al. 2008), changes in the expression of proteins 136 involved with calcium handling (Bruton et al. 2002; Warmington et al. 2000), fibre type shift (Eshima et al. 2017; 137 Shortreed et al. 2009) and an altered metabolic profile (Ciapaite et al. 2015; Seebacher et al. 2017; Tallis et al. 138 2017b) have been suggested as potential mechanisms for the obesity associated reduction in muscle performance. 139 The onset and magnitude of changes in muscle performance seems likely to be muscle specific and to vary with 140 feeding duration.

141

The present study used the work loop technique to uniquely examine the effects of 2, 4, 8 or 12 weeks of HFD 142 143 feeding on the mechanical performance of isolated soleus, extensor digitorum longus (EDL) and diaphragm 144 muscle to provide an insight into the muscle specific onset and magnitude of the obesity response. This has been 145 prioritised as an important area for future work in a recent review by Tallis et al (2018). The range of feeding 146 durations used in this study was selected as previous work has shown mechanistic changes, such as altered fibre 147 type composition, from as little as 3 weeks of a HFD consumption (Thomas et al. 2014). Results gained from this 148 study will provide clarity on the magnitude of the effects of a HFD over different durations of feeding and will help rationalise previous ambiguous finding untimely providing a better understanding of the effects of obesity 149 on isolated skeletal muscle function. Furthermore, previous studies examining the obesity effect on isolated 150 151 muscle performance have primarily used assessments of isometric force (Bott et al. 2017; Ciapaite et al. 2015; 152 Eshima et al. 2017; Matsakas et al. 2015; Shortreed et al. 2009; Thomas et al. 2014; Warmington et al. 2000) and 153 to date only two studies have used the work loop technique to examine obesity effect on muscular power 154 (Seebacher et al. 2017; Tallis et al. 2017b). Studies assessing isometric muscle force are limited with respect to 155 their in vivo application given that dynamic power producing muscle activity is needed for the completion of 156 locomotor tasks (Josephson, 1985). Although isotonic and isovelocity assessments of muscle function have been 157 used to assed isolated skeletal muscle power in previous work (Barclay, 1996; James et al, 1996), such techniques 158 have been criticised and suggested to substantially overestimate muscle power output (James et al, 1996). The 159 work loop techniques considers muscle force generation during dynamic activity and accounts for the interaction 160 between force produced during shortening, passive resistance to re-lengthening and changes in activation and

(Josephson, 1985; James *et al*, 1995). *In vivo*, locomotor muscles are used to produce and absorb work during shortening and lengthening, meaning examination of obesity effects on muscle performance using the work loop technique provides a valuable insight into the effects of obesity on a contraction type fundamental for tasks of daily living previous work has only examined the effect of obesity on work loop power at a fixed cycle frequencies (number of length changes per unit time) (Tallis *et al* .2017b; Seebacher *et al*. 2017). Here we uniquely measure obesity effects on work loop power over a range of cycle frequencies in order to gain a better understanding of obesity effects on power across and range of length change velocities. This offers a greater insight on how obesity

relaxation time using length change magnitudes and waveform that more closely replicate those used in vivo

- 169 effects range of length change velocities used during locomotor tasks *in vivo*.
- 170 171

172

161

Results

Morphology

173 Whole animal body mass, absolute gonadal fat pad mass, gonadal fat pad mass as a percentage of whole body 174 mass and abdominal circumference were significantly affected by treatment (Table 2; ANOVA P<0.004 in each 175 case). More specifically 8 and 12 weeks of feeding resulted in significantly greater body mass, absolute gonadal 176 fat pad mass, gonadal fat pad mass as a percentage of body mass and abdominal circumference compared to lean 177 controls (Dunnett's P<0.01). Muscle mass for soleus was significantly greater in treatments groups of 8 and 12 178 weeks of feeding when compared to lean controls (Table 2, ANOVA P<0.001, Dunnett's P<0.001 in both cases). 179 Muscle mass of the EDL was significantly greater in treatment groups of 4, 8 and 12 weeks of feeding when compared to lean controls (Table 2, ANOVA P<0.01, Dunnett's P>0.01). Diaphragm mass is not compared 180 181 between treatment groups as it was not possible to consistently remove the exact same section of the diaphragm 182 on each occasion

- 183
- 184

Isometric Tetanus Force & Stress

For the soleus and the EDL, maximal isometric stress and force was not significantly affected by treatment (Figure 1A, 1B, 1C, 1D; ANOVA P>0.1 in each case). The maximal isometric stress of diaphragm was significantly lower than lean controls in the 8-week HFD treatment group (Figure 1E ANOVA P=0.01; Dunnett's P=0.04).

188

Time to half peak tetanus (THPT) was not significantly different between the treatment groups for soleus, EDL or diaphragm (ANOVA P>0.5 in each case). Similarly, time from the last stimulus to half relaxation (LSHR) for EDL and diaphragm was not significantly affected (Table 3 ANOVA P=0.7). However, LSHR for soleus was significantly affected by feeding duration (Table 3, ANOVA P=0.01, Dunnett's P>0.04), with LSHR being significantly greater in the 8 and 12 week feeding groups

194 195

Work loop power, fatigue and recovery.

There was no significant interaction between treatment and the optimal cycle frequency CF of work loop WL (two-way ANOVA P=1.00) on any of the muscles tested, the shape of the force velocity curve remained unchanged. For both the soleus and EDL, absolute work loop power output was not significantly different in any of the treatment groups when compared to the lean control (Figure 2B & D ANOVA P>0.13 in both muscles). When work loop power output was normalised to muscle mass: treatment group of 8 weeks of HFD for the soleus

- ANOVA P<0.019, Dunnett's P=0.03 & P=0.06 respectively). For the EDL, treatment groups of 8 and 12 weeks
- of HFD were significantly lower than the lean control group (Figure 2C, ANOVA, P<0.01, Dunnett's P>0.03).
- For the diaphragm, 12 weeks of a HFD resulted in significantly lower normalised work loop power output than
- 205 the lean control group (Figure 2E, ANOVA P=0.01, Dunnett's P=0.019). Absolute work loop power output for
- the diaphragm was not recorded as only a section of the whole muscle was taken.
- 207

There was an overall significant difference in time to fatigue for soleus (Figure 3A, ANOVA P=0.02), however there was no differences among treatments groups and the lean control group in post hoc analysis. EDL from treatment groups 8 and 12 fatigued significantly quicker than the lean control group (Figure 3B, ANOVA, P=0.02, Dunnett's P<0.03). Diaphragm had no significant changes in time to fatigue between any of the treatment groups and the lean control group (Figure 3C, ANOVA P>0.36). The ability for the muscle to recover over a 30-minute period post fatigue protocol was not significantly affected by any treatment for all the muscles examined (ANOVA, P>0.06).

215

216 Morphological data indicated that body fat accumulation is specific to the individual (table 4) and independent of 217 the feeding duration. For example, one individual from the 2-week feeding group had a body mass of 40.96g with 218 a FPM of 3.86g in comparison to another individual from the 12-week feeding group which had a body mass of 219 30.78g with a FPM of 0.5g. As such, correlations between gonadal fat pad mass and contractility measures were 220 analysed. For soleus, there was a significant positive correlation between body mass and gonadal fat pad mass, 221 body mass and absolute force (Table 4, R>0.42; P<0.001), and a significant negative relationship between body 222 mass and normalised work loop power output (R=-0.3; P=0.045). For EDL, there was a significant positive 223 relationship between body mass and absolute work loop power output (Table 4 R=0.385; P=0.011) and a 224 significant negative relationship between body mass and time to fatigue (Table 4 R=-0.323; P=0.035). For the diaphragm, there was a significant correlation between body mass and gonadal fat pad mass (R=0.613; P<0.001) 225 226 and a significant negative relationship between body mass and maximal tetanus stress and time to fatigue (Table 227 4. R=-0.339; P=0.025).

228 229

Discussion

230 The present study is the first to assess the effect of a range of high fat feeding durations on the contractile 231 performance of skeletal muscle using a method which more closely replicates in vivo muscle function. When 232 considered with respect to feeding duration, the results indicated that HFD consumption for a period shorter than 233 8 weeks has no significant effect on animal morphology or muscle isometric performance, results that agree with 234 previous findings showing short durations (3-8 weeks) of feeding have little effect on muscle contractility (Eshima 235 et al. 2017; Shortreed et al. 2009; Thomas et al. 2014). Only after 8 weeks of a HFD consumption affected 236 contractile performance, with this duration being the first associated with a significant increase in whole body 237 mass, gonadal fat pad mass (Table 2) and gonadal fat pad mass as a percentage of body mass (Table 2). These 238 data therefore indicate that changes in skeletal muscle performance are related to changes in body composition. 239 When gonadal fat pad mass was correlated with skeletal muscle mechanical performance variables and considered 240 irrespective of feeding duration, it was evident that for some individuals changes in contractile performance

ensued prior to 8 weeks HFD consumption. As such, the present findings infer that quantity of fat accumulation rather than feeding duration is the key factor driving obesity associated changes in muscle contractility and rationalising previous ambiguous findings in this area of research where studies report outcomes following fixed feeding durations.

- 245
- 246

Obesity effects on force and power

247 There were no changes in peak isometric force or peak isometric stress (force/cross section area) of SOL in any of treatment groups when compared to the lean control group. This is contradictory to the findings of previous 248 249 work which has demonstrated obese individuals have increased absolute force of postural muscles (Tallis et al. 250 2017b). In human studies, this has been attributed to an added training effect from increased body mass 251 (Maffiuletti et al. 2013). One possible reason for the discrepancy in our results is the body mass of the animals 252 used; body masses of the 8 and 12-week treatment groups were 41.2±7.1 and 38.7±6 respectively. However, the 253 body mass of the obese animals in the work by Tallis et al (2017b), was 52.7±2.30, with this additional body mass 254 likely from the longer duration of feeding the animal received (16 weeks). A greater body mass is likely to evoke 255 a greater training stimulus, given the higher force requirement needed to overcome body inertia. Absolute power 256 output was also unaffected by treatment in the present study, however when normalised to muscle mass, power 257 output from the 8-week HFD group was significantly less than their lean counterparts and was approaching 258 significance (P=0.06) in the 12-week HFD group. This reduction in the normalised power of soleus was not 259 evident in our previous work using a longer feeding duration (Tallis et al. 2017b), further exacerbating the complexity with respect to the effect of feeding duration and fat gain has on muscle performance. 260

261

262 When compared to controls, isometric stress, absolute force or absolute power output in the EDL or soleus was 263 unchanged in the HFD groups. Shortreed et al (2009), also reported no force changes in the EDL after 8 weeks of 264 feeding. Previous literature has reported a decrease in both tetanic force and stress (Eshima et al. 2017; Matsakas 265 et al. 2015), although variation in methodology (both age and assessment temperature) may account for these discrepancies. In the EDL normalised power output was significantly reduced in animals from HFD treatment 266 groups of 8 and 12-week duration, indicating that the increase in muscle mass induced by obesity was not 267 268 associated with an improvement in contractile performance. These data also demonstrate that the reduction in muscle quality may be reduced earlier than 8 weeks and that the magnitude of the decline is dependent on fat 269 270 accumulation rather than feeding duration (see table 4). As such, findings from the present study infer that an 271 obesity associated reduction in muscle quality is likely to occur following much shorter feeding durations than 272 those previously cited (Tallis et al. 2017b). When considered in vivo, larger muscles of poorer quality will be 273 produced, raising metabolic demand in protein synthesis and further increasing body mass and body inertia.

274

275 Our results infer that the magnitude and onset of the obesity associated changes in contractile performance are

276 muscle specific, and likely related to mechanical function, anatomical location and fibre type distribution. These

277 findings are the first to indicate that obesity may cause a reduction in skeletal muscle quality across all muscles,

- which has not been the case in previous work. For example, Tallis *et al* (2017b), demonstrated that muscle quality
- was relatively well maintained in the soleus, which is not the case in the present findings. Such results further
- 280 highlight the complexity and ambiguity in relation to this area of research. However, we propose that the changes

on body composition, or more specifically fatness, are the most important mediator of obesity associated changes in muscle performance. Given the rate of fat accumulation is not uniform across all individuals, assessing the effects of obesity using HFD during a fixed feeding duration and failing to consider individual results may be the cause of ambiguity in recent findings.

285

286 Correlation data (table 4) between morphological and contractility measurements plotted against fat pad mass 287 support the theory that fatness rather than feeding duration is the main influence on the decline on muscle 288 performance.

289 290

Obesity effects on fatigue resistance

291 The effect of obesity on fatigue resistance of skeletal muscle has been previously assessed in vivo (Maffiuletti et 292 al. 2007). However, in vivo work does not give an accurate assessment of skeletal muscle fatigue as muscle from 293 an obese individual will have to move a bigger mass resulting in faster fatigue. Only one study has previously 294 assessed the effect of obesity on the fatigue of muscle power output. Tallis et al (2017b), reported that soleus 295 fatigued significantly quicker in the obese group when compared to the lean controls, given the postural support 296 role of the Sol this would have detrimental effects on sustained locomotion. Results from the current study showed 297 little change in time to fatigue for the soleus and no change in the diaphragm. Contrary to Tallis et al (2017b), the 298 present study reported a significant reduction in time to fatigue in the EDL in both 8 and 12-week treatment 299 groups, as a longer duration of feeding was used in the Tallis et al (2017b) study, it is possible the fiber type of 300 the EDL began to take on the metabolic formula of the adjacent fibers in preparation to alter fiber type in 301 accordance with the nearest neighbor theory (Pette and Staron. 1997). As per previous work (Tallis et al. 2017b) 302 it would be expected that when such muscle performance is considered *in vivo*, that all muscles would experience 303 greater fatigue given the obesity induced reduction in normalized work loop power and the elevated segmental 304 mass. This point is particularly important when considering the *in vivo* role of the diaphragm; reduced fatigue 305 resistance could lead to reduced oxygen delivery, deteriorating fatigability for all muscles. Although significant 306 differences were shown in the time to fatigue in the EDL, correlation data (table 4) shows individuals from all 307 treatment groups can be found at both ends of the trend line, again suggesting that fatness, not duration of feeding 308 if the key underlying issue.

309 310

Mechanisms for an obesity associated change in skeletal muscle contractile performance

311 There are a number of mechanisms which have been suggested to affect the decline in muscle performance due 312 to obesity, such as a change in fiber type composition (Eshima et al. 2017; Shortreed et al. 2009), a reduction in 313 protein synthesis (Akhmedov and Berdeaux. 2013; Anderson et al. 2008), altered calcium handling (Bruton et al. 314 2002; Warmington et al. 2000) and changes to metabolic profile (Ciapaite et al. 2015; Seebacher et al. 2017; 315 Shortreed et al. 2009; Tallis et al. 2017b). The variation in mechanisms proposed for how obesity contributes to 316 a decline in muscular contractility may at least partially be due to variation in methodology used in previous 317 literature and the dearth of research which attempts to couple mechanistic changes with effects on contractile 318 performance. The changes in mechanisms relating to muscle function are likely to be muscle specific given that 319 the changes in performance are, there is also some relation to the magnitude of fatness. These mechanisms may

320 be further exacerbated by long duration feeding.

322 The findings regarding changes in fiber type expression as a result of obesity in rodents are equivocal, with some 323 literature reporting no change (de Wilde et al. 2008; Denies et al. 2014; Shortreed et al. 2009; Tallis et al. 2017b; 324 Trajcevski et al. 2013), and some reporting a shift to either slow twitch or fast twitch fibers (Eshima et al. 2017; 325 Kemp et al. 2009; Shortreed et al. 2009; Tanner et al. 2002; Thomas et al. 2014; Warmington et al. 2000). Part 326 of this ambiguity may be related to evidence derived from dietary induced obesity models compared genetically 327 obese rodent models. Generally, studies using OB/OB rodents show a shift to slower, more oxidative fibers 328 (Warmington et al. 2000) with a lower muscle mass and those using dietary induced obesity demonstrate a shift 329 towards faster fibers (Eshima et al. 2017; Thomas et al. 2014) and a larger muscle mass (Ciapaite et al. 2015; 330 Tallis et al. 2017b). Although fiber type composition and metabolic capacity was not analyzed, a decrease in slow 331 twitch fibers or altered AMPK activation may go some way in explaining the reduction in time to fatigue in obese individuals, however a reduction in contractile performance without a change in fiber type composition has also 332 333 been reported (Tallis et al. 2017b). Ciapaite et al (2015), reported a decrease in tetanus force in the soleus with 334 no change in MyHC, however an increase in slow Tnnt1 isoform and decrease in fast Tnnt3 isoform was reported. This change would increase Ca²⁺ sensitivity and decrease Ca²⁺ cooperativity of force production, therefore, 335 reducing force production of the muscle (Brotto et al. 2006). Changes in both fibre type and speed of actin-myosin 336 337 cycling need further investigation.

338

339 Obesity has been associated with -chronic inflammation leading to elevated pro-inflammatory cytokines TNF- α 340 and IL-6 which has been linked to a decline in skeletal muscle contractile protein synthesis (Akhmedov and 341 Berdeaux. 2013; D'Souza *et al.* 2015; Tallis *et al.* 2017b). The increase in muscle mass and decrease in normalised 342 muscle power output from the obese group in the current study suggest that although skeletal muscle remodeling 343 is continued in obese individuals, the quality of the contractile proteins being synthesized is likely to be reduced.

344 345

Broader applications of the findings

Elevated fatness has the potential to cause individuals to fall into a negative cycle, as body mass increases and muscle quality decreases, obese individuals may have their movement reduced due to earlier fatigue and reduced ability to produce muscular power. Obese individuals may also have reduced ability to oxidize fat due to metabolic changes, further adding to the negative cycle. This lack of movement will not only decrease quality of life, but also lead to further decreases in calorie expenditure which may lead to further increases in body mass. An obesity associated reduction in the performance of the diaphragm, may affect pulmonary function and oxygen delivery to working tissue thus further limiting muscle performance.

- 353
- This data has shown important results with respect of the effect of obesity on muscle function at the muscle level, but to consider effect on whole body functional performance, contractile function in relation to elevated body inertia should be considered. The effect of a reduction in muscle quality on locomotor performance has already been outlined. In addition, the limited effects on the absolute force and power of soleus and EDL muscles would also present substantial problems for functional performance *in vivo* given that such musculature is expected to cause movement in a system that has a greater inertia. As a result, the decline in fatigue resistance will also be on a greater magnitude given that musculature of the obese group would have to work at a greater intensity to sustain

- movement at the same speed as their lean counterparts. Importantly, this data adds growing evidence that indicates that the obesity associated reduction in locomotor performance is caused by factors further to elevated inertia induced by greater fat mass, but a significant reduction in the decline of the contractile performance of skeletal muscle. Furthermore, the effects seen at a muscle level may be further exaggerated as obesity has reportedly caused changes in neuromuscular recruitment (Yoshida *et al.* 2012).
- 366 367

Limitations and future work

368 One limitation for this study is normalising work loop power to whole muscle mass in attempt to assess muscle 369 quality. Goodpaster et al (2001), reported that skeletal lipid content can double in obese individuals; this would 370 mean a smaller proportion of the muscle mass would be contractile protein; therefor these results may overestimate 371 the decline in contractile performance. Normalising work loop power to lean muscle mass would allow for a more 372 accurate representation of muscle quality. Although obese individuals may have a higher intramuscular lipid 373 content than their lean counterparts, Machann et al (2003) reported this difference was limited with intramuscular 374 fat in the soleus rising from 2.5% in lean individuals to 3.8% in obese individuals. It would be possible to quantify 375 the volume of the contractile machinery in the muscle fibre using transmission electron microscopy (Singh et al, 376 2009). This would provide a more detailed assessment of muscle quality in an obese model.

377

The current study used gonadal fat pad mass as an indicative measure of whole animal body fat. Although this is a valuable measurement for determining differences in fatness between the experimental groups, evidence has indicated that mechanistic responses to obesity may be influenced by the specific location of lipid accumulation (Wronska *et al*, 2012). The effect of lipid accumulation location on skeletal muscle contractile performance is yet to be explored and should be a specific focus of future work. In particular, a focus on muscular lipid accumulation and skeletal muscle function should be preauthorised.

384

The current study, reports the effect of a range of short term feeding durations on muscle contractility, however, it would be valuable to look at the effects of longer durations of feeding which reflect a 'lifetime' of a high fat diet and more sustained obesity. This would also take into consideration both obesity and sarcopenic effects. Given that this study replicates the negative effects of a high fat diet on skeletal muscle contractility, reversibility of these effects should be investigated further. Both exercise and calorie restrictive diet models should be implemented on dietary induced obese animals.

391

As mentioned in a recent review article (Tallis *et al*, 2018), a key direction for future work should be to add to the dearth of littering examining the underpinning mechanisms which cause the decline in performance onset by obesity.

395 396

Conclusion

In summary, our findings offer a complex and novel insight into the effects of a HFD on skeletal muscle mechanics. The data suggests that although feeding duration may have some influence, it is the amount of fat gained during a HFD that affects muscle contractility. An increase in muscle mass in the soleus and EDL along with no change in absolute tetanic force or absolute power output shows a decrease in muscle quality as bigger 401 muscles are required to produce the same amount of force. Such effects would be exacerbated in vivo when considering the additional body mass gained during obesity, which could lead to a negative cycle of further weight 402 403 gain, reduced mobility and a reduction in quality of life. Our data suggest that it may be possible for some 404 individuals to see substantial negative changes at low durations of a HFD if considerable fat gain occurs, 405 compromising both function performance and health. The decline in muscular performance due to obesity may be 406 caused by a range of morphological and biochemical changes and future work should aim to correlate these 407 changes to better identify the underpinning mechanisms.

408

420

409 **Material and Methods**

410 Animals

411 Following ethics approval from Coventry University, 150 female CD1 mice (Charles River, UK; Harlan 412 Laboratories, UK) were randomly assigned to 5 different experimental groups - 4 treatment groups of differing feeding durations and 1 lean control group. The treatment groups were given 2, 4, 8 or 12 weeks of a high fat 413 414 forage diet. Mice were housed in a 12-hour light/dark cycle at 50% humidity. Animals were aged matched and kept in cages containing 8-10 individuals. All animals were given water and standard lab chow (SDS RM-1 415 416 Maintenance) ad libitum. Mice in the treatment groups also had ad libitum access to a HFD, in the form of a laboratory supplied forage diet (Advanced Protocol PicoLab Natural Sunflower). Nutritional information for both 417 418 the lab chow and the sunflower seeds are given in Table 1.

- 419 Mice where aged to 20 weeks of age, with the high fat diet delivered at the defined period prior to 20 weeks (e.g.
- 12 weeks of feeding began at 8 weeks of age). Once an experimental group was of age they were weighed and 421 sacrificed via cervical dislocation in accordance with British Home Office Animals (Scientific Procedures) Act
- 422 1986, Schedule 1. Gonadal fat pad mass was dissected and weighed as an indicative measure of total body fat
- 423 (Rodgers et al, 1980). Either the rib cage or hind limbs were removed and placed in chilled oxygenated (95% O₂;
- 5% CO₂) Krebs- Henseleit solution ([mM] NaCl 118; KCl 4.75; MgSO₄ 1.18; NaHCO₃ 24.8; KH₂PO₄ 1.18; 424
- glucose 10; CaCl₂ 2.54; pH 7.55 at room temperature [James et al. 2005]). The target muscle was then isolated 425
- 426 from either the right hind limb or the rib cage and dissected under microscope.

427 **Dissection and preparation**

- 428 The limb containing the target muscle (EDL or soleus) or the rib cage containing the whole, intact diaphragm 429 (n=8-10 per experimental group) was isolated and pinned out on an sylgard dish containing frequently changed, 430 chilled, oxygenated (95% O₂, 5%CO₂) Krebs-Henseleit solution. These muscles were used due to their different anatomical location, function and fibre type composition, as well as to replicate previous literature where these 431 432 muscles are commonly used for contractile performance studies. Once the muscle (whole Sol, whole EDL, section
- of the DIA) had been isolated, aluminium foil T clips were used to wrap around the tendons at the distal end of 433
- 434 the muscle whilst a small piece of bone was left at the proximal end of the muscle, these were placed into crocodile
- 435 clips to secure the muscle in the bath.

436 **Experimental set up**

- and stimulation parameters. Each muscle was placed in an organ bath and attached at one end to a force transducer
- 439 (UF1, Pioden Controls Ltd, UK) and at the other to a motor arm (V201, Ling Dynamic Systems, UK). Position of
- the motor arm was detected via a Linear Variable Displacement Transformer (LVDT, DFG5.0, Solartron
- 441 Metrology, UK). The motor arm was used to change the length of the muscle during assessments of work loop
- 443 a central reservoir maintained at a constant temperature by a heater/cooler (Grant LTD6G, Grant Instruments Ltd,

power. The muscle was immersed in circulated oxygenated Krebs solution $(37^{\circ}C \pm 0.2)$ which was pumped from

- 444 UK). A digital thermometer (Checktemp C, Harvard Apparatus, UK) was used inside the bath to monitor
- temperature throughout the duration of the experiment.
- 446 The muscle was stimulated to produce force via parallel platinum electrodes which were submerged in the Krebs
- solution inside the bath, the amplitude of the stimulation was controlled by an external power source (PL320,
- 448 Thurlby Thandar Instruments, Huntingdon, UK). Visual representation of the force and length data was provided
- 449 by a storage oscilloscope (2211, Tektronix, Marlow, UK). Length change and stimulation parameters were
- 450 controlled by a custom programme within Testpoint software (Testpoint, CEC, Massachusetts, USA).

451 Experimental testing of skeletal muscle

- 452 The procedure for assessing the contractile performance of isolated skeletal muscle aligns with that of previous work (James et al. 1996; Tallis et al. 2014; Tallis et al. 2017b). Once the muscle was successfully clamped in the 453 454 bath it was allowed 10 minutes to equilibrate to the new temperature. Following this, the resting length of the muscle and stimulation amplitude (typically 14-16V, 14-16V and 12-14V for soleus, EDL and diaphragm 455 456 respectively) was altered to attain maximal twitch force. Assessment of isometric tetanus force was then measured. Electrical stimulation was delivered at a fixed burst duration (350ms, 200ms and 250ms for soleus, EDL and 457 458 diaphragm respectively), and stimulation frequency (typically 120-140Hz, 200-220-Hz and 120-140Hz for soleus, 459 EDL and diaphragm respectively) was optimised to elicit maximal tetanus force. Each tetanus assessment was separated by a 5-minute recovery period. Measures of activation time (time to half peak tetanus; THPT) and 460 461 relaxation time (time from last stimulus to half tetanus relaxation; LSHR) were obtained from the peak isometric
- tetanus force trace.

- The muscle length that elicited maximal isometric force was measured by an eyepiece graticule fitted to a microscope and was defined as L_0 . As per previous work, mean muscle fiber length was calculated as 85% and 75% of L_0 for soleus and EDL muscle respectively (James *et al.* 1995). Due to the variances in each diaphragm dissection, no such estimation of muscle fibre length exists. Therefore, the exact diaphragm preparation length reading gained from the graticule was used as per previous studies (Tallis *et al.* 2014; Tallis *et al.* 2017a).
- 468 Post isometric optimisation and a 5-minute rest, the work loop technique was used. This technique assesses the
- 469 ability of the muscle to produce power whilst undergoing cyclical length changes (James et al. 2005; James et al.
- 470 1995; Josephson. 1985). During the work loop, optimal length and stimulation parameters determined during the
- 471 isometric assessments were used. Each muscle was subjected to 4 sinusoidal length changes at a symmetrical
- 472 strain of 0.10 (10% of L_0), lengthening the muscle from its original L_0 length by 5% and then shortening the
- 473 muscle to 5% less than L_0 , before returning to L_0 length. Cycle frequency, phase and burst duration were altered

to attain maximal net work loop power and to construct a power output- cycle frequency curve. Between each

475 work loop, a 5-minute rest period was used to ensure recovery (Tallis *et al.* 2012).

- 476 The cycle frequency determines the speed at which the muscle undergoes changes in length. Due to the differences
- 477 in fibre type composition, the different muscles tested produced maximal power at different length change
- velocities. Previous work demonstrates that the cycle frequency to elicit maximal work loop power is 5Hz, 7Hz
- and 10Hz for soleus, diaphragm and EDL respectively in healthy young female mice (Altringham and Young.
- 480 1991; James *et al.* 1995). Given that the optimal cycle frequency to elicit maximal power may be affected by 481 obesity, here we assessed work loop power over a range of cycle frequency (2, 3, 4, 5, 6, 7, 8 and 10 Hz for soleus;
- 482 4, 6, 8, 10, 12, 14 and 16 Hz for EDL; 3, 4, 5, 6, 7, 8, 10 and 12 Hz for diaphragm). This methodological approach
- 483 has been used in previous studies (James *et al.* 1996; James *et al.* 1995; Tallis *et al.* 2014; Tallis *et al.* 2017b). To
- 484 optimise work at each cycle frequency, strain was also altered. Typically, in order to obtain maximal work loop
- 485 power strain had to be reduced slightly at higher cycle frequency and increased slightly at lower cycle frequency.
- In order to optimise the net work produced during the work loop cycle, stimulation parameters were adjusted. 486 487 Burst duration equates to the number of electrical stimuli which the muscle receives during the work loop. The optimal burst duration is one that maximises net work during the whole work loop cycle. If the burst duration is 488 489 too short, then maximal work will not be achieved due to limited Ca²⁺ release limiting the duration of force production during shortening. If the burst duration is too long, then the muscle will be too active during muscle 490 491 lengthening and at rest, causing increased work done on the muscle during lengthening and as result decreased 492 net work (net work = work done by the muscle during shortening –work done on the muscle during lengthening 493 (Josephson, 1993). The phase shift denotes the time at which the electrical stimulation is applied, relative to peak 494 muscle length. Negative values were used (-10 ms for soleus, -5 ms for Diaphragm and -2 ms for EDL). The value 495 of -10 ms used for the soleus meant that 10 ms prior to maximal length, stimulation of the muscle began. This 496 meant that 55 ms (based on a burst duration of 65) of stimulation occurred during shortening. The values for phase 497 are typical of previous work (Tallis et al. 2014; Tallis et al. 2017b). The values for strain, cycle frequency, burst 498 duration and stimulus phase were altered after interpreting the work loop shapes and net work value.
- It is recognised that the performance of isolated mouse muscle will decrease very slowly over time due to the development of an anoxic core (Barclay. 2005). In accordance with previous studies using a similar protocol (James *et al.* 1996; James *et al.* 1995), a series of 'control' assessments of work were made at various time points across the protocol. As such the performance of the muscle across the time course of the experiment could be corrected by considering the small decline in muscle performance over time.
- Following the final work loop, each muscle was rested for 10 minute before being subjected to 50 consectutive workloops (at 5Hz, 7Hz and 10Hz cycle frequency for SOL, DIA and EDL respectively) to assess fatigue resistance. Net work was analysed for every second work loop until the muscle began to produce negative work. This protocol has been used previously to assess fatigability in isolated skeletal msucle (Tallis *et al.* 2014; Tallis *et al.* 2017b).
- Each muscle underwent a 30-minute recovery period with 4 work loops being performed at each of 10, 20 and 30
 minutes. These were used to measure the ability of the muscle to recover following fatigue (Tallis *et al.* 2014;

Tallis *et al.* 2017b). Each experiment lasted between 2.5 and 3 hours, this was dependent time taken to optimise
work loop power at each cycle frequency.

14

513 514

Muscle Mass Measurements

515 Upon completion, the muscle was removed from the crocodile clips and the remaining tendon, bone and 516 aluminium clips were removed. Any excess fluid remaining on the muscle was removed by blotting the tissue on 517 absorbent tissue paper. The muscle was then weighed to attain wet muscle mass to the nearest 0.0001g. Muscle 518 cross sectional area was then calculated from the wet muscle mass, fibre length and an assumed muscle density 519 of 1060 kg m⁻³ (Mendez and Keys. 1960) Isometric stress was calculated as force divided by mean muscle cross-520 sectional area. Muscle power output was normalised to muscle mass to express power as W.kg⁻¹.

521 522

Statistical Methods

Following appropriate checks of normality and homogeneity, data for animal morphology, isometric stress, force, work loop power output, and fatigue was analysed using a one-way ANOVA with planned contrasts with all treatment groups compared against the lean control group in SPSS (SPSS, IL, USA), and Dunnett's post hoc analysis was used with P<0.05 to indicate significant differences between treatment groups. Two-way ANOVA were used to assess the effects of feeding duration and cycle frequency on maximal power output. Correlations were performed on all muscles between fat free mass and body mass, isometric force, isometric stress, absolute work loop power output, normalised work loop power output and fatigue resistance.

530

531 Acknowledgements

532 We wish to acknowledge Mark Bodycote and Bethan Grist for technical assistance.

The authors have no conflict of interest to declare.

- 533 Conflict of interests
- 534 535

536 537

538

552 Tables:

Table 1: Nutritional value of the diets given to the animals.

		SDS RM-1 Maintenance	Advanced Protocol PicoLab Natural Sunflower
	Calories provided by:		
	Protein (%)	17.49	17.95
	Fat (%)	7.42	63.66
	Carbohydrates (%)	75.09	18.39
	Gross energy (kcal.g ⁻¹)	3.52	5.24
	Digestible energy (kcal.g ⁻¹)	2.57	3.80
	Fatty acids content:		
	Saturated (%)	0.51	2.61
	Monounsaturated (%)	0.88	5.36
554			
555			
556			

Treatment Group	Control		2 week		4 week		8 week		12 week	
	Mean	SEM	Mean	SEM	Mean	SEM	Mean	SEM	Mean	SEM
Body Mass (g)	33.3	4.1	36.4	4.9	35.9	5.6	41.2*	7.1	38.7*	6.0
Body Length (cm)	10.4	0.4	10.6	0.3	10.8	0.4	10.9	0.5	10.9	0.3
Gonadal Fat Pad Mass (g)	0.91	0.79	1.91	1.01	1.99	1.28	2.92*	2.39	2.30*	1.39
Gonadal Fat Pad Mass (%	2.86	0.5	5.39	0.6	5.06	0.6	7.34*	1.0	6.00*	0.5
of body mass)										
Circumference (cm)	7.48	0.49	8.52	0.79	8.49	0.84	9.24*	1.03	9.03*	0.93
SOL Mass (mg)	8.50	0.2	10.1	0.5	9.90	0.4	10.8*	0.5	12.3*	0.4
EDL Mass (mg)	10.4	0.6	11.9	0.7	12.9*	0.3	12.6*	0.3	13.7*	0.3

TABLE 2 - THE EFFECT OF HFD FEEDING DURATION ON THE WHOLE ANIMAL AND MUSCLE **MORPHOLOGY**

[Data represented as Mean + SEM; N= 24 for groups control, 2 and 4weeks; N= 30 for groups 8 and 12 weeks; ** indicate significant differences between treatment group and lean control group]*

TABLE 3 – THE EFFECT OF DIFFERENT DURATIONS OF A HFD ON ISOMETRIC TIME FROM LAST STIMULUS TO HALF TETANUS RELAXATION (LSHR) OF ISOLATED MOUSE SOLEUS, EDL AND DIAPHRAGM

Treatment	Control		2 weeks		4 weeks		8 weeks	8 weeks		12 weeks	
Group											
	Mean	SEM	Mean	SEM	Mean	SEM	Mean	SEM	Mean	SEM	
Soleus (ms)	45.0	2.9	45.9	4.8	47.8	6.2	52.7*	5.2	55.1*	4.0	
EDL (ms)	12.9	0.9	12.8	1.0	13.2	1.2	14.6	0.7	14.4	1.1	
Diaphragm	25.1	2.5	25.8	2.1	25.5	3.0	26.2	2.8	24.9	3.1	

(ms)

[Data represented as Mean + SEM N= 8 for SOL, DIA and EDL in groups Control, 2 and 4 weeks, and N= 10 for Soleus, Diaphragm and EDL in groups 8 and 12 weeks; * represents significant difference between treatment group and lean control group]

TABLE 4 – CORRELATION BETWEEN GONADAL FAT PAD MASS AND: BODY MASS, TETANUS FORCE, TETANUS STRESS, ABSOLUTE POWER OUTPUT, NORMALISED POWER OUTPUT AND FATIGUE FOR SOLEUS, EDL AND DIAPHRAGM. ALL EXPERIMENTAL GROUPS WERE POOLED FOR THIS ANALYSIS.

		Body mass	Tetanus	Tetanus	Absolute PO	Normalised	Fatigue (% of
			Force	Stress		РО	max)
SOLEUS	R Value	0.792	0.326	0.081	0.151	-0.216	0.191
	P Value	0.001*	0.03*	0.606	0.331	0.164	0.221
EDL	R Value	0.155	0.068	0.082	-0.008	-0.267	-0.357
	P Value	0.321	0.664	0.603	0.957	0.074	0.019*
DIAPHRAGM	R Value	0.613		-0.334		-0.021	-0.413
	P Value	0.001*		0.03*		-0.891	0.005*

[N=44 for each group; * represents significant correlation between body mass and variable]

- 604
- 605
- 606
- 607
- - -
- 608 609

610 Legends to figures:

- 611
- 612

Figure 1 – The effect of different durations of a HFD on the maximal isometric tetanus force and stress of isolated
mouse Soleus (a & b), EDL (c & d) and Diaphragm (e) [Data represented as Mean+SEM N= 8 for Soleus,
Diaphragm and EDL in groups Control, 2 and 4 weeks, and N= 10 for Soleus, Diaphragm and EDL in groups 8
and 12 weeks; * represent significant differences between treatment group and lean control group]

617

Figure 2 – The effect of different durations of a HFD on the normalised net WL PO and absolute net WL PO of
isolated mouse Soleus (a & b), EDL (c & d) and Diaphragm (e) over a range of cycle frequencies. [Data
represented as Mean; N= 8 for Soleus, Diaphragm and EDL in groups Control, 2 and 4 weeks, and N= 10 for
Soleus, Diaphragm and EDL in groups 8 and 12 weeks * represents significant difference between treatment group
and lean control group]

623

Figure 3 - The effect of different durations of a HFD on the fatigue resistance of maximally stimulated mouse
Soleus (a), EDL (b) and Diaphragm (c) [Data represented as Mean±SEM; N= 8 for Soleus, Diaphragm and EDL

626 in groups Control, 2 and 4 weeks, and N= 10 for Soleus, Diapragm and EDL in groups 8 and 12 weeks]

- 627
- 628



8 Weeks



Experimental group

4 Weeks



4 Weeks

Experimental group

2 Weeks

637

0

Control

629

Figures: Figure One:

639

638

- 640
- 641

8 Weeks

12 Weeks





- 662 References
- Abdelmoula A, Martin V, Bouchant A *et al* (2012) Knee extension strength in obese and nonobese male
 adolescents. Applied Physiology, Nutrition, and Metabolism 37:269-275
- Akhmedov D, Berdeaux R (2013) The effects of obesity on skeletal muscle regeneration. Front Physiol 4:371.
 doi:10.3389/fphys.2013.00371 [doi]
- Altringham JD, Young IS (1991) Power output and the frequency of oscillatory work in mammalian diaphragm
 muscle: the effects of animal size. J Exp Biol 157:381-389
- Anderson SR, Gilge DA, Steiber AL, Previs SF (2008) Diet-induced obesity alters protein synthesis: tissue specific effects in fasted versus fed mice. Metab Clin Exp 57:347-354
- Aucouturier J, Lazaar N, Dore E, Meyer M, Ratel S, Duche P (2007) Cycling peak power in obese and lean 6-to
 8-year-old girls and boys. Applied Physiology, Nutrition, and Metabolism 32:367-371
- Barclay C (2005) Modelling diffusive O 2 supply to isolated preparations of mammalian skeletal and cardiac
 muscle. J Muscle Res Cell Motil 26:225-235
- Barclay C (1996) Mechanical efficiency and fatigue of fast and slow muscles of the mouse. The Journal of
 Physiology, 497:781-794.
- Blimkie CJ, Ebbesen B, MacDougall D, Bar-Or O, Sale D (1989) Voluntary and electrically evoked strength
 characteristics of obese and nonobese preadolescent boys. Human biology :515-532
- Bott KN, Gittings W, Fajardo VA *et al* (2017) Musculoskeletal structure and function in response to the
 combined effect of an obesogenic diet and age in male C57BL/6J mice. Molecular Nutrition & Food Research
- Brotto MA, Biesiadecki BJ, Brotto LS, Nosek TM, Jin JP (2006) Coupled expression of troponin T and troponin
 I isoforms in single skeletal muscle fibers correlates with contractility. Am J Physiol Cell Physiol 290:C567-76.
 doi:00422.2005 [pii]
- Bruton JD, Katz A, Lännergren J, Abbate F, Westerblad H (2002) Regulation of myoplasmic Ca 2 in genetically
 obese (ob/ob) mouse single skeletal muscle fibres. Pflügers Archiv European Journal of Physiology 444:692699
- Capodaglio P, Vismara L, Menegoni F, Baccalaro G, Galli M, Grugni G (2009) Strength characterization of
 knee flexor and extensor muscles in Prader-Willi and obese patients. BMC Musculoskeletal Disorders 10:47
- 689 Ciapaite J, Van Den Berg, Sjoerd A, Houten SM, Nicolay K, van Dijk KW, Jeneson JA (2015) Fiber-type-
- specific sensitivities and phenotypic adaptations to dietary fat overload differentially impact fast-versus slow twitch muscle contractile function in C57BL/6J mice. J Nutr Biochem 26:155-164
- de Wilde J, Mohren R, van den Berg S *et al* (2008) Short-term high fat-feeding results in morphological and metabolic adaptations in the skeletal muscle of C57BL/6J mice. Physiol Genomics 32:360-369. doi:00219.2007
- 694 [pii]
- Denies MS, Johnson J, Maliphol AB *et al* (2014) Diet-induced obesity alters skeletal muscle fiber types of male
 but not female mice. Physiol Rep 2:e00204. doi:10.1002/phy2.204 [doi]
- 697 D'Souza DM, Trajcevski KE, Al-Sajee D et al (2015) Diet-induced obesity impairs muscle satellite cell
- activation and muscle repair through alterations in hepatocyte growth factor signaling. Physiol Rep
- 699 3:10.14814/phy2.12506. doi:10.14814/phy2.12506 [doi]

- 700 Eshima H, Tamura Y, Kakehi S et al (2017) Long-term, but not short-term high-fat diet induces fiber
- composition changes and impaired contractile force in mouse fast-twitch skeletal muscle. Physiol Rep
 5:10.14814/phy2.13250. doi:e13250 [pii]
- Goodpaster BH, He J, Watkins S, Kelley DE (2001) Skeletal muscle lipid content and insulin resistance:
 evidence for a paradox in endurance-trained athletes. The Journal of Clinical Endocrinology & Metabolism
 86:5755-5761
- Hulens M, Vansant G, Lysens R, Claessens A, Muls E, Brumagne S (2001) Study of differences in peripheral
 muscle strength of lean versus obese women: an allometric approach. Int J Obes 25:676
- James RS, Kohlsdorf T, Cox VM, Navas CA (2005) 70 μM caffeine treatment enhances in vitro force and
 power output during cyclic activities in mouse extensor digitorum longus muscle. Eur J Appl Physiol 95:74-82
- James RS, Young IS, Cox VM, Goldspink DF, Altringham JD (1996) Isometric and isotonic muscle properties
 as determinants of work loop power output. Pflügers Archiv European Journal of Physiology 432:767-774
- James RS, Altringham JD, Goldspink DF (1995) The mechanical properties of fast and slow skeletal muscles of
 the mouse in relation to their locomotory function. J Exp Biol 198:491-502
- Josephson RK (1993) Contraction dynamics and power output of skeletal muscle. Annu Rev Physiol 55:527 546
- Josephson RK (1985) Mechanical power output from striated muscle during cyclic contraction. J Exp Biol
 114:493-512
- Kemp J, Blazev R, Stephenson DG, Stephenson G (2009) Morphological and biochemical alterations of skeletal
 muscles from the genetically obese (ob/ob) mouse. Int J Obes 33:831
- 720 Kopelman P (2007) Health risks associated with overweight and obesity. Obesity reviews 8:13-17
- 721 Machann J, Bachmann OP, Brechtel K et al (2003) Lipid content in the musculature of the lower leg assessed by
- fat selective MRI: Intra-and interindividual differences and correlation with anthropometric and metabolic data.
- 723 Journal of Magnetic Resonance Imaging 17:350-357
- Maffiuletti NA, Ratel S, Sartorio A, Martin V (2013) The impact of obesity on in vivo human skeletal muscle
 function. Current Obesity Reports 2:251-260
- Maffiuletti NA, Jubeau M, Munzinger U *et al* (2007) Differences in quadriceps muscle strength and fatigue
 between lean and obese subjects. Eur J Appl Physiol 101:51-59
- Matsakas A, Prosdocimo DA, Mitchell R *et al* (2015) Investigating mechanisms underpinning the detrimental
 impact of a high-fat diet in the developing and adult hypermuscular myostatin null mouse. Skeletal muscle 5:38
- Mendez J, Keys A (1960) Density and composition of mammalian muscle
 Metabolism :184-188
- Miyatake N, Fujii M, Nishikawa H *et al* (2000) Clinical evaluation of muscle strength in 20–79-years-old obese
 Japanese. Diabetes Res Clin Pract 48:15-21
- Paolillo FR, Milan JC, Bueno Pde G *et al* (2012) Effects of excess body mass on strength and fatigability of
 quadriceps in postmenopausal women. Menopause 19:556-561. doi:10.1097/gme.0b013e3182364e80 [doi]
- 736 Pette D, Staron RS (1997) Mammalian skeletal muscle fiber type transitions. Int Rev Cytol 170:143-223

- Rogers P,Webb G.P (1980) Estimation of body fat in normal and obese mice. British Journal of Nutrition, 43(1),
 pp.83-86.
- Rolland Y, Lauwers-Cances V, Pahor M, Fillaux J, Grandjean H, Vellas B (2004) Muscle strength in obese
 elderly women: effect of recreational physical activity in a cross-sectional study. Am J Clin Nutr 79:552-557
- Seebacher F, Tallis J, McShea K, James R (2017) Obesity-induced decreases in muscle performance are not
 reversed by weight loss. Int J Obes
- Shortreed KE, Krause MP, Huang JH *et al* (2009) Muscle-specific adaptations, impaired oxidative capacity and
 maintenance of contractile function characterize diet-induced obese mouse skeletal muscle. PloS one 4:e7293
- Singh R, Kaushik S, Wang Y, Xiang Y, Novak I, Komatsu M, Tanaka K, Cuervo A.M, Czaja, M.J (2009)
 Autophagy regulates lipid metabolism. *Nature*, 458(7242), p.1131.
- Szymura J, Gradek J, Maciejczyk M, Więcek M, Cempla J (2011) The anaerobic capacity in obese children.
 Gastroenterologia Polska 18:147-153
- Tallis J, Higgins MF, Seebacher F, Cox VM, Duncan MJ, James RS (2017a) The effects of 8 weeks voluntary
 wheel running on the contractile performance of isolated locomotory (soleus) and respiratory (diaphragm)
 skeletal muscle during early ageing. J Exp Biol 220:3733-3741. doi:10.1242/jeb.166603 [doi]
- 752 Tallis J, Hill C, James RS, Cox VM, Seebacher F (2017b) The effect of obesity on the contractile performance
- of isolated mouse soleus, EDL, and diaphragm muscles. J Appl Physiol (1985) 122:170-181.
- 754 doi:10.1152/japplphysiol.00836.2016 [doi]
- Tallis J, James, RS, Seebacher F. The effects of obesity on skeletal muscle contractile function. Journal of
 Experimental Biology. (Accepted for publication on 24/4/18).
- 757 Tallis J, James RS, Little AG, Cox VM, Duncan MJ, Seebacher F (2014) Early effects of ageing on the
- mechanical performance of isolated locomotory (EDL) and respiratory (diaphragm) skeletal muscle using the
 work-loop technique. Am J Physiol Regul Integr Comp Physiol 307:R670-84. doi:10.1152/ajpregu.00115.2014
 [doi]
- Tallis J, James RS, Cox VM, Duncan MJ (2012) The effect of physiological concentrations of caffeine on the
 power output of maximally and submaximally stimulated mouse EDL (fast) and soleus (slow) muscle. J Appl
 Physiol (1985) 112:64-71. doi:10.1152/japplphysiol.00801.2011 [doi]
- Tanner CJ, Barakat HA, Dohm GL *et al* (2002) Muscle fiber type is associated with obesity and weight loss.
 Am J Physiol Endocrinol Metab 282:E1191-6. doi:10.1152/ajpendo.00416.2001 [doi]
- 766 Thomas MM, Trajcevski KE, Coleman SK *et al* (2014) Early oxidative shifts in mouse skeletal muscle
- morphology with high-fat diet consumption do not lead to functional improvements. Physiological reports2:e12149
- Tomlinson D, Erskine R, Morse C, Winwood K, Onambélé-Pearson G (2016) The impact of obesity on skeletal
 muscle strength and structure through adolescence to old age. Biogerontology 17:467-483
- Trajcevski KE, O'Neill HM, Wang DC *et al* (2013) Enhanced lipid oxidation and maintenance of muscle insulin
 sensitivity despite glucose intolerance in a diet-induced obesity mouse model. PLoS One 8:e71747
- Tuttle LJ, Sinacore DR, Mueller MJ (2012) Intermuscular adipose tissue is muscle specific and associated with
 poor functional performance. Journal of aging research 2012
- Ward DS, Trost SG, Felton G *et al* (1997) Physical Activity and Physical Fitness in African-American Girls
 With and Without Obesity. Obesity 5:572-577

- Warmington S, Tolan R, McBennett S (2000) Functional and histological characteristics of skeletal muscle and
 the effects of leptin in the genetically obese (ob/ob) mouse. Int J Obes 24:1040
- 779 WHO (2017) Obesity and overweight, World Health, Organisation . 2017
- 780 WOF (2017) Obesity Country Profiles, World Obesity Federation. 2017
- Wronska A, Kmiec Z (2012) Structural and biochemical characteristics of various white adipose tissue depots.
 Acta Physiologica, 205(2), pp.194-208.
- Yoshida Y, Marcus RL, Lastayo PC (2012) Intramuscular adipose tissue and central activation in older adults.
 Muscle Nerve 46:813-816
- 785 Zurlo F, Larson K, Bogardus C, Ravussin E (1990) Skeletal muscle metabolism is a major determinant of
- resting energy expenditure. J Clin Invest 86:1423-1427. doi:10.1172/JCI114857 [doi]