

Quantification of MRI and MRS characteristics changes in a rat model at different stage of cerebral ischemia

Zhang, Y., Hao, D., Lv, X., Li, S., Tian, Y., Zheng, D. & Zeng, Y.

Author post-print (accepted) deposited by Coventry University's Repository

Original citation & hyperlink:

Zhang, Y, Hao, D, Lv, X, Li, S, Tian, Y, Zheng, D & Zeng, Y 2016, 'Quantification of MRI and MRS characteristics changes in a rat model at different stage of cerebral ischemia' *Neurological Research*, vol. 38, no. 7, pp. 640-646.
<https://dx.doi.org/10.1080/01616412.2016.1181345>

DOI 10.1080/01616412.2016.1181345

ISSN 0161-6412

ESSN 1743-1328

Publisher: Maney Publishing

Copyright © and Moral Rights are retained by the author(s) and/ or other copyright owners. A copy can be downloaded for personal non-commercial research or study, without prior permission or charge. This item cannot be reproduced or quoted extensively from without first obtaining permission in writing from the copyright holder(s). The content must not be changed in any way or sold commercially in any format or medium without the formal permission of the copyright holders.

This document is the author's post-print version, incorporating any revisions agreed during the peer-review process. Some differences between the published version and this version may remain and you are advised to consult the published version if you wish to cite from it.

Quantification of MRI and MRS characteristics changes in a rat model at different stage of cerebral ischemia

Yan Zhang¹, Dongmei Hao¹, Xiuhua Lv¹, Shaowu Li², Yunqing Tian³, Dingchang Zheng⁴, Yanjun Zeng^{2,*}

¹College of Life Science and Bioengineering, Beijing University of Technology, Beijing, China, ²Beijing Neurosurgical Institute, Beijing Tiantan Hospital, Capital Medical University, Beijing, China, ³State Intellectual Property Office of the P.R.C, Beijing, China, ⁴Faculty of Medical Science, Health and Wellbeing Academy, Anglia Ruskin University, Chelmsford, UK

Background: A better understanding the mechanisms of cerebral ischemia is important both for diagnosis and treatment.

Objective: The study aimed to quantify several characteristics of magnetic resonance imaging (MRI) and magnetic resonance spectroscopy (MRS) to indicate the brain tissue changes at different stage of cerebral ischemia in rats.

Methods: In the present study, a rat model of cerebral ischemia was established by middle cerebral artery occlusion (MCAO) in the left hemisphere. MRI and MRS were performed on 15 Sprague Dawley rats 4 H, 24 H, and 1 W after MCAO. Apparent diffusion coefficient (ADC), relative ADC including FNR, PNR, PNF, and metabolite ratio NCC were proposed to reflect the changes of water diffusion and metabolism in brain tissue.

Results: ADCs of focal zone and penumbra zone from 1 W group were significantly larger than those from 4H group, respectively (both $p < 0.05$). PNR and PNF of 24H and 1 W groups were significantly less than 4H group (all $p < 0.01$). NCCs of focal zone and penumbra zone were significantly less than the normal zone within 4H, 24H, and 1 W groups, respectively (both $p < 0.01$). While NCCs of penumbra zone from 24H and 1 W groups were significantly larger than 4H group (both $p < 0.01$).

Conclusion: We conclude that combination of MRI and MRS characteristics can provide significant indicators for ischemic damage at different stage of cerebral ischemia in a rat model.

Keywords: Magnetic resonance imaging (MRI), Magnetic resonance spectroscopy (MRS), Cerebral ischemia, Middle cerebral artery occlusion (MCAO), Rat

Introduction

Cerebral ischemia is a condition in which there is insufficient blood flow to the brain to meet metabolic demand. This leads to poor oxygen supply or cerebral hypoxia and thus to ischemic stroke, which may involve impairments in body movement, vision and speaking. Neurological deficits are common sequelae of cerebral ischemia and neurological recovery has been accompanied by changes in brain activation patterns. What happen in the brain structure and metabolism accompany by cerebral ischemia are not very clear.

Rodent models of cerebral ischemia have contributed greatly to our understanding of the mechanisms of the occurrence and evolution of ischemic stroke. Magnetic Resonance Imaging (MRI) has been used previously in

animal models to non-invasively and quantitatively evaluate the progression of focal ischemic lesions.^{1,2} The neurological impairments and restoration have been linked to the presence of post ischemic lesions on T2-weighted MRI and diffusion weighted imaging (DWI).³

T2-weighted MRI provides a degree of certainty in assigning irreversible ischemia. DWI monitors disturbances of ion homeostasis, water distribution, and tissue microstructure.³ The intensity of each image voxel reflects the best estimate of the rate of water diffusion at that location. DWIs visualize the ischemic core that is likely to proceed into infarction within a few hours from the onset of ischemia.⁴ Following an ischemic stroke, DWI is more sensitive to the changes occurring in the lesion than the traditional MRI measurements such as T1 or T2 relaxation rates. Although the apparent diffusion coefficient (ADC) measured from DWI has been shown to decrease immediately after the onset of ischemia,⁵ more investigations would be helpful to support the viewpoint that quantitative diffusion coefficient

Correspondence to: Dongmei Hao, College of Life Science and Bioengineering, Beijing University of Technology, Beijing 100124, China. Email: haodongmei@bjut.edu.cn; Yanjun Zeng, Beijing Neurosurgical Institute, Beijing Tiantan Hospital, Capital Medical University, Beijing, China. Email: yjzeng@bjut.edu.cn

*Present affiliation: College of Life Science and Bioengineering, Beijing University of Technology, Beijing 100124, China.

changes can be successfully applied for the essential distinction between the ischemic core and a potentially salvageable penumbra zone.

Several analytical approaches have been explored to predict the severity of the brain lesion from multiparametric MRI data acquired in acute stroke.⁶ These studies exploited diffusion, perfusion, and T2 MR data according to the biophysical and pathologic information from the ischemic regions provided by these parameters. Multiparameter studies using quantitative T1, T2, and diffusion MRI have shown the potential to predict the infarct size with prior knowledge of the location of stroke.^{7,8} These observations are significant as far as the prediction of the ischemic core is concerned in the hyperacute phase of the stroke. However, diffusion mismatch is not sufficient to predict the outcome of ischemic brain.^{9,10}

Magnetic resonance spectroscopy (MRS), which is based on nuclear magnetic resonance can give information on the severity of ischemic injury by measuring metabolites after cerebral ischemia. MRS provides a non-invasive diagnostic tool for the biochemical characterization of pathophysiological processes in the brain, which allows assessment of several metabolite concentrations known to be altered following cerebral ischemia.^{11,12} In brain MRS, the resonances of interest are N-acetyl aspartate (NAA), choline (Cho), creatine (Cr), myo-inositol (mI), lactate (Lac), lipids (Lip), glutamine and glutamate (Glx), and amino acids.¹³ In particular, a decrease in NAA has been shown to correlate with neuronal loss which was used to predict outcome after stroke.^{14,15}

Most of the previous study investigated cerebral ischemia by partial combination of MRI, immunohistochemistry, behavioral status, MRS, and histopathology.^{1,2} They mainly focused on the characteristic changes in the lesion volume^{3,16} or specific sensorimotor areas² in transient cerebral ischemia, for example, at time points between 3 and 48 h,^{16,17} which is not sufficient to analyze the cerebral ischemia comprehensively.

The purpose of this study was to quantify and compare both MRI and MRS characteristics of focal, penumbra, and normal zones 4 h, 24 h, and 1 week after the onset of ischemia in rats. It helps to elucidate the mechanisms underlying brain impairment and variation, and aids in optimization of diagnostic methods.

Materials and methods

Animals

Adult male Sprague–Dawley (SD) rats aged 8–10 weeks, weighing 200–250 g were purchased from BEIJING HFK BIOSCIENCE CO., LTD in this experiment (License No. SCXK (Beijing), 2009- 0007). They were housed in transparent Makrolon cages during 12 h day/night cycle in a temperature-controlled room (25 °C) with free access to food and water. This study received permission from

the Animal Care and Research Committee of Beijing University of Technology, Beijing, China.

Middle cerebral artery occlusion

Rats were anesthetized by an intramuscular injection of 10% chloral hydrate (0.4 ml/100 g, provided by department of pharmacology, Peking union medical college hospital) and placed in the prone position on a stereotaxic system (DW-5, Chengdu Taimeng Technology CO., Ltd., China). The rat's left carotid region was exposed through a midline cervical incision (about 20 mm). The left common carotid artery (CCA), internal carotid artery (ICA), and external carotid artery (ECA) were isolated from the surrounding tissue. A nylon suture (product code: 2432-50, Beijing ShaDong biological technology CO., LTD) coated with poly-L-lysine, with its tip shaped round, was inserted into the left ICA and advanced to a point approximately 18–20 mm distal to the bifurcation of ECA and ICA, thereby occluding the origin of the middle cerebral artery (MCA). During surgery, the core temperature of the rat was maintained at 37 ± 1 °C by a heating lamp.

After recovery from anesthesia, rats were scored as follows: (0) no neurological deficit was observed, (1) failure to extend the forepaw of the right side, (2) circling to the right side, (3) falling to the right, and (4) unable to walk and had a depressed level of consciousness. Only the rats with a Longa's score 1–2 were selected for our study.¹⁸

Group of rats

The qualified 15 SD rats were selected and randomly divided into three groups: 4 h after MCAO (4 H group), 24 h after MCAO (24 H group), and 1 week after MCAO (1 W group) with 5 rats in each group.

Magnetic resonance scanning

MRI experiments were performed in a horizontal 7 T magnet (Bruker, Germany). A surface coil (2.3 cm × 1.5 cm) was used for brain imaging. Multislice coronal spin-echo diffusion-weighted images (DWI) were acquired with an echo planar (EP) diffusion-trace MRI sequence (matrix size = 108 × 128, TR = 4500 ms, TE = 35 ms, FOV = 33 × 40 mm², FA = 90°, b0 = 600 s/mm²).

T2-weighted images were acquired using a SE sequence (TE = 41 ms, TR = 3140 ms, FOV = 40 × 40 mm², matrix size = 240 × 320, slice thickness = 0.7 mm, FA = 180°).

Spectra acquisition was performed using a chemical shift imaging (CSI) (TE/TR = 135/1500 ms) spectroscopy method. Measurements were made in the regions of interest (ROI).

Data analysis

The normal, penumbra and focal zone were determined on T2WI by an experienced radiologist. Focal zone was selected in the highlight region and normal zone was selected in the contralateral region.




Figure 1 Example of focal, penumbra, and normal zone in T2 image

In pathology and anatomy, the penumbra was the area surrounding an ischemic focal zone, shown in Fig. 1.

ADC reflects the degree of diffusion of water molecules through different tissues. ADCs of three ROIs selected, respectively, in the normal, penumbra and focal zone were calculated and averaged by software Functool (GE AW 4.2). Relative ADCs were proposed to overcome the between-rat differences and reflect the MRI characteristics at different stage of ischemic stroke. Here, three parameters were defined as:

$$FNR = \frac{FZ - NZ}{NZ} \quad (1)$$

$$PNR = \frac{PZ - NZ}{NZ} \quad (2)$$

$$PNF = \frac{PZ - NZ}{FZ} \quad (3)$$

Where FZ, NZ, and PZ are ADCs in focal, normal, and penumbra zones, respectively.

FNR reflects the ADC in focal zone relative to normal zone, PNR reflects the ADC in penumbra zone relative to normal zone and PNF reflects the ADC in penumbra zone relative to normal zone normalized by focal zone. FNR, PNR, and PNF were calculated in 4H, 24H, and 1 W group, respectively.

Choline(Cho), N-Acetylaspartate (Naa) and Creatine (Cr) of three ROIs selected, respectively, in the normal, penumbra, and focal zone were measured on MRS and averaged. Then a new parameter NCC was defined to reduce the effect of between-rat variability.

$$NCC = \frac{Naa - Cho}{Cr} \quad (4)$$

Statistical analysis

Experimental measurement data was expressed as Mean \pm SD and was statistically processed by SPSS 18.0 software (IBM Corporation, New York, United States) using

one-Way ANOVA analysis. The ADCs were compared within and between groups and the relative parameters of focal and penumbra zones were compared between the three groups. $p < 0.05$ was considered statistically significant.

Results

MRI assessment

ROI selection

Figure 2 shows the DWI (left column) and T2WI (right column) in 4H, 24H and 1 W groups. The circles in DWI are ROIs selected in normal, penumbra, and focal zones, in which 1, 2, 3 are ROIs in focal zone, 4, 5, 6 are in normal zone and 7, 8, 9 are in penumbra zone.

Comparison of ADC between the three ROIs

Table 1 shows that the ADC in focal zone was significantly smaller than normal zone within 4H, 24H, and 1 W groups (all $p < 0.05$). ADC in penumbra zone was significantly smaller ($p < 0.05$) than normal zone within 4H and 24H group, but not significantly different in 1 W group.

ADC changes with ischemic time following MCAO for the three ROIs

From Table 1, it can also be seen that the ADCs in focal and penumbra zones from 1 W group were significantly larger than those from 4H group, respectively (both $p < 0.05$).

Relative ADC changes with ischemic time following MCAO

The relative ADCs are summarized in Table 2. FNR was not significantly different between 4H, 24H, and 1 W groups ($p > 0.05$). PNR and PNF of 24H and 1 W groups were significantly less than 4H group (all $p < 0.01$).

The results indicated that FNR increased, PNR and PNF decreased with ischemic time following MCAO.

MRS changes with ischemic time following MCAO

Figure 3 gives an example of MRS and its corresponding ROI from penumbra zone in 4 H, 24H, and 1 W group, where ROI was marked using a red rectangle.




Figure 2 DWI and T2 from (a) & (b) 4H group; (c) & (d) 24H group; (e) & (f) 1 W group. 1, 2, 3 are in focal zone 4, 5, 6 are in normal zone and 7, 8, 9 are in penumbra zone

Table 1 ADC in focal, penumbra and normal zones from 4H, 24H, and 1 W groups

Group	Focal zone	Penumbra zone	Normal zone
4H	492 ± 21	593 ± 33	791 ± 13
24H	517 ± 83	644 ± 39	756 ± 46
1 W	588 ± 93*	764 ± 42*	714 ± 24

* $p < 0.05$ with comparison referred to the 4H group.

Table 2 Relative ADC changes with ischemic time following MCAO

Group	FNR	PNR	PNF
4H	37.8 ± 3.2	25.0 ± 1.2	40.3 ± 2.7
24H	44.1 ± 2.1	14.8 ± 1.2*	26.5 ± 1.3*
1 W	47.7 ± 5.3	6.9 ± 0.9*	8.3 ± 0.3*

* $p < 0.01$, with comparison referred to the 4H group.




Figure 3 ROI from penumbra zone and its corresponding MRS. (a) and (b) 4H group; (c) and (d) 24H group; (e) and (f) 1 W group

Table 3 NCC in focal, penumbra and normal zones from 4H, 24H, and 1 W group

Group	Focal zone	Penumbra zone	Normal zone
4H	3.4 ± 0.3	7.1 ± 1.2	102.8 ± 5.3
24H	4.4 ± 0.2	$29.0 \pm 2.3^*$	107.8 ± 7.5
1 W	7.7 ± 1.3	$148.0 \pm 13.1^*$	160.8 ± 11.1

* $p < 0.01$ with comparison referred to the 4H group.

The NCCs are summarized in Table 3. NCCs of focal zone and penumbra zone were significantly less than the normal zone within 4H, 24H, and 1 W groups, respectively ($p < 0.01$). Both NCCs of focal zone and normal zone presented no significant difference between 4H, 24H, and 1 W groups ($p > 0.05$). While NCCs of penumbra zone from 24H and 1 W groups were significantly larger than 4H group ($p < 0.01$). NCC increased with ischemic time following MCAO.

Discussion

Brain edema is one of the main complications in cerebral ischemia.³ In DWI, the intensity of each image voxel reflects the best estimate of the rate of water diffusion at that location. Because the mobility of water is driven by thermal agitation and highly dependent on its cellular environment, diffusion changes are correlated to the clinical deficit and are potentially useful for early diagnosis and

longitudinal evaluation. For instance, DWI is more sensitive to early changes after a stroke than more traditional MRI measurements such as T1 or T2 relaxation rates. To elucidate diffusion and relaxation effects on image contrast, one may obtain quantitative images of the diffusion coefficient, or more exactly the ADC.

In our study, the reduced ADC of water in focal and penumbra zones are in agreement with the previous studies.¹⁷ The relative ADC ratios presented are helpful to reflect the water diffusion of focal and penumbra zone relative to normal zone after MCAO. It was observed that the relative ADC in the focal zone (FNR) did not change significantly as time elapsed, which indicated the injury in the focal zone might be irreversible. Immediately following the stroke, blood flow and therefore oxygen transport is reduced locally, leading to hypoxia of the cells near the location of the original insult. This can lead to hypoxic cell death (infarction) and amplify the original damage from the ischemia; however, the penumbra area may remain viable after an ischemic event due to the collateral arteries that supply the penumbral zone. PNR and PNF being significantly decreased with time suggested that water diffusion of penumbra zone approached to normal zone. Penumbra is under an elevated risk of infarction, and monitoring its variation may provide important information for clinicians.¹⁹

N-acetylaspartate (Naa) is the second most abundant metabolite in the human central nervous system (CNS). Naa emits the strongest signal in MRS of the human brain, which has clinical significance. Many neurological disorders involving neuronal loss or dysfunction result in reductions in brain Naa levels. Choline (Cho) is the precursor molecule for the neurotransmitter acetylcholine, which is involved in many functions including memory and muscle control. It is used in the synthesis of the constructional components in the body's cell membranes. Creatine (Cr) is a nitrogenous organic acid that occurs naturally in vertebrates and can be a reservoir to store energy as phosphocreatine.²⁰

Hirt's study compared NAA, taurine, Glu, and Gln between 3 h and 8 h and considered NAA + Glu + Tau as a predictor of an irreversible lesion.¹⁶ To reduce the effect of between-rat variability, our study focused on the within-rat ratio of NAA, Cho, and Cr and their changes with time (4H, 24H, and 1 W). The proposed NCC in this study shows the evident differences of normal, penumbra, and focal zones within 4H, 24H, and 1 W groups and indicates the significant changes in penumbra zone between 24H, 1 W, and 4H groups, which may be useful to reflect the metabolic changes at different stage of cerebral ischemia.

MRI and MRS characteristics can be performed in animal models of 'lacunar' stroke, which may have practical implications to the treatment of cerebral small vessel disease with drugs to improve endothelial function given the non-atherothrombotic etiology of small vessel disease.²¹ The prior presence of a transient ischemic attack

(TIA) is associated with an early good outcome in non-lacunar ischemic strokes, which suggesting a neuroprotective effect of TIA possibly by inducing a phenomenon of ischemic tolerance.²² TIA model will be studied in the next step, and the penumbra location and focal size could be measured and investigated combined with the present characteristics in a further study.

In summary, our investigation illustrates that ADC measured using DWI is a viable surrogate marker of the damage and variation of brain tissue after a stroke. The relative ratio of Naa, Cho, and Cr can reflect the metabolic changes in normal, focal, and penumbra zones at different stage of cerebral ischemia. The combination of MRI and MRS techniques provides a good understanding of the evolution of ischemic damage and to probe the efficacy of future therapeutic interventions in a rat model of stroke.

Conflict of interest

The authors declare no conflict of interest.

Acknowledgments

The authors give special thanks to Dr Wangsheng Lu for his instruction to MCAO operation.

Funding

This work was supported by Beijing Natural Science Foundation [grant numbers 7132028 and 7132021] and National Natural Science Foundation of China [grant number 81441053].

Notes on contributors

Yan Zhang is a graduate student in the college of Life Science and Bioengineering, Beijing University of Technology. The author's areas of research interests include Ischemic Stroke, Physiological signals measurement. The author's recent publication is Somatosensory evoked potential changes in a rat model of cerebral ischemia. Chinese Journal of Biomedical Engineering 2014.

Dongmei Hao is an associate professor in the college of Life Science and Bioengineering, Beijing University of Technology. The author's areas of research interests include Ischemic Stroke, physiological signals detection and analysis. The author's recent publication is Applying support vector regression analysis on grip force level-related corticomuscular coherence. J Comput Neurosci 2014.

Xiuhua Lv is an associate professor in the college of Life Science and Bioengineering, Beijing University of Technology. The author's areas of research interests include cerebral palsy functional rehabilitation, Ischemic Stroke and traditional Chinese medicine. The author's recent publication is Somatosensory evoked potential changes in a rat model of cerebral ischemia. Chinese Journal of Biomedical Engineering 2014.

Shaowu Li is a professor in Beijing Neurosurgical Institute, Beijing Tiantan Hospital, Capital Medical University. The author's areas of research interests include Diagnosis of neural system disease with CT, DSA and MRI. The author's recent publication is Anatomical specificity of vascular endothelial growth factor expression in glioblastomas: a voxel-based mapping analysis. *Neuroradiology* 2016.

Yunqing Tian is a senior patent examiner in State Intellectual Property Office of the P.R.C. The author's areas of research interests include Medical electronic device development and medical signal processing. The author's recent publication is investigating the modulation of brain activity associated with handgrip force and fatigue. *Technology and Health Care* 2015.

Dingchang Zheng is a research fellow in Faculty of Medical Science, Health and Wellbeing Academy, Anglia Ruskin University. The author's areas of research interests include innovative medical technology, physiological measurement, cardiovascular signal processing and modeling. The author's recent publication is assessing the complexity of short-term heartbeat interval series by distance distribution entropy. *Medical & Biological Engineering & Computing* 2015.

YanJun Zeng is a professor in the college of Life Science and Bioengineering, Beijing University of Technology. The author's areas of research interests include Biomechanics, Neural engineering. The author's recent publication is the role of coupling strength and internal delay between compartments in shaping the bursting behavior of cortical neuron. *Neurological Sciences* 2014.

References

- 1 van de Looij Y, Chatagner A, Quairiaux C, Gruetter R, Hüppi PS, Sizonenko SV. Multi-modal assessment of long-term erythropoietin treatment after neonatal hypoxic-ischemic injury in rat brain. *Plos One*. 2014;9(4):e95643:1–10.
- 2 Sicard KM, Henninger N, Fisher M, Duong TQ, Ferris CF. Long-term changes of functional MRI-Based brain function, behavioral status, and histopathology after transient focal cerebral ischemia in rats. *Stroke*. 2006;37:2593–600.
- 3 Ashioti M, Beech JS, Lowe AS, Hesselink MB, Modo M, Williams SCR. Multi-modal characterisation of the neocortical clip model of focal cerebral ischaemia by MRI, behaviour and immunohistochemistry. *Brain Res*. 2007;1145:177–89.
- 4 Jokivarsi KT, Hiltunen Y, Tuunanen PI, Kauppinen RA, Gröhn OHJ. Correlating tissue outcome with quantitative multiparametric MRI of acute cerebral ischemia in rats. *J Cerebr. Blood Flow Metab*. 2010;30:415–27.
- 5 Weber R, Ramos-Cabrer P, Hoehn M. Present status of magnetic resonance imaging and spectroscopy in animal stroke models. *J Cerebr Blood Flow Metab*. 2006;26:591–604.
- 6 Shen Q, Meng X, Fisher M, Sotak CH, Duong TQ. Pixel-by-pixel spatiotemporal progression of focal ischemia derived using quantitative perfusion and diffusion imaging. *J Cerebr Blood Flow Metab*. 2003;23:88–1479.
- 7 Lu M, Mitsias PD, Ewing JR, Soltanian-Zadeh H, Bagher-Ebadian H, Zhao Q, *et al.* Predicting final infarct size using acute and subacute multiparametric MRI measurements in patients with ischemic stroke. *J Magn Reson Imaging*. 2005;21:495–502.
- 8 Mitsias PD, Jacobs MA, Hammoud R, Pasnoor M, Santhakumar S, Papamitsakis NI, *et al.* Multiparametric MRI ISODATA ischemic lesion analysis: correlation with the clinical neurological deficit and single-parameter MRI techniques. *Stroke*. 2002;33:44–2839.
- 9 Mezzapesa DM, Petruzzellis M, Lucivero V, Prontera M, Tinelli A, Sancilio M, *et al.* Multimodal MR examination in acute ischemic stroke. *Neuroradiology*. 2006;48:46–238.
- 10 Rivers CS, Wardlaw JM, Armitage PA, Bastin ME, Carpenter TK, Cvorov V, *et al.* Do acute diffusion- and perfusion-weighted MRI lesions identify final infarct volume in ischemic stroke? *Stroke*. 2006;37:98–104.
- 11 van de Looij Y, Chatagner A, Hüppi PS, Gruetter R, Sizonenko SV. Longitudinal MR assessment of hypoxic ischemic injury in the immature rat brain. *Magnet Reson Med*. 2011;65:305–12.
- 12 Tkáč I, Rao R, Georgieff MK, Gruetter R. Developmental and regional changes in the neurochemical profile of the rat brain determined by *in vivo* 1H NMR spectroscopy. *Magnet Reson Med*. 2003;50:24–32.
- 13 Gujar SK, Maheshwari S, Björkman-Burtscher I, Sundgren PC. Magnetic resonance spectroscopy. *J Neuro-Ophthalmol*. 2005;25(3):217–26.
- 14 Clark JB. N-acetyl aspartate: a marker for neuronal loss or mitochondrial dysfunction. *Int J Dev Neurosci*. 1998;20:271–6.
- 15 Pereira AC, Saunders DE, Doyle VL, Bland JM, Howe FA, Griffiths JR, *et al.* Measurement of initial N-Acetyl aspartate concentration by magnetic resonance spectroscopy and initial infarct volume by MRI predicts outcome in patients with middle cerebral artery territory infarction. *Stroke*. 1999;30:1577–82.
- 16 Berthet C, Lei HX, Gruetter R, Hirt L. Early predictive biomarkers for lesion after transient cerebral ischemia. *Stroke*. 2011;42:799–805.
- 17 Sobrado M, Delgado M, Fernández-Valle E, García-García L, Torres M, Sánchez-Prieto J, *et al.* Longitudinal studies of ischemic penumbra by using 18F-FDG PET and MRI techniques in permanent and transient focal cerebral ischemia in rats. *NeuroImage*. 2011;57:45–54.
- 18 Longa EZ, Weinstein PR, Carlson S, Cummins R. Reversible middle cerebral artery occlusion without craniectomy in rats. *Stroke*. 1989;20:84–91.
- 19 Guadagno JV, Calautti C, Baron JC. Progress in imaging stroke: emerging clinical applications. *Brit Med Bull*. 2003;65(1):145–57.
- 20 Miller BL. A review of chemical issues in 1H NMR spectroscopy: N-acetyl-L-aspartate, creatine and choline. *NMR Biomed*. 1991;4(2):47–52.
- 21 Arboix A, Blanco-Rojas L, Martí-Vilalta JL. Advancements in understanding the mechanisms of symptomatic lacunar ischemic stroke: translation of knowledge to prevention strategies. *Expert Rev Neurother*. 2014;14(3):261–76.
- 22 Arboix A, Cabeza N, García-Eroles L, Massons J, Oliveres M, Targa C, *et al.* Relevance of transient ischemic attack to early neurological recovery after nonlacunar ischemic stroke. *Cerebrovasc Dis*. 2004;18(4):304–11.