

# Caspase inhibition via A3 adenosine receptors: A new cardioprotective mechanism against myocardial infarction

Hussain, A. , Gharanei, A. , Nagra, A. and Maddock, H.L.  
Author pre-print (submitted version) deposited in CURVE February 2015

**Original citation & hyperlink:**

Hussain, A. , Gharanei, A. , Nagra, A. and Maddock, H.L. (2014) Caspase inhibition via A3 adenosine receptors: A new cardioprotective mechanism against myocardial infarction. *Cardiovascular Drugs and Therapy*, volume 28 (1): 19-32.

<http://dx.doi.org/10.1007/s10557-013-6500-y>

**Publisher statement:**

The final publication is available at Springer via <http://dx.doi.org/10.1007/s10557-013-6500-y>.

**Copyright © and Moral Rights are retained by the author(s) and/ or other copyright owners. A copy can be downloaded for personal non-commercial research or study, without prior permission or charge. This item cannot be reproduced or quoted extensively from without first obtaining permission in writing from the copyright holder(s). The content must not be changed in any way or sold commercially in any format or medium without the formal permission of the copyright holders.**

**This document is the submitted version of the journal article, as originally submitted to the journal prior to the peer-review process. Some differences between the published version and this version may remain and you are advised to consult the published version if you wish to cite from it.**

**CURVE is the Institutional Repository for Coventry University**  
<http://curve.coventry.ac.uk/open>

1 **Caspase inhibition via A<sub>3</sub> adenosine receptors: a new cardioprotective mechanism against myocardial**  
2 **infarction**

3  
4  
5  
6 **Running Title: 2-CL-IB-MECA mediated cardioprotection**

7  
8 **Authors:** <sup>1</sup>Afthab Hussain B.Sc (Hons) Ph.D Lecturer

9  
10 <sup>1</sup>Ahmed Mayel Gharane B.Sc (Hons) Ph.D

11  
12 <sup>1</sup>Aarondeep Singh Nagra B.Sc (Hons)

13  
14 <sup>1</sup>Helen L Maddock B.Sc (Hons) Ph.D. Principal Lecturer

15  
16 9

17  
18 **Corresponding Author: Dr H. L. Maddock**

19  
20 **Biomolecular Sciences Applied Research Group**

21  
22 **Faculty of Health and Life Sciences**

23  
24 **Coventry University**

25  
26 **Coventry**

27  
28 **CV1 5FB**

29  
30 **Tel: 02476 8881 70**

31  
32 **Fax: 02476 888 702**

33  
34 **E-mail: [apx281@coventry.ac.uk](mailto:apx281@coventry.ac.uk)**

35  
36 19

37  
38 20

39  
40 21

41  
42

43  
44

45  
46

47  
48

49  
50

51  
52

53  
54

55  
56

57  
58

59  
60

61  
62

63  
64

65

22 **Abstract**

1  
2 23 **Purpose.** 2-CL-IB-MECA, ( $A_3$  adenosine receptor agonist)( $A_3$ AR) mediated cardioprotection is well  
3  
4 24 documented although the intracellular signalling pathways associated remain unclear. Here we demonstrate a  
5  
6 25 role of the pro-survival signalling pathways MEK1/2-ERK1/2 and PI3K/AKT and their effect on modifying  
7  
8 26 Caspase-3 activity in  $A_3$ AR mediated cardioprotection.

9  
10 27 **Methods.** Isolated perfused rat hearts or primary adult rat cardiac myocytes were subjected to ischaemia/hypoxia  
11  
12 28 and reperfusion/reoxygenation, respectively. 2-CL-IB-MECA (1nM) was administered at the onset of  
13  
14 29 reperfusion/reoxygenation in the presence and absence of either the PI3K inhibitor Wortmannin (5nM) or  
15  
16 30 MEK1/2 inhibitor UO126 (10 $\mu$ M). Heart tissues were harvested for assessment of p-ERK1/2<sub>(Thr202/Tyr204)</sub> or p-  
17  
18 31 AKT<sub>(ser473)</sub> status or underwent infarct size assessment. Cardiac myocytes underwent flow-cytometric analysis  
19  
20 32 for apoptosis, necrosis, cleaved-caspase 3 / p-BAD (ser112) activity post-reoxygenation.

21  
22 33 **Results.** 2-CL-IB-MECA significantly reduced infarct size compared to non-treated controls, where co-  
23  
24 34 administration with either of the kinase inhibitors abolished the infarct sparing effects. Administration of 2-CL-  
25  
26 35 IB-MECA at reperfusion significantly upregulated the status of p-ERK1/2 and p-AKT compared to time  
27  
28 36 matched controls in a UO126 and Wortmannin sensitive manner respectively. 2-CL-IB-MECA when  
29  
30 37 administered throughout reoxygenation significantly reduced apoptosis, necrosis, cleaved-caspase 3 activity and  
31  
32 38 increased p-BAD (ser112) and p-BAD (ser136) activity in myocytes subjected to hypoxia/reoxygenation injury.  
33  
34 39 The cytoprotective effect was abolished by co-administration with the kinase inhibitors Wortmannin and/or  
35  
36 40 UO126.

37  
38 41 **Conclusions.** We have described the molecular mechanisms associated with  $A_3$ AR mediated cardioprotection  
39  
40 42 indicating a role for the pro-survival signalling pathways that decrease caspase-3 activity. These observations  
41  
42 43 provide novel insight into the pharmacological effects of  $A_3$ ARs in ameliorating myocardial  
43  
44 44 ischaemia/reperfusion injury.

45  
46 45  
47  
48 46 **Keywords:**  $A_3$  Adenosine receptor, Reperfusion injury, MEK1/2, PI3K, Apoptosis, Caspase 3, BAD.

## 50 **Introduction**

1 Adenosine is a purine nucleoside that is highly abundant throughout body where it plays a critical role in cellular  
2  
3  
4 52 function. Previous studies have shown adenosine, a metabolite of adenosine triphosphate (ATP), to accumulate  
5  
6 53 within the interstitial fluid within the ischaemic myocardium(1). Increased adenosine has profound effects in the  
7  
8 54 cardiovascular system including vasodilation, hypotension and bradycardia. Adenosine exerts its physiological  
9  
10 55 responses via four purinergic G-protein coupled adenosine receptors designated A<sub>1</sub>, A<sub>2a</sub>, A<sub>2b</sub> and A<sub>3</sub>(2).

11 56 Research over the past decade has implicated the putative role of adenosine in mediating an innate role in  
12  
13 57 protecting the myocardium from the deleterious consequences of ischaemia-reperfusion (I/R) injury (3-6). A<sub>1</sub>,  
14  
15 58 A<sub>2a</sub>, A<sub>2b</sub> and A<sub>3</sub> have been shown to exhibit cardioprotective properties, although controversy remains regarding  
16  
17 59 the precise timing of activation of these receptors required to induce cardioprotection, as well as their  
18  
19 60 involvement in ischaemic preconditioning and postconditioning(7-11). The ability of adenosine and adenosine  
20  
21 61 analogues to mediate cardioprotection is of clinical relevance as a possible adjunctive therapy to treat patients  
22  
23 62 with myocardial infarction or during cardiac surgery to limit the consequences of reperfusion injury(12).

24 63 Maddock *et al.* have previously shown activation of A<sub>3</sub>AR at the onset of reperfusion to protect the myocardium  
25  
26 64 from post ischaemic injury via an anti-apoptotic/necrotic mechanism that was abolished in the presence of A<sub>3</sub>AR  
27  
28 65 antagonist MRS 1191(8). A<sub>3</sub> adenosine receptor-mediated cardioprotection has also been reported to limit  
29  
30 66 myocardial I/R injury in a number of animal models including therat (10), rabbit(13), guinea pig (14), mice (7,  
31  
32 67 15)and dog (16). Adenosine receptors have previously been shown to be functionally expressed in a number of  
33  
34 68 organs including the myocardium (17, 18). The protection afforded by A<sub>3</sub>AR agonists can be directly attributed to  
35  
36 69 A<sub>3</sub>ARs, as in A<sub>3</sub>AR gene knock-out mice subjected to myocardial ischaemia reperfusion injury, administration  
37  
38 70 of the A<sub>3</sub>AR agonist failed to induce cardioprotection(10).

39 71 A<sub>3</sub>AR agonists have been reported to mediate protection from I/R injury in a number of organs including the  
40  
41 72 feline lung (19-21) and rat brain(22, 23). A<sub>3</sub>AR agonists have also been shown to promote cell survival in HL-60  
42  
43 73 leukaemia and U-937 lymphoma cells exposed to UV light by means of restricting apoptosis(24). Surprisingly,  
44  
45 74 there is limited data available that indicates the mechanisms responsible for this protection in the intact heart,  
46  
47 75 despite some information obtained in cellular models.

48  
49 76 Recent studies have shown that mitogen activated protein kinases MEK1/2 - extracellular regulated kinase  
50  
51 77 (ERK1/2) and phosphatidylinositol 3 kinase (PI3K) to become activated by a range of stimuli like growth factors  
52  
53 78 and hormones following receptor stimulation (25-28). Mitogen activated protein kinases (MAPKs) play a role in  
54  
55 79 cell survival, growth and differentiation and apoptosis (29). In this context, the MEK1/2-ERK1/2-BAD and  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

80 PI3K-AKT cell survival pathways have been shown in numerous studies to regulate cardioprotection upon  
1 81 receptor stimulation where it is referred to as the reperfusion injury salvage kinase pathway (RISK) (30).  
2  
3 82 Studies by Germack and Dickenson showed direct stimulation of A<sub>3</sub>ARs in neonatal rat cardiomyocytes to be  
4  
5 83 directly linked to the serine/threonine kinase ERK1/2 and AKT in a dose and time dependent manner in  
6  
7 84 normoxic conditions (31, 32). These further showed that preconditioning with the A<sub>3</sub>AR agonist 2-CL-IB-  
8  
9 85 MECA protected neonatal myocytes from hypoxia/reoxygenation injury in neonatal myocytes. Interestingly, a  
10  
11 86 number of cardioprotective agents including insulin, bradykinin and erythropoietin have been shown to have  
12  
13 87 infarct sparing effects when administered alone either before or during reperfusion via upregulation of AKT and  
14  
15 88 ERK pro-survival kinases (30).  
16  
17 89 The aim of this study was to evaluate the putative role of PI3K and MEK1/2 signalling pathways in A<sub>3</sub>AR  
18  
19 90 receptor mediated cardioprotection, as previous studies have shown A<sub>3</sub>ARs to activate PI3K and MEK1/2 in  
20  
21 91 normoxic conditions. Therefore, it was feasible that the cardioprotection exerted by 2-CL-IB-MECA when  
22  
23 92 administered at reperfusion may involve potential recruitment of the PI3K and MEK1/2 MAPK signalling  
24  
25 93 pathways.  
26  
27 94 Infarct sizes were measured after 2-CL-IB-MECA (1nM) was administered and the mechanisms responsible for 2-  
28  
29 95 CL-IB-MECA dependent cardioprotection in the isolated perfused heart were assessed using the MEK1/2  
30  
31 96 inhibitor UO126 (10µM) and the PI3K inhibitor Wortmannin (5nM). Tissues were also harvested at various time  
32  
33 97 points post reperfusion to determine the phosphorylation status of AKT<sub>(Ser473)</sub>, ERK<sub>(Thr202/Tyr204)</sub> and BAD<sub>(Ser112)</sub>.  
34  
35 98 Furthermore, we investigated the effect of the A<sub>3</sub>AR agonist 2-CL-IB-MECA in the presence and absence of  
36  
37 99 UO126 (10µM) or Wortmannin (5nM) when administered at reoxygenation on apoptosis, necrosis and cleaved-  
38  
39 100 caspase 3 activity in adult rat cardiomyocytes subjected to 6 hours of hypoxia and 18 hours of reoxygenation.  
40  
41  
42  
43

101

## 102 **Materials and Methods**

103 **Chemicals:** A<sub>3</sub>AR agonist 1-[2-Chloro-6-[[[(3-iodophenyl)methyl]amino]-9H-purin-9-yl]-1-deoxy-N-methyl-*b*-D-  
104 ribofuranuronamide (2-CL-IB-MECA), 1,4-Diamino-2,3-dicyano-1,4-bis[2-aminophenylthio]butadiene  
105 (UO126), (1S,6bR,9aS,11R,11bR) 11-(Acetyloxy)-1,6b,7,8,9a,10,11,11b-octahydro-1-(methoxymethyl)-9a,11b-  
106 dimethyl-3H-furo[4,3,2-de]indeno[4,5,-h]-2-h]-2-benzopyran-3,6,9-trione (Wortmannin) were purchased from  
107 Tocris Cookson (Bristol, UK). Total ERK, AKT, p-ERK1/2 (Thr<sup>202</sup> / Tyr<sup>204</sup>) p-AKT (Ser<sup>473</sup>), Cleaved-Caspase-3  
108 (Asp<sup>175</sup>) (5A1), p-BAD (Ser<sup>112</sup>) and p-BAD (Ser<sup>136</sup>) Rabbit mAb antibodies were purchased from New England  
109 Biolabs (Hertfordshire, UK). Alexa Fluor 488 F(ab')<sub>2</sub> fragment of goat anti-rabbit IgG (H+L) was purchased  
110  
111

110 from Invitrogen (Paisley, UK).  $\beta$ -actinantibody was purchased from Abcam (Cambridge, UK). Drugs were  
111 dissolved in DMSO and stored at  $-20^{\circ}\text{C}$ .

## 112 **Animals**

113 Male Sprague-Dawley rats were used in all experiments (350-400g b/w). Experiments were conducted in  
114 accordance with the Guidelines on the operation of Animals (Scientific Procedures) Act 1986. Animals were  
115 obtained from Charles River (Margate, UK).

## 116 **Isolated perfused heart preparation.**

117 Animals were sacrificed by cerebral dislocation. Hearts were rapidly excised and placed into ice-cold Krebs  
118 Heinsleit buffer (KH). Hearts were quickly mounted onto the Langendorff system and retrogradely perfused with  
119 modified Krebs Henseleit (KH) bicarbonate buffer containing (in mmol) NaCl 118.5,  $\text{NaHCO}_3$  25, KCL 4.8,  
120  $\text{MgSO}_4$  1.2,  $\text{KH}_2\text{PO}_4$  1.2,  $\text{CaCl}_2$  1.7, and glucose 11. KH buffer was gassed using 95%  $\text{O}_2$  and 5%  $\text{CO}_2$  and  
121 maintained at  $37^{\circ}\text{C}$ .

122 The left atrium was trimmed away and a latex isovolumic balloon was carefully introduced into the left ventricle  
123 and inflated up to 5-10mmHg. Functional recording was via a physiological pressure transducer connected to a  
124 bridge amp and a Powerlab (AD Instruments Ltd. Chalgrove, UK). Left ventricular pressure (LVDP), heart rate  
125 (HR) and coronary flow (CF) were measured at regular intervals. To induce regional ischaemia a surgical needle  
126 was inserted under the left main coronary artery and the ends of the thread were passed through a tube to form a  
127 snare. Tightening of the snare induced regional ischaemia and releasing the thread initiated reperfusion. At the  
128 end of reperfusion, the snare was tightened to re-occlude the coronary artery branch.

129 A solution of 0.25% Evans blue in saline was then infused slowly via the aorta to delineate the non risk zone of  
130 the myocardium, which stained dark blue. The hearts were weighed and frozen at  $-20^{\circ}\text{C}$ . Frozen hearts were then  
131 sliced into 2mm thick transverse sections and incubated in triphenyl-tetrazolium chloride solution (1% in  
132 phosphate buffer) at  $37^{\circ}\text{C}$  for 10-12minutes and fixed in 10% Formalin for at least 4 hours.

133 In the risk zone the viable issue stained red and infarct tissue appeared pale. The risk zone and infarct areas were  
134 traced onto acetate sheets. Using computerised planimetry (Summasketch II, Summagraphics) the percentage of  
135 infarct tissue within the volume of the myocardium at risk was calculated (I/R%).

## 136 **Experimental Groups.**

137 Isolated hearts were split into two groups. Group 1 hearts underwent a protocol for infarct size assessment as  
138 shown in Fig 1a. Group 2 hearts underwent protocols for western blotting as summarised in Fig 1b.

139 **Protocol 1:** All hearts were allowed to stabilise for 20min prior to being subjected to 35min regional ischaemia  
140 followed by 120min reperfusion. Hearts were randomly assigned to the following groups 1min before  
141 reperfusion. a) hearts perfused with KH buffer alone with no drugs(control); b) hearts perfused with 2-CL-IB-  
142 MECA (1nM) throughout the duration of reperfusion, previously shown to be cardioprotective by Maddock et al.  
143 c) hearts perfused with 2-CL-IB-MECA (1nM) in the presence and absence of UO126 (10µM) (highly specific  
144 MEK1/2 inhibitor) or Wortmannin (5nM)(selective irreversible inhibitor of PI3K) throughout the reperfusion  
145 period.

146 **Protocol 2:** Hearts were allowed to stabilise for 20min after which they were randomly assigned to one of the  
147 following groups a) perfused for 60min b) 35min ischaemia with 5,10, or 20min reperfusion in the presence and  
148 absence of 2-CL-IB-MECA (1nM) c) 35min ischaemia followed by 10min reperfusion with 2-CL-IB-MECA in  
149 the presence and absence of U0126 (10µM) or Wortmannin (5nM)d) 35 min ischaemia followed administered at  
150 the onset 10 min reperfusion in the presence and absence of UO126 (10 µM) or Wortmannin (5nM).

#### 151 **Protein Extraction**

152 At the end of each experiment tissues were isolated from the risk zone and snap frozen in liquid nitrogen.  
153 Tissues from the ischaemic zone were homogenised in suspension buffer containing (in mM) 0.1 NaCl, 10 Tris,  
154 1 EDTA (pH8), 2 sodium pyrophosphate, 2 NaF, 2 β-glycerophosphate, 0.1mg/ml 4-(2-Aminoethyl)  
155 benzenesulfonyl fluoride hydrochloride (AEBSF) and protease inhibitor cocktail mixture (Roche, UK) at 4°C  
156 using the IKA-Ultra-Turrax homogeniser. Samples were then centrifuged for 5min at 11,000 rpm and stored at -  
157 80°C until further analysis. Samples were assayed for protein content using the bicinchoninic acid (BCA) protein  
158 assay reagent kit (Pierce,Northumberland,UK).

#### 159 **Western blot analysis**

160 Protein samples were mixed with an equal volume of sample buffer containing 100mM Tris (pH6.8), 200mM  
161 DTT, 2%SDS, 0.2% Bromophenol blue, 20% glycerol and heated to 95°C for 10min and stored at -20°C.  
162 Samples were defrosted on ice and 60µg of protein were separated by SDS-PAGE (12.5% acrylamide gel) using  
163 a Bio-Rad mini protean II system.  
164 Proteins were transferred to polyvinylidenedifluoride (PVDF) membrane using the Bio-Rad mini Trans blot  
165 system. After transfer the membranes were washed with Ambo Life Tris buffered saline (TBS) and blocked for 1  
166 hour using TBS, 5%w/v non fat milk, 0.1% Tween-20. Blots were incubated overnight at 4°C with the primary  
167 antibody to phosphor-AKT(Ser<sup>473</sup>) phospho-ERK1 / ERK2 (Thr<sup>202</sup>/Tyr<sup>204</sup>)at a dilution of 1:1000 in TBS, 5%w/v  
168 bovine serum albumin fraction V, 0.1% Tween-20. Blots were washed and incubated with anti rabbit

169 IgG antibody conjugated to horseradish peroxidase at a concentration of 1:2000 at room temperature for 1 hour.  
170 Blots were further washed and were incubated for a few seconds in enhanced chemiluminescence substrates  
171 (AMBO Lumifast ECL Kit).

172 Blots were exposed to Hyperfilm ECL (Amersham, Buckinghamshire, UK) and developed using Kodak  
173 developing/fixing solution (Sigma, Poole, UK). Equal loading was confirmed by developing the blots for either  
174 total AKT, ERK1/2, antibody. The transfer efficiency was assessed by Ponceau S staining prior to primary  
175 antibody application. Films were scanned and the band densitometry was assessed using the NIH Image J  
176 (v1.33) software.

### 177 **Rat cardiac myocyte model**

178 Adult rat cardiomyocytes were isolated by conventional enzymatic dissociation. Briefly, Male Sprague Dawley  
179 rats (350-500g body weight) were sacrificed by cerebral dislocation. Hearts were quickly excised and mounted  
180 on a modified Langendorff apparatus. Hearts were perfused with modified Krebs Heinsleit bicarbonate buffer  
181 (NaCl 116mM, NaHCO<sub>3</sub> 25.0mM, KCl 5.4mM, MgSO<sub>4</sub> .7.H<sub>2</sub>O 0.4mM, CaCl<sub>2</sub> 1.7mM, glucose 10mM, taurine  
182 20mM, pyruvate 5mM and Na<sub>2</sub>HPO<sub>4</sub>.12H<sub>2</sub>O 0.9mM). The buffer was oxygenated with 95% O<sub>2</sub> and 5% CO<sub>2</sub>  
183 (BOC Gases) and heated to 37°C (pH 7.4).

184 Hearts were perfused for 5 minutes with Krebs buffer containing calcium followed by 5 minutes of perfusion  
185 with calcium free Krebs Heinsleit buffer. Upon perfusion with calcium free buffer the heart ceased to contract.  
186 The hearts underwent a final 5 minute perfusion cycle with modified Krebs Heinsleit digestion buffer containing  
187 BSA 0.5%, Worthingtons Type II Collagenase 0.075%, CaCl<sub>2</sub> 4.4µM pH 7.4 with NaOH. During perfusion with  
188 collagenase the effluent was collected and reused.

189 After perfusion with collagenase the heart was removed and the atria were cut away. The ventricles were teased  
190 apart and incubated with fresh digestion buffer for 10 minutes on an orbital shaker. The digestion buffer was  
191 aspirated and was passed through a nylon mesh into a sterile falcon tube and centrifuged at 400 rpm for 2  
192 minutes.

193 The supernatant was removed using a sterile pipette and the pellet was redistributed in 25ml freshly prepared  
194 restoration buffer (NaCl 116mM, NaHCO<sub>3</sub> 25.0mM, KCl 5.4mM, MgSO<sub>4</sub> .7.H<sub>2</sub>O 0.4mM, glucose 10mM,  
195 taurine 20mM, pyruvate 5mM Na<sub>2</sub>HPO<sub>4</sub>.12H<sub>2</sub>O 0.9mM, 1% BSA and 1% Pen-Strep). The calcium  
196 concentration was gradually brought back to 1.25mM. Myocyte viability was assessed using trypan blue and  
197 were discarded if the viability was below 70% or myocytes were incubated in suspension buffer for 24 hours  
198 before being used in experiments.



199 **Experimental Protocol.**

200 Cardiac myocytes were centrifuged at 500rpm for 5 min and buffer was replaced with hypoxic buffer (12 mM  
201 KCL, 0.49mM MgCl<sub>2</sub>, 0.9 mM CaCl<sub>2</sub>, 4 mM HEPES, 10 mMdeoxyglucose and 20 mMlactate). Myocytes were  
202 incubated in a pre-heated (37°C) hypoxic chamber for 6 hours where the O<sub>2</sub> was replaced with 5% CO<sub>2</sub> balanced  
203 in Argon (BOC Gases). After hypoxia, cells were removed from the chamber and centrifuged at 500 rpm for 5  
204 minutes. The supernatant was removed and replaced with restoration buffer.The A<sub>3</sub> agonist 2-Cl-IB-MECA  
205 (1nM) and the MEK1/2 inhibitor UO126 (10µM) or PI3K inhibitor Wortmannin (5nM) were placed into the  
206 appropriate wells in the 24 well plate. Cells were also kept for the normoxic group and unstained group.  
207 Cells were reoxygenated in the incubator for 18 hours before undergoing fluorescence-activated cell sorting  
208 (FACS) analysis for assessment of cellular apoptosis , necrosis, cleaved-caspase3 and p-BAD (Ser<sup>112</sup>) and p-  
209 BAD (Ser<sup>136</sup>) activity.

210 **Assessment of apoptosis and necrosis by FACS Analysis**

211 The Vybrant Apoptosis Assay Kit #10 (Invitrogen, Paisley,UK) provides a three-colour fluorescence assay that  
212 distinguishes live, apoptotic, and late apoptotic cells from one another. Cells were incubated with fluorochromes  
213 according to the manufacturer's instructions. Briefly, cells were pelleted by centrifugation and resuspended in  
214 Annexin V buffer. The appropriate volumes of fluorescent probes were added to the cells and incubated at 37°C  
215 for 15 minutes. The samples were analysed on the FL-2 and FL-4 channels on the BD FACS Calibur® Flow  
216 cytometer and setup to count 10,000 events. Quantitative assessment of live, apoptotic and necrotic cell  
217 populations was made using the Cell Quest software.

218 **Assessment of p-BAD (Ser<sup>112</sup>) and p-BAD (Ser<sup>136</sup>) and cleaved-caspase 3 activity by FACS Analysis**

219 To determine the activity of p-BAD (Ser<sup>112</sup>), p-BAD (Ser<sup>136</sup>) and cleaved-caspase 3 in adult rat myocytes at the  
220 end of reoxygenation cells were centrifuged at 1200rpm for 2min(33). The supernatant was aspirated and the  
221 pellet was resuspended in 250µl phosphate buffered saline andfixed by the addition of 250µl of 6%  
222 formaldehyde. The cells were incubated for 10min at 37°C and placed on ice for 1 min.

223 The eppendorffs were centrifuged at 1200 rpm for 2min after which the supernatant was aspirated and the cells  
224 were permeabilised by resuspending in ice-cold methanol (90%). Cells were kept on ice for 30min and  
225 centrifuged at 1200 rpm for 2min. The supernatant was removed and the cells were with incubation buffer (0.5%  
226 BSA in PBS stored @ 4°C). The eppendorffs were centrifuged at 1200 rpm for 2min followed by the removal of  
227 the supernatant. The cells were blocked by the addition of 100µl of incubation buffer for 10 minutes at room  
228 temperature. The cleaved-Caspase 3<sub>ASP175</sub> (5A1) rabbit monoclonal primary antibody or p-BAD (ser112) (New

229 England Biolabs, Hitchin, UK) was added to the blocking buffer to give a final concentration of 1:100 and  
1 230 incubated at room temperature for 60 minutes.

2 231 Theeppendorffs were centrifuged again at 1200 rpm for 2 minutes followed by the removal of supernatant and  
3  
4 232 washed twice in incubation buffer. The cells were resuspended in 200µl of incubation buffer containing the  
5  
6 233 Alexa Fluor® 488 secondary antibody (Invitrogen®) to give a final concentration of 1:1000. The samples were  
7  
8 234 covered in foil and incubated for 30 minutes at room temperature. The cells were centrifuged at 1200 rpm for 2  
9  
10 235 min and washed in incubation buffer. Finally, the cells were resuspended in 500µl of phosphate buffered saline  
11  
12 236 and analysed on the BD FACS Calibur® flow cytometer on the FL-1 channel. Histograms were plotted for each  
13  
14 237 of the groups showing the mean fluorescence for 10,000 counts.

### 18 238 **Statistical Analysis**

19  
20 239 All values are expressed as mean ±SEM. Infarct size, cell death and band densities were tested for group  
21  
22 240 differences using one way analysis of variance (ANOVA) with Fishers post hoc tests. A P-value of P<0.05 was  
23  
24 241 considered statistically significant. Left ventricular developed pressure, heart rate and coronary flow were  
25  
26 242 assessed for statistical difference using a Two way ANOVA for repeated measures.

### 28 243 **Exclusion Criteria**

29  
30 244 In total 5 hearts were excluded from the study where heart rate or left ventricular developed pressure failed to  
31  
32 245 recover during the stabilisation phase.

34 246

### 36 247 **Results**

#### 38 248 *Homogeneity among the experimental groups*

39  
40 249 The results from 68successful experiments were included. Heart rate, coronary flow and left ventricular  
41  
42 250 developed pressure were not significantly different during stabilisation, ischaemia and reperfusion in all groups  
43  
44 251 (data not included).

#### 46 252 *Protection afforded by the A<sub>3</sub>AR agonist 2-CL-IB-MECA when administered at reperfusion is mediated via* 48 49 253 *MEK1/2 activity*

50  
51 254 Isolated perfused rat hearts underwent 35 minutes of ischaemia followed by 120min of reperfusion where the A<sub>3</sub>  
52  
53 255 agonist 2-CL-IB-MECA (1nM) was administered throughout the reperfusion period. Our studies showed that the  
54  
55 256 administration of the highly specific A<sub>3</sub>AR agonist 2-Cl-IB-MECA (1nM) throughout the reperfusion period  
56  
57 257 significantly attenuated infarct development (% of the risk area) (I/R %) from 65 ± 1.7% in control hearts to 32

258 ±4% in 2-CL-IB-MECA treated hearts, a reduction of 51% (P<0.01, Fig.2). The cardioprotective effect of 2-CL-  
1 IB-MECA has previously been demonstrated by Maddock et al.(8).

259  
3  
4 260 To determine whether the protection afforded by the A<sub>3</sub>AR agonist 2-Cl-IB-MECA (1 nM) when administered at  
5  
6 261 reperfusion was mediated via MEK1/2 activity, we used the MEK1/2 inhibitor UO126 (10 μM). Alone, UO126  
7  
8 262 had no significant effect on infarct size compared to control (55 ± 8% vs. 65 ± 1.5% control P>0.05, Fig.2).

10 263 Co-administration of UO126 in the presence of 2-CL-IB-MECA significantly abrogated the protection afforded  
11  
12 264 by 2-CL-IB-MECA (59 ± 9% vs. 32 ± 4% 2-CL-IB-MECA P<0.05, Fig.2). These data suggest that MEK1/2  
13  
14 265 dependent signalling pathways are involved in A<sub>3</sub>AR mediated cardioprotection when administered at  
15  
16 266 reperfusion.

18  
19 267 ***Protection afforded by the A<sub>3</sub>AR agonist 2-CL-IB-MECA at reperfusion is mediated via PI3K activity***

20 268 To determine whether the protection afforded by the A<sub>3</sub>AR agonist 2-Cl-IB-MECA (1 nM) when administered at  
21  
22 269 reperfusion was mediated by enhanced PI3K activity, we used the PI3K inhibitor Wortmannin (5nM). Alone,  
23  
24 270 Wortmannin had no significant effect on infarct size compared to control (65 ± 8% vs. 65 ± 2% respectively,  
25  
26 271 P>0.05, Fig.3).

28  
29 272 Co-administration of Wortmannin with 2-CL-IB-MECA significantly abrogated the protection afforded by 2-  
30  
31 273 CL-IB-MECA (68 ± 12% vs. 32 ± 4% P<0.05; respectively, Fig.3). These data suggest that PI3K dependent  
32  
33 274 signalling pathways are also involved in A<sub>3</sub>AR mediated cardioprotection when administered at reperfusion.

34  
35 275 ***The role of MEK1/2 – ERK1/2 MAPK pathway in A<sub>3</sub> adenosine receptor mediated cardioprotection in adult***  
36  
37 276 ***rat cardiomyocytes subjected to hypoxia/ reoxygenation injury***

38  
39 277 As shown in Figure 4 and 6, isolated adult rat cardiac myocytes underwent 6 hours of hypoxia followed by 18  
40  
41 278 hours of reoxygenation resulting in a 3.2 fold increase in the number of apoptotic cells (41.8 ± 5.4% vs. 12.9 ±  
42  
43 279 1.3% normoxic group, P<0.001), and a 3.1 fold increase in the number of necrotic cells (33.2 ± 2 vs. 10.7 ±2.1%  
44  
45 280 normoxic group, P<0.001, Fig 4).

46  
47 281 To ascertain the role of A<sub>3</sub>ARs in limiting the deleterious consequences of reoxygenation injury the A<sub>3</sub>AR  
48  
49 282 agonist 2-Cl-IB-MECA (1 nM) was administered throughout the reoxygenation period. Administration of the A<sub>3</sub>  
50  
51 283 agonist 2-Cl-IB-MECA (1 nM) significantly decreased the percentage of apoptotic myocytes (26 ± 6% vs. 41.8 ±  
52  
53 284 5.4 % Hyp/Reox group, P<0.001, Fig 4), and necrotic myocytes (16 ± 4 vs. 33.2 ± 2 Hyp/Reox group, P<0.001,  
54  
55 285 Fig 4).

56  
57 286 Administration of the A<sub>3</sub> adenosine receptor agonist 2-Cl-IB-MECA (1 nM) at reoxygenation significantly  
58  
59 287 decreased the number of apoptotic and necrotic myocytes compared to non-treated groups. To determine which  
60  
61  
62  
63  
64  
65

288 mechanisms are involved in 2-CL-IB-MECA mediated cardioprotection we assessed the role of the MEK1/2 –  
1 ERK1/2 cell survival pathway using the MEK1/2 inhibitor UO126 (10  $\mu$ M). Cardiac myocytes were incubated  
2 289  
3  
4 290 with the MEK1/2 inhibitor UO126 (10  $\mu$ M) in the presence and absence of the A<sub>3</sub> agonist 2-CL-IB-MECA (1  
5  
6 291 nM).

7  
8 292 Co-administration of the A<sub>3</sub> agonist 2-CL-IB-MECA (1 nM) with the MEK 1/2 inhibitor UO126 (10  $\mu$ M)  
9  
10 293 significantly abolished the anti-apoptotic (38  $\pm$  4 % vs. 26  $\pm$  6 % 2-CL-IB-MECA, P<0.001) and anti-necrotic  
11  
12 294 potential of 2-Cl-IB-MECA (25  $\pm$  3 vs. 16  $\pm$  4 2-CL-IB-MECA, P<0.001, Fig 4 ). UO126 alone had no  
13  
14 295 significant effect of cellular apoptosis when compared to the hypoxic reoxygenated group (42  $\pm$  6 % vs. 42  $\pm$  5  
15  
16 296 % Hyp/Reox, P>0.05, Fig 4). UO126 when administered at reoxygenation was seen to significantly decrease the  
17  
18 297 number of cells dying by cellular necrosis when compared the hypoxic reoxygenated group (23  $\pm$  3 % vs. 32.2 %  
19  
20 298 vs. Hyp/Reox, P<0.05, Fig 4).

21  
22 299 ***The role of PI3K survival pathway in A<sub>3</sub> adenosine receptor mediated cardioprotection in adult rat***  
23  
24 300 ***cardiomyocytes subjected to hypoxia/ reoxygenation injury.***

25 301 Administration of the A<sub>3</sub> adenosine receptor agonist 2-Cl-IB-MECA (1 nM) at reoxygenation significantly  
26  
27 302 decreased the number of apoptotic and necrotic myocytes compared to non-treated groups. To determine which  
28  
29 303 cellular mechanisms are involved in 2-CL-IB-MECA mediated cytoprotection we assessed the role of the PI3K  
30  
31 304 cell survival pathway using the PI3K inhibitor Wortmannin (5 nM).

32  
33 305 Administration of the A<sub>3</sub> agonist 2-CL-IB-MECA (1 nM) in the presence of the PI3K inhibitor Wortmannin (5  
34  
35 306 nM) significantly abolished the anti-apoptotic effect of 2-CL-IB-MECA (1 nM) compared to when administered  
36  
37 307 alone throughout reoxygenation (37  $\pm$  6 % 2-CL-IB-MECA + Wortmannin vs. 26  $\pm$  6 % 2-CL-IB-MECA,  
38  
39 308 P<0.05, Fig 5).

40  
41 309 Administration of 2-CL-IB-MECA (1 nM) in the presence of the PI3K inhibitor Wortmannin (5 nM) also  
42  
43 310 significantly abolished the anti-necrotic effect of 2-CL-IB-MECA (1 nM) when administered alone throughout  
44  
45 311 reoxygenation (28  $\pm$  3 % 2-CL-IB-MECA + Wortmannin vs. 16  $\pm$  4 % 2-CL-IB-MECA, P<0.05, Fig 5).

46  
47 312 Wortmannin (5 nM) alone had no significant effect of cellular apoptosis when compared to the Hyp/Reox group  
48  
49 313 (36  $\pm$  5% Wortmannin vs. 42  $\pm$  5 % Hyp/Reox, P>0.05) Fig 5. Interestingly, Wortmannin (5 nM) when  
50  
51 314 administered alone throughout reoxygenation was seen to have a significant effect on reducing cellular necrosis  
52  
53 315 when compared to the Hyp/Reox group (22  $\pm$  3% Wortmannin vs. 32  $\pm$  2 % Hyp/Reox P<0.001, Fig 5).

54 316 ***Time course of ERK 1/2 phosphorylation in non treated and 2-CL-IB-MECA ischaemic reperfused rat hearts***

317 In order to investigate the role of MAPKs ERK1/2 in A<sub>3</sub>AR mediated cardioprotection 2-CL-IB-MECA (1 nM)  
318 was administered at reperfusion and the heart tissues were harvested at different time intervals in both treated  
319 and non-treated groups and assessed by western blot analysis for total and Phospho-ERK1/2.

320 Administration of the A<sub>3</sub> agonist 2-CL-IB-MECA (1nM) at reperfusion significantly upregulated the  
321 phosphorylation of ERK1/2 after 10 minutes of reperfusion (P<0.05; Fig 6) compared to non-treated time  
322 matched controls. This upregulation of ERK 1/2 phosphorylation by 2-CL-IB-MECA (1nM) after 10 minutes of  
323 reperfusion was abrogated by the co-administration of the MEK1/2 inhibitor UO126 (10µM) (P<0.01; Fig 6).  
324 Administration of UO126 (10µM) for 10 minutes of reperfusion had no significant effect of p-ERK expression  
325 compared to time matched controls (P>0.05; Fig 6). Administration of Wortmannin alone or in the presence of  
326 2-CL-IB-MECA had no significant effect p-ERK levels.

327 ***Time course of AKT<sub>(ser 473)</sub> phosphorylation in non-treated and 2-CL-IB-MECA treated ischaemic reperfused***  
328 ***rat hearts.***

329 In order to determine the role of the PI3K/AKT pro-survival pathway in A<sub>3</sub>AR mediated cardioprotection we  
330 assessed the phosphorylation status of p-AKT<sub>(Ser 473)</sub>, at various time intervals post reperfusion in the presence  
331 and absence of the A<sub>3</sub> agonist 2-CL-IB-MECA (1 nM).

332 Administration of 2-CL-IB-MECA (1nM) for 10 minutes of reperfusion significantly upregulated the  
333 phosphorylation of AKT<sub>(ser473)</sub> compared totime matched controls (P<0.01 vs. time matched controls, Fig 7).2-  
334 CL-IB-MECA dependent increase in p-AKT at 10 minutes of reperfusion was significantly reversed in the  
335 presence of the PI3K Inhibitor Wortmannin (P<0.001;Fig 7).Administration of Wortmannin (5nM) for 10  
336 minutes of reperfusion significantly decreased p-AKT expression compared to time matched controls (P<0.001;  
337 Fig 7). Administration of UO126 alone had no significant effect p-AKT compared to time matched control  
338 (P>0.05; Fig 7). Administration of UO126 + 2CL-IB-MECA significantly increased p-AKT compared to time  
339 matched control (P<0.01; Fig 7).

340 ***Role of 2-CL-IB-MECA when administered at reoxygenation on p-BAD (ser112)and p-BAD (ser136) activity***  
341 ***in isolated rat cardiomyocytes subjected to 6 hours of hypoxia and 18 hours of reoxygenation***

342 Recruitment of pro-survival pathways has been shown to inhibit by phosphorylation downstream pro-apoptotic  
343 protein such as BAD. BADis involved in initiating apoptosis, but when phosphorylated at Ser112 and Ser136  
344 BAD is inactivated and apoptosisisreduced. In order to investigate whether the cardioprotection observed by the  
345 administration of 2-CL-IB-MECA involved phosphorylation of p-BAD (Ser<sup>112</sup>) and p-BAD (Ser<sup>136</sup>),cardiac  
346 myocytes were subjected to hypoxia and reoxygenation injury where 2-CL-IB-MECA (1nM) was administered  
347 at reoxygenation in the presence and absence of the kinase inhibitors.

348 Administration of the A<sub>3</sub> agonist 2-Cl-IB-MECA (1 nM) at reoxygenation significantly up regulated the  
1  
2 349 phosphorylation of BAD<sub>ser112</sub> and BAD<sub>ser136</sub> compared to the Hyp/Reox group (P<0.05) (Fig 8). Co-  
3  
4 350 administration of 2-CL-IB-MECA (1nM) with UO126 (10uM) or Wortmannin (5nM) significantly abolished 2-  
5  
6 351 CL-IB-MECA (1nM) mediated upregulation of p-BAD(ser112) and p-BAD (Ser136) respectively (P<0.05) (Fig  
7  
8 352 8). The kinase inhibitors Wortmannin or UO126 alone had no significant effect on p-BAD(ser112) and p-BAD  
9  
10 353 (Ser136) compared to the Hyp/Reox group (Fig 8).

11  
12 354 ***Role of 2-CL-IB-MECA when administered at reoxygenation on cleaved-caspase 3 activity in isolated rat***  
13  
14 355 ***cardiomyocytes subjected to 6 hours of hypoxia and 18 hours of reoxygenation***

15  
16 356 Isolated adult rat cardiomyocytes subjected to 6 hours of hypoxia followed by 18 hours of reoxygenation  
17  
18 357 resulted in a 2.5 fold increase in cleaved-caspase 3 activity compared to the normoxic group P<0.001 (Fig 9).  
19  
20 358 Administration of the A<sub>3</sub> agonist 2-Cl-IB-MECA (1nM) at reoxygenation significantly decreased Hyp/Reox  
21  
22 359 dependent increases in cleaved-caspase 3 activity (P<0.001, Fig 9). The data strongly suggests that the  
23  
24 360 cardioprotective effects of 2-CL-IB-MECA are via anti-apoptotic mechanisms.

25  
26 361 ***Role of MEK1/2-ERK1 and PI3K 2 cell survival pathway in cleaved-caspase 3 activity upon administration of***  
27  
28 362 ***2-CL-IB-MECA (1nM) at reoxygenation in isolated myocytes subjected to hypoxia/reoxygenation injury.***

29  
30 363 Administration of 2-Cl-IB-MECA (1nM) at reoxygenation significantly reduced cleaved-caspase 3 activity  
31  
32 364 compared to non-treated isolated adult rat cardiomyocytes subjected to 6 hours of hypoxia and 18 hours of  
33  
34 365 reoxygenation (P<0.001, Fig 9). To determine whether the decrease in cleaved - caspase 3 activity involved the  
35  
36 366 MEK1/2 – ERK1/2 cell survival pathway we used the MEK1/2 inhibitor UO126 (10µM) or PI3K inhibitor  
37  
38 367 Wortmannin (5nM) in the presence and absence of 2-C-IB-MECA (1nM).

39  
40 368 Administration of the MEK1/2 inhibitor UO126 with the A<sub>3</sub> agonist 2-CL-IB-MECA (1nM) at reoxygenation  
41  
42 369 failed to block the decrease in cleaved-caspase 3 activity afforded when 2-CL-IB-MECA was administered alone  
43  
44 370 at reoxygenation (P>0.05, Fig 9). UO126 when administered alone had no significant effect on cleaved-caspase  
45  
46 371 3 activity compared to the Hyp/Reox group (P>0.05, Fig 9).

47  
48 372 Activation of A<sub>3</sub>ARs throughout reoxygenation significantly decreased cleaved-caspase 3 activity which was  
49  
50 373 significantly abolished in the presence of the PI3K inhibitor Wortmannin (5 nM) (P<0.05, Fig 10).

51  
52 374 Administration of Wortmannin (5nM) throughout reoxygenation alone had no significant effect of cleaved-  
53  
54 375 caspase 3 activity compared to the non-treated Hyp/Reox (P>0.05, Fig 10).

55  
56 376  
57  
58 377 **Discussion**

378 The findings from the current study demonstrate that activation of adenosine A<sub>3</sub> receptors at reperfusion protects  
1  
2 379 the isolated perfused rat heart from ischaemia/reperfusion injury and cardiac myocytes from  
3  
4 380 hypoxia/reoxygenation injury via recruitment of the survival kinases PI3K and MEK1/2.  
5  
6 381 Our data are in agreement with those of Maddock and others which have previously shown that activation of  
7  
8 382 A<sub>3</sub>AR at reperfusion significantly attenuated infarct development when administered during reperfusion (7, 8, 10,  
9  
10 383 13, 14). The selective A<sub>3</sub>AR antagonist I-ABOPX has previously been co-administered with the A<sub>3</sub>AR agonist 2-  
11  
12 384 CL-IB-MECA confirming that the protection was directly attributed to A<sub>3</sub>ARs and not other adenosine receptor  
13  
14 385 subtypes (14). 2-CL-IB-MECA dependent cerebral protection has also been described in the gerbil model of  
15  
16 386 ischaemia / reperfusion injury leading to neuronal protection and decreased microglial infiltration (23). 2-CL-IB-  
17  
18 387 MECA has been shown to protect against simulated models of ischaemia reperfusion injury in the brain (34) and  
19  
20 388 feline lung (19, 21).  
21  
22 389 Our results confirm those of Maddock and colleagues showing 2-CL-IB-MECA (1nM) attenuated apoptosis and  
23  
24 390 necrosis cell death in cardiac myocytes subjected to 6 hours of hypoxia and 18 hours of reoxygenation (8).  
25  
26 391 Indeed, activation of A<sub>3</sub>ARs has been shown to protect rat basophilic leukaemia mast cells exposed to UV light  
27  
28 392 reducing cellular apoptosis by 50% (24).  
29  
30 393 MAPKs play an essential role in myocardial ischaemia reperfusion injury (35, 36). Adenosine A<sub>3</sub> receptors have  
31  
32 394 been shown to activate MAPKs ERK1/2 in Chinese hamster ovary cells exhibiting the human A<sub>3</sub> adenosine  
33  
34 395 receptor (37, 38). Our data revealed that the infarct sparing effects associated with 2-CL-IB-MECA was via  
35  
36 396 recruitment of the reperfusion injury salvage kinase MEK1/2 – ERK 1/2 cell survival pathway in the isolated  
37  
38 397 perfused rat heart. These results are in accordance with other studies showing that activation of A<sub>3</sub>ARs at  
39  
40 398 reperfusion can limit the deleterious consequences of ischaemia reperfusion injury in a number of organs  
41  
42 399 including the heart (8, 13-15, 39), lung (19, 21) and brain (23). Our studies show that co-administration of 2-  
43  
44 400 CL-IB-MECA in the presence of the MEK1/2 inhibitor UO126 abolished A<sub>3</sub> agonist dependent cardioprotection  
45  
46 401 in the isolated heart. In concurrence with other cardioprotective agents, previous studies have shown  
47  
48 402 cardioprotection with erythropoietin or adipocytokine visfatin protects against ischaemic injury via activation of  
49  
50 403 MEK1/2, where inhibition with UO126 an inhibitor of MEK1/2 abolished protection (40, 41).  
51  
52 404 Our data showed that the administration of 2-CL-IB-MECA at reoxygenation significantly decreased the number  
53  
54 405 of apoptotic and necrotic cells compared to non-treated cells. Interestingly, the MEK1/2 inhibitor when  
55  
56 406 administered together with 2-CL-IB-MECA abolished 2-CL-IB-MECA-dependent anti apoptotic/necrotic  
57  
58 407 effects. Our results are in accordance with numerous studies that have shown activation of A<sub>3</sub>ARs to protect  
59  
60  
61  
62  
63  
64  
65

408 myocytes from hypoxia reoxygenation injury (32, 42). They further showed that the cytoprotection was  
1  
2 409 abolished in the presence of the A<sub>3</sub>AR antagonist MRS 1191 implicating a direct role for A<sub>3</sub>ARs. Matot and  
3  
4 410 colleagues have shown the A<sub>3</sub> agonist 2-CL-IB-MECA when administered at reperfusion attenuated ischaemia  
5  
6 411 reperfusion injury in the in vivo feline lung via anti apoptotic/necrotic signalling pathways (19). They further  
7  
8 412 showed that the A<sub>3</sub> antagonist MRS 1191 abolished this protective effect. A number of groups have shown  
9  
10 413 activation of A<sub>3</sub>ARs to protect cells from ischaemic injury (23, 24). This is the first study to show that activation  
11  
12 414 of adenosine A<sub>3</sub> receptors at reperfusion/reoxygenation to activate the MEK1/2-ERK1/2-BAD cell survival  
13  
14 415 pathways in isolated hearts and cardiomyocytes.  
15  
16 416 Our results support those of Germack and Dickenson who showed preconditioning with adenosine protected  
17  
18 417 myocytes from hypoxia reoxygenation injury measured by decreased caspase 3 releases and lactate  
19  
20 418 dehydrogenase release(32). They further showed this cardioprotection was associated with recruitment of  
21  
22 419 MEK1/2-ERK1/2 signalling pathways where inhibition of MEK1 with PD98059 completely abolished  
23  
24 420 adenosine-dependent decreases in LDH and caspase 3.  
25  
26 421 Our data showed that 2-Cl-IB-MECA when administered at reperfusion significantly upregulated ERK1/2  
27  
28 422 phosphorylation at the time points assessed compared to time matched controls in a UO126 sensitive manner.  
29  
30 423 Our results confirm those of Dickenson and colleagues who showed recruitment of the MEK1/2 – ERK1/2 cell  
31  
32 424 survival pathway in neonatal myocytes preconditioned with 2-CL-IB-MECA (32). Importantly, Matot and  
33  
34 425 colleagues showed that the administration of the A<sub>3</sub> agonist IB-MECA or the newly synthesised A<sub>3</sub> agonist  
35  
36 426 MRS3558 at the onset of reperfusion significantly protected the feline ischaemic lung from reperfusion  
37  
38 427 injury(19). They further showed upregulation of ERK1/2 phosphorylation with the A<sub>3</sub> agonists where ERK1/2  
39  
40 428 phosphorylation was abolished in the presence of the A<sub>3</sub> antagonist MRS 1191(19).  
41  
42 429 Our findings showed 2-CL-IB-MECA mediated infarct sparing effects were abolished in the presence of the  
43  
44 430 PI3K inhibitor Wortmannin in the isolated perfused heart. 2-CL-IB-MECA mediated protection involved  
45  
46 431 recruitment of PI3K /AKT pathway, where the PI3K inhibitor Wortmannin abolished 2-CL-IB-MECA mediated  
47  
48 432 protection. Similarly, Park and colleagues have shown that the protection afforded by the administration of 2-  
49  
50 433 CL-IB-MECA (1µM) at reperfusion involves recruitment of the PI3K/AKT survival pathway as inhibition of this  
51  
52 434 pathway with the PI3K inhibitor Wortmannin abolished 2-CL-IB-MECA mediated protection in the isolated  
53  
54 435 perfused rat heart (43). Our findings may be explained as 2-CL-IB-MECA has been shown to activate AKT in a  
55  
56 436 dose dependent manner, where 2-CL-IB-MECA increases AKT phosphorylation (31). We show for the first time  
57  
58 437 that the anti-apoptotic and anti-necrotic effects of 2-CL-IB-MECA were dependent on PI3K signalling where the  
59  
60  
61  
62  
63  
64  
65



438 cytoprotective effects were reversed in the presence of the PI3K inhibitor Wortmannin in cardiac myocytes  
1  
2 439 subjected to hypoxia reoxygenation injury.

3  
4 440 Recruitment of the MEK1/2-ERK1/2 and PI3K/AKT pathways has been shown to recruit downstream effector  
5  
6 441 proteins including the pro-apoptotic protein BAD(41, 44, 45). BAD remains in the cytosol when phosphorylated  
7  
8 442 on either of its phosphorylation sites BAD<sub>ser 112</sub> and BAD<sub>ser 136</sub> bound to 14-3-3. In its phosphorylated state BADs  
9  
10 443 pro-apoptotic potential is diminished and upon dephosphorylation BAD translocates to mitochondria where it  
11  
12 444 leads to the release of cytochrome c from the mitochondria initiating the apoptosis(46). Our results indicate that  
13  
14 445 activation of A<sub>3</sub>ARs at reperfusion upregulates BAD (Ser<sub>112</sub>) and BAD (ser136) phosphorylation thereby  
15  
16 446 limiting cellular apoptosis.

17  
18 447 Caspase 3, a marker of cellular apoptosis, undergoes a conformational change from its inactive form to its active  
19  
20 448 cleaved form upon receiving apoptotic signals(47). Administration of 2-CL-IB-MECA at the onset of  
21  
22 449 reoxygenation reversed the increase in cleaved-caspase 3 observed in the non-treated Hyp/Reox group. These  
23  
24 450 findings further support our findings that activation of A<sub>3</sub>ARs protects the myocardium from  
25  
26 451 reperfusion/reoxygenation injury in the isolated rat heart/myocyte in an anti-apoptotic manner.

27  
28 452 To determine whether the decrease in cleaved-caspase 3 was dependent on the recruitment of cell survival  
29  
30 453 kinases MEK1/2-ERK1/2 myocytes were incubated with 2-CL-IB-MECA in the presence and absence of the  
31  
32 454 selective MEK1/2 inhibitor UO126. Our data showed that 2-CL-IB-MECA dependent decreases in caspase 3  
33  
34 455 were ameliorated by the PI3K inhibitor Wortmannin and not by the MEK1/2 inhibitor UO126. Previous studies  
35  
36 456 have shown that preconditioning with adenosine (non-specific adenosine receptor agonist) or 2-CL-IB-MECA  
37  
38 457 (highly specific A<sub>3</sub>AR agonist) decreased caspase 3 activity in neonatal cardiac myocytes subjected to 4 hours of  
39  
40 458 hypoxia and 18 hours of reoxygenation(32).

41  
42 459 We have shown that 2-CL-IB-MECA when administered at reoxygenation protects cardiac myocytes from  
43  
44 460 hypoxia/reoxygenation injury via anti-apoptotic pathways. This protective anti-apoptotic effect of 2-CL-IB-  
45  
46 461 MECA is supported in a number of diverse models of ischaemia/reperfusion including the lung (19-21), heart (8)  
47  
48 462 and brain (23). Furthermore, 2-CL-IB-MECA has also been implicated in reducing cleaved-caspase 3 activity in  
49  
50 463 the ischaemic/reperfused lung (19, 20).

51  
52 464 Numerous studies have implicated growth factors, hormones, cytokines and neurotrophic factors to protect  
53  
54 465 against ischaemia reperfusion injury via recruitment of the MEK1/2-ERK1/2/BAD and PI3K/AKT/BADcell  
55  
56 466 survival pathway(28). Urocortin a hormone related to the hypothalamic corticotrophin releasing factor hormone  
57  
58 467 has been shown to protect the ischaemic myocardium from ischaemia reperfusion injury via upregulation of  
59  
60  
61  
62  
63  
64  
65

468 ERK1/2 in a PD98059 (MEK1/2 inhibitor) sensitive manner as has visfatinanadipocytokine(41, 48, 49). Other  
1 469 studies have shown insulin and Neuregulin-1 mediated cytoprotection involved recruitment of the  
2  
3 470 PI3K/AK/BAD cell survival pathway (50,51).Previous studies have shown that the cAMP Response Element-  
4  
5 471 Binding Protein is downstream target of the PI3K and MAPK cell survival pathway and may play in A<sub>3</sub>AR  
6  
7 472 mediated cardioprotection via activation of the transcription factor CREB, which induces the expression of the  
8  
9 473 antiapoptotic factor Bcl-2 (52, 53). Furthermore, studies have shown that A<sub>3</sub>AR induced cardioprotection  
10  
11 474 involves ATP-sensitive potassium channels (K<sub>ATP</sub>) indicating their role A<sub>3</sub>AR mediated protection (54). It is  
12  
13 475 feasible that a number of pathways converge including the PI3K/AKT, MEK1-2/ERK1/2/ K<sub>ATP</sub> channels  
14  
15 476 mediating 2-CL-IB-MECA induced cardioprotection(17, 18).

16 477 Interestingly, Ge and colleagues have shown A<sub>3</sub>AR mediated cardioprotection to be dependent on activation of  
17  
18 478 A<sub>3</sub>AR on bone marrow cells via supressing inflammatory reactions in an *in vivo* model of ischaemia/reperfusion  
19  
20 479 injury (55).The current in-vitro study shows 2-CL-IB-MECA to mediate protection for ischaemia/ reperfusion in  
21  
22 480 a cell free model. Our findings may be explained as A<sub>3</sub>ARs have been shown to functionally present on  
23  
24 481 cardiomyocytes where upon activation elicit protection via cell survival signalling pathways.

25  
26 482 The A<sub>1</sub>/A<sub>2</sub> AR agonist 5'-(N-ethylcarboxamido) adenosine (NECA) has been shown to protect the ischaemic  
27  
28 483 reperfused rabbit heart from reperfusion injury when administered at reperfusion via upregulation of ERK1/2  
29  
30 484 whereby inhibition of MEK1 with PD98059 blocked the protection(56).  
31  
32  
33

34 485  
35  
36 486 In conclusion we show for the first time that 2-CL-IB-MECA protects the ischaemic reperfused rat heart from  
37  
38 487 ischaemia reperfusion injury when administered at reperfusion via recruitment of the MEK1-2/PI3K cell survival  
39  
40 488 pathway. Furthermore, activation of A<sub>3</sub>ARs on the onset of reoxygenation significantly protected the hypoxic /  
41  
42 489 reoxygenated adult rat cardiac myocyte from reoxygenation injury in an anti-apoptotic/necrotic manner via the  
43  
44 490 MEK1/2 - PI3K cell survival pathway. Finally, 2-CL-IB-MECA reduced caspase 3 activityin a PI3K dependent  
45  
46 491 and MEK1/2 independent manner. The potential pathways involved in A<sub>3</sub>AR mediated cardioprotection are  
47  
48 492 summarised in Fig 11.  
49

50 493

#### 51 494 **Acknowledgements**

52  
53  
54 495 This work was funded by the British Heart Foundation PhD scholarship grantFS/03/018/15145.  
55

#### 56 496 57 497 **Conflicts of Interest**

58 498  
59 499 The authors declare that they have no conflict of interest.  
60  
61 500  
62  
63  
64  
65

501 **References:**

- 1 502 1. Van Wylen DG, Schmit TJ, Lasley RD, Gingell RL, Mentzer RM, Jr. Cardiac  
2 503 microdialysis in isolated rat hearts: interstitial purine metabolites during ischemia. *Am J*  
3 504 *Physiol.* 1992;262(6 Pt 2):H1934-8.
- 4 505 2. Tucker AL, Linden J. Cloned receptors and cardiovascular responses to adenosine.  
5 506 *Cardiovasc Res.* 1993;27:62-7.
- 6 507 3. Ely SW, Mentzer RM, Jr., Lasley RD, Lee BK, Berne RM. Functional and metabolic  
7 508 evidence of enhanced myocardial tolerance to ischemia and reperfusion with adenosine. *J*  
8 509 *Thorac Cardiovasc Surg.* 1985;90:549-56.
- 9 510 4. Olafsson B, Forman MB, Puett DW, et al. Reduction of reperfusion injury in the canine  
10 511 preparation by intracoronary adenosine: importance of the endothelium and the no-reflow  
11 512 phenomenon. *Circulation.* 1987;76:1135-45.
- 12 513 5. Toombs CF, McGee S, Johnston WE, Vinten-Johansen J. Myocardial protective effects of  
13 514 adenosine. Infarct size reduction with pretreatment and continued receptor stimulation  
14 515 during ischemia. *Circulation.* 1992;86:986-94.
- 15 516 6. Lasley RD, Mentzer RM, Jr. Protective effects of adenosine in the reversibly injured heart.  
16 517 *Ann Thorac Surg.* 1995;60:843-6.
- 17 518 7. Ge ZD, van der Hoeven D, Maas JE, Wan TC, Auchampach JA. A(3) adenosine receptor  
18 519 activation during reperfusion reduces infarct size through actions on bone marrow-derived  
19 520 cells. *J Mol Cell Cardiol.* 2010;49:280-6.
- 20 521 8. Maddock HL, Mocanu MM, Yellon DM. Adenosine A(3) receptor activation protects the  
21 522 myocardium from reperfusion/reoxygenation injury. *Am J Physiol Heart Circ Physiol.*  
22 523 2002;283:H1307-13.
- 23 524 9. Liu GS, Thornton J, Van Winkle DM, Stanley AW, Olsson RA, Downey JM. Protection  
24 525 against infarction afforded by preconditioning is mediated by A1 adenosine receptors in  
25 526 rabbit heart. *Circulation.* 1991;84:350-6.
- 26 527 10. Ge ZD, Peart JN, Kreckler LM, et al. Cl-IB-MECA [2-chloro-N6-(3-  
27 528 iodobenzyl)adenosine-5'-N-methylcarboxamide] reduces ischemia/reperfusion injury in  
28 529 mice by activating the A3 adenosine receptor. *J Pharmacol Exp Ther.* 2006;319:1200-10.
- 29 530 11. McIntosh VJ, Lasley RD. Adenosine receptor-mediated cardioprotection: are all 4  
30 531 subtypes required or redundant? *J Cardiovasc Pharmacol Ther.* 2012;17:21-33.
- 31 532 12. Headrick JP, Lasley RD. Adenosine receptors and reperfusion injury of the heart. *Handb*  
32 533 *Exp Pharmacol.* 2009;193:189-214.
- 33 534 13. Jordan JE, Thourani VH, Auchampach JA, Robinson JA, Wang NP, Vinten-Johansen J.  
34 535 A(3) adenosine receptor activation attenuates neutrophil function and neutrophil-mediated  
35 536 reperfusion injury. *Am J Physiol.* 1999;277(5 Pt 2):H1895-905.
- 36 537 14. Maddock HL, Gardner NM, Khandoudi N, Bril A, Broadley KJ. Protection from  
37 538 myocardial stunning by ischaemia and hypoxia with the adenosine A3 receptor agonist,  
38 539 IB-MECA. *Eur J Pharmacol.* 2003;477:235-45.
- 39 540 15. Wan TC, Ge ZD, Tampo A, Mio Y, Bienengraeber MW, Tracey WR, et al. The A3  
40 541 adenosine receptor agonist CP-532,903 [N6-(2,5-dichlorobenzyl)-3'-aminoadenosine-5'-  
41 542 N-methylcarboxamide] protects against myocardial ischemia/reperfusion injury via the  
42 543 sarcolemmal ATP-sensitive potassium channel. *J Pharmacol Exp Ther.* 2008;324:234-43.
- 43 544 16. Auchampach JA, Ge ZD, Wan TC, Moore J, Gross GJ. A3 adenosine receptor agonist IB-  
44 545 MECA reduces myocardial ischemia-reperfusion injury in dogs. *Am J Physiol Heart Circ*  
45 546 *Physiol.* 2003;285:H607-13.
- 46 547 17. Dixon AK, Gubitza AK, Sirinathsinghji DJ, Richardson PJ, Freeman TC. Tissue  
47 548 distribution of adenosine receptor mRNAs in the rat. *Br J Pharmacol* 1996;118:1461-8.
- 48 549
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60
- 61
- 62
- 63
- 64
- 65

- 550 18. Morrison RR, Teng B, Oldenburg PJ, Katwa LC, Schnermann JB, Mustafa SJ. Effects of  
1 551 targeted deletion of A1 adenosine receptors on postischemic cardiac function and  
2 552 expression of adenosine receptor subtypes. *Am J Physiol Heart Circ Physiol*.  
3 553 2006;291:H1875-82.
- 4 554 19. Matot I, Weiniger CF, Zeira E, Galun E, Joshi BV, Jacobson KA. A3 adenosine receptors  
5 555 and mitogen-activated protein kinases in lung injury following in vivo reperfusion. *Crit*  
6 556 *Care*. 2006;10:R65.
- 7 557 20. Rivo J, Zeira E, Galun E, Matot I. Activation of A3 adenosine receptor provides lung  
8 558 protection against ischemia-reperfusion injury associated with reduction in apoptosis. *Am*  
9 559 *J Transplant*. 2004;4:1941-8.
- 10 560 21. Rivo J, Zeira E, Galun E, Matot I. Activation of A3 adenosine receptors attenuates lung  
11 561 injury after in vivo reperfusion. *Anesthesiology* 2004;101:1153-9.
- 12 562 22. Von Lubitz DK, Simpson KL, Lin RC. Right thing at a wrong time? Adenosine A3  
13 563 receptors and cerebroprotection in stroke. *Ann N Y Acad Sci*. 2001;939:85-96.
- 14 564 23. Von Lubitz DK, Lin RC, Boyd M, Bischofberger N, Jacobson KA. Chronic  
15 565 administration of adenosine A3 receptor agonist and cerebral ischemia: neuronal and glial  
16 566 effects. *Eur J Pharmacol*. 1999;367:157-63.
- 17 567 24. Gao Z, Li BS, Day YJ, Linden J. A3 adenosine receptor activation triggers  
18 568 phosphorylation of protein kinase B and protects rat basophilic leukemia 2H3 mast cells  
19 569 from apoptosis. *Mol Pharmacol*. 2001;59:76-82.
- 20 570 25. Kyriakis JM, Avruch J. Mammalian mitogen-activated protein kinase signal transduction  
21 571 pathways activated by stress and inflammation. *Physiol Rev*. 2001;81:807-69.
- 22 572 26. Hausenloy DJ. Signalling pathways in ischaemic postconditioning. *Thromb Haemost*.  
23 573 2009;101:626-34.
- 24 574 27. Hausenloy DJ, Mocanu MM, Yellon DM. Cross-talk between the survival kinases during  
25 575 early reperfusion: its contribution to ischemic preconditioning. *Cardiovasc Res*.  
26 576 2004;63:305-12.
- 27 577 28. Hausenloy DJ, Yellon DM. Reperfusion injury salvage kinase signalling: taking a RISK  
28 578 for cardioprotection. *Heart Fail Rev* 2007;12:217-34.
- 29 579 29. Shiryayev A, Moens U. Mitogen-activated protein kinase p38 and MK2, MK3 and MK5:  
30 580 menage a trois or menage a quatre? *Cell Signal*. 2010;22:1185-92.
- 31 581 30. Hausenloy DJ, Yellon DM. New directions for protecting the heart against ischaemia-  
32 582 reperfusion injury: targeting the Reperfusion Injury Salvage Kinase (RISK)-pathway.  
33 583 *Cardiovasc Res*. 2004;61:448-60.
- 34 584 31. Germack R, Griffin M, Dickenson JM. Activation of protein kinase B by adenosine A1  
35 585 and A3 receptors in newborn rat cardiomyocytes. *J Mol Cell Cardiol*. 2004;37:989-99.
- 36 586 32. Germack R, Dickenson JM. Adenosine triggers preconditioning through MEK/ERK1/2  
37 587 signalling pathway during hypoxia/reoxygenation in neonatal rat cardiomyocytes. *J Mol*  
38 588 *Cell Cardiol*. 2005;39:429-42.
- 39 589 33. Fox R, Aubert M. Flow cytometric detection of activated caspase-3. *Methods Mol Biol*.  
40 590 2008;414:47-56.
- 41 591 34. Abbracchio MP, Cattabeni F. Brain adenosine receptors as targets for therapeutic  
42 592 intervention in neurodegenerative diseases. *Ann N Y Acad Sci*. 1999;890:79-92.
- 43 593 35. Abe J, Baines CP, Berk BC. Role of mitogen-activated protein kinases in ischemia and  
44 594 reperfusion injury : the good and the bad. *Circ Res*. 2000;86:607-9.
- 45 595 36. Yue TL, Wang C, Gu JL, et al. Inhibition of extracellular signal-regulated kinase enhances  
46 596 Ischemia/Reoxygenation-induced apoptosis in cultured cardiac myocytes and exaggerates  
47 597 reperfusion injury in isolated perfused heart. *Circ Res*. 2000;86:692-9.
- 48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 598 37. Schulte G, Fredholm BB. Signaling pathway from the human adenosine A(3) receptor  
1 599 expressed in Chinese hamster ovary cells to the extracellular signal-regulated kinase 1/2.  
2 600 *Mol Pharmacol.* 2002;62:1137-46.
- 3 601 38. Graham S, Combes P, Crumiere M, Klotz KN, Dickenson JM. Regulation of p42/p44  
4 602 mitogen-activated protein kinase by the human adenosine A3 receptor in transfected CHO  
5 603 cells. *Eur J Pharmacol.* 2001;420:19-26.
- 6 604 39. Tracey WR, Magee W, Masamune H, Oleynek JJ, Hill RJ. Selective activation of  
7 605 adenosine A3 receptors with N6-(3-chlorobenzyl)-5'-N-methylcarboxamidoadenosine  
8 606 (CB-MECA) provides cardioprotection via KATP channel activation. *Cardiovasc Res.*  
9 607 1998;40:138-45.
- 10 608 40. Bullard AJ, Govewalla P, Yellon DM. Erythropoietin protects the myocardium against  
11 609 reperfusion injury in vitro and in vivo. *Basic Res Cardiol.* 2005;100:397-403.
- 12 610 41. Lim SY, Davidson SM, Paramanathan AJ, Smith CC, Yellon DM, Hausenloy DJ. The  
13 611 novel adipocytokine visfatin exerts direct cardioprotective effects. *J Cell Mol Med.*  
14 612 2008;12:1395-403.
- 15 613 42. Maddock HL, Mocanu MM, Yellon DM. Adenosine A(3) receptor activation protects the  
16 614 myocardium from reperfusion/reoxygenation injury. *Am J Physiol Heart Circ*  
17 615 *Physiol*2002;283:H1307-13.
- 18 616 43. Park SS, Zhao H, Jang Y, Mueller RA, Xu Z. N6-(3-iodobenzyl)-adenosine-5'-N-  
19 617 methylcarboxamide confers cardioprotection at reperfusion by inhibiting mitochondrial  
20 618 permeability transition pore opening via glycogen synthase kinase 3 beta.. *J Pharmacol*  
21 619 *Exp Ther.* 2006;318:124-31.
- 22 620 44. Jin YC, Kim KJ, Kim YM, et al. Anti-apoptotic effect of magnolol in myocardial  
23 621 ischemia and reperfusion injury requires extracellular signal-regulated kinase1/2 pathways  
24 622 in rat in vivo. *Exp Biol Med* 2008;233:1280-8.
- 25 623 45. Hausenloy DJ, Tsang A, Mocanu MM, Yellon DM. Ischemic preconditioning protects by  
26 624 activating prosurvival kinases at reperfusion. *Am J Physiol Heart Circ Physiol.*  
27 625 2005;288:H971-6.
- 28 626 46. Danial NN. BAD: undertaker by night, candyman by day. *Oncogene.* 2008;27 Suppl  
29 627 1:S53-70.
- 30 628 47. Scabini M, Stellari F, Cappella P, Rizzitano S, Texido G, Pesenti E. In vivo imaging of  
31 629 early stage apoptosis by measuring real-time caspase-3/7 activation. *Apoptosis.*  
32 630 2011;16:198-207.
- 33 631 48. Schulman D, Latchman DS, Yellon DM. Urocortin protects the heart from reperfusion  
34 632 injury via upregulation of p42/p44 MAPK signaling pathway. *Am J Physiol Heart Circ*  
35 633 *Physiol.* 2002;283:H1481-8.
- 36 634 49. Brar BK, Jonassen AK, Stephanou A, Santilli G, Railson J, Knight RA, et al. Urocortin  
37 635 protects against ischemic and reperfusion injury via a MAPK-dependent pathway. *J Biol*  
38 636 *Chem.* 2000;275:8508-14.
- 39 637 50. Jonassen AK, Sack MN, Mjos OD, Yellon DM. Myocardial protection by insulin at  
40 638 reperfusion requires early administration and is mediated via Akt and p70s6 kinase cell-  
41 639 survival signaling. *Circ Res.* 2001;89:1191-8.
- 42 640 51. Guo WP, Fu XG, Jiang SM, Wu JZ. Neuregulin-1 regulates the expression of Akt, Bcl-2,  
43 641 and Bad signaling after focal cerebral ischemia in rats. *Biochem Cell Biol.* 2010;88:649-  
44 642 54.
- 45 643 52. Mehrhof FB, Muller FU, Bergmann MW, et al. In cardiomyocyte hypoxia, insulin-like  
46 644 growth factor-I-induced antiapoptotic signaling requires phosphatidylinositol-3-OH-  
47 645 kinase-dependent and mitogen-activated protein kinase-dependent activation of the  
48 646 transcription factor cAMP response element-binding protein. *Circulation* 2001;104:2088-  
49 647 94.
- 50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 648 53. Thourani VH, Nakamura M, Ronson RS, Jordan JE, Zhao ZQ, Levy JH, et al. Adenosine  
1 649 A(3)-receptor stimulation attenuates postischemic dysfunction through K(ATP) channels.  
2 650 Am J Physiol. 1999;277(1 Pt 2):H228-35.  
3 651 54. Wan TC, Ge ZD, Tampo A, Mio Y, Bienengraeber MW, Tracey WR, et al. The A3  
4 652 adenosine receptor agonist CP-532,903 [N6-(2,5-dichlorobenzyl)-3'-aminoadenosine-5'-  
5 653 N-methylcarboxamide] protects against myocardial ischemia/reperfusion injury via the  
6 654 sarcolemmal ATP-sensitive potassium channel. J Pharmacol Exp Ther 2008;324:234-43.  
7 655 55. Ge ZD, van der Hoeven D, Maas JE, Wan TC, Auchampach JA. A(3) adenosine receptor  
8 656 activation during reperfusion reduces infarct size through actions on bone marrow-derived  
9 657 cells. J Mol Cell Cardiol. 2010;49:280-6.  
10 658 56. Yang XM, Krieg T, Cui L, Downey JM, Cohen MV. NECA and bradykinin at reperfusion  
11 659 reduce infarction in rabbit hearts by signaling through PI3K, ERK, and NO. J Mol Cell  
12 660 Cardiol. 2004;36:411-21.  
13 661  
14 662  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65





Figure 1a



**Figure 1b**






Figure 2







Figure 3




**Figure 4**



**Figure 5**



**Figure 6**



**Figure 7**




Figure 8




Figure 9




Figure 10




Figure 11

## List of Figures

**Figure 1a Shows the treatment protocol used for infarct size assessment studies**

**Figure 1b. Shows the treatment protocol used for tissue collection for western blot analysis**

**Figure 2. The A<sub>3</sub>AR agonist 2-Cl-IB-MECA (1nM) administered at reperfusion significantly limits infarct size in a MEK 1/2 –ERK 1/2-dependent manner. \*P<0.01 vs. Control. \*\*P<0.05 vs. 2-Cl-IB-MECA. Results are shown as Mean ± SEM.**

**Figure 3. The A<sub>3</sub>AR agonist 2-Cl-IB-MECA (1 nM) administered at reperfusion significantly limits infarct in the presence of PI3K inhibitor Wortmannin (5nM). #P<0.05 vs. Control, \$P<0.05 vs. 2-CL-IB-MECA. Results are shown as Mean ± SEM.**

**Figure 4. Assessment of the MEK 1/2 – ERK 1/2 cell survival pathway in 2-CL-IB-MECA mediated cardioprotection in isolated adult rat cardiomyocytes subjected to 6 hours hypoxia and 18 hours of reoxygenation. The MEK 1/2 inhibitor UO126 (10 µM) was administered at reoxygenation in the presence and absence of the A<sub>3</sub>AR agonist 2-CL-IB-MECA (1 nM). Results are shown as Mean ± SEM and are expressed as a percentage of the total cells counted. #P<0.001 vs. Normoxia. £P<0.001 vs. Hyp/Reox. \$ <0.001 vs. 2-CL-IB-MECA.**

**Figure 5. Assessment of the PI3K - AKT cell survival pathway in isolated adult rat cardiomyocytes subjected to 6 hours hypoxia and 18 hours of reoxygenation. The A<sub>3</sub>AR agonist 2-CL-IB-MECA (1 nM) was administered at reoxygenation in the presence and absence of the PI3K inhibitor Wortmannin (5 nM). Results are shown as Mean ± SEM and are expressed as a percentage of the total cells counted. # P<0.001 vs. Normoxia. £ P<0.001 vs. Hyp/Reox. \$ P<0.05 vs. 2-CL-IB-MECA.**

**Figure 6. Assessment of p-ERK 1/2 status in isolated hearts to subjected ischaemia followed by 5, 10 or 20 minutes of reperfusion in the presence and absence of the A<sub>3</sub> Agonist 2-CL-IB-MECA (1nM) (MECA). The MEK 1/2 inhibitor UO126 (10 µM) was administered at reperfusion in the presence and absence of the A<sub>3</sub>AR agonist 2-CL-IB-MECA (1 nM). Results are shown as Mean ± SEM three individual experiments. \*P<0.05 vs. Basal. \*\*P<0.01 vs. Basal. £ P<0.05 vs. Control 5 min Reperfusion. # P<0.05 vs. Control 10. ### P<0.001 vs. MECA 10 min Reperfusion.**

**Figure 7. Assessment of AKT<sub>(ser473)</sub> phosphorylation in isolated hearts to subjected ischaemia followed by 5, 10 or 20 minutes of reperfusion in the presence and absence of the A<sub>3</sub> Agonist 2-CL-IB-MECA (1 nM) (MECA). Basal hearts were perfused for 60 minutes with no treatment. Results are shown as Mean ± SEM of three individual experiments. \*\* P<0.05 vs. Basal. ## P<0.01 vs.**

**Control 10 10'R/F. \*\*\*P<0.001 vs. MECA 10'R/F. ###P<0.01 vs. MECA 10min Reperfusion.**

**Figure 8. Assessment of p-BAD<sub>(Ser 112)</sub> and p-BAD<sub>(ser136)</sub> activity in isolated adult rat cardiac myocytes subjected to 6 hours of hypoxia followed by 18 hours of reoxygenation. The A<sub>3</sub> agonist 2-CL-IB-MECA (1 nM) was administered at the onset of reoxygenation in the presence and absence of the MEK1/2 inhibitor UO126 (10 μM) or the PIK inhibitor Wortmannin (5nM). \*P<0.05 vs. Hyp/Reox. # P<0.05 vs. M2-CL-IB-MECA. Mean ± SEM of 4 individual experiments.**

**Figure 9. Cleaved-caspase 3 activity in isolated adult rat cardiac myocytes subjected to 6 hours of hypoxia followed by 18 hours of reoxygenation. The A<sub>3</sub> agonist 2-CL-IB-MECA (1 nM) was administered at the onset of reoxygenation in the presence and absence of the MEK1/2 inhibitor UO126(10 μM). #P<0.001 vs. Normoxia. \$P<0.001 vs. Hyp/Reox. Mean ± SEM of 5 individual experiments.**

**Figure 10. Cleaved-caspase 3 activity in isolated adult rat cardiac myocytes subjected to 6 hours of hypoxia followed by 18 hours of reoxygenation. The A<sub>3</sub> agonist 2-CL-IB-MECA (1 nM) was administered throughout reoxygenation in the presence and absence of the PI3 kinase inhibitor Wortmannin (5 nM). # P<0.001 vs. Normoxia. £ P<0.001 vs. Hyp/Reox. \$ P<0.05 vs. 2-CL-IB-MECA. Mean ± SEM of 5 individual experiments.**

**Figure 11. Shows a hypothetical schematic representation of the cell survival signalling cascades that may be involved in 2-CL-IB-MECA mediated cardioprotection when administered at reperfusion.**