

Structure-based modeling and dynamics of MurM, a *Streptococcus pneumoniae* penicillin resistance determinant present at the cytoplasmic membrane

York, A., Lloyd, A. J., del Genio, C. I., Shearer, J., Hinxman, K. J., Fritz, K., Fulop, V., Dowson, C. G., Khalid, S. & Roper, D. I.

Author post-print (accepted) deposited by Coventry University's Repository

Original citation & hyperlink:

York, A, Lloyd, AJ, del Genio, CI, Shearer, J, Hinxman, KJ, Fritz, K, Fulop, V, Dowson, CG, Khalid, S & Roper, DI 2021, 'Structure-based modeling and dynamics of MurM, a *Streptococcus pneumoniae* penicillin resistance determinant present at the cytoplasmic membrane', *Structure*, vol. 29, no. 7, 1, pp. 731-742.e6.

<https://dx.doi.org/10.1016/j.str.2021.03.001>

DOI 10.1016/j.str.2021.03.001

ISSN 0969-2126

ESSN 1878-4186

Publisher: Elsevier

NOTICE: this is the author's version of a work that was accepted for publication in *Structure*. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in *Structure*, 7:1, (2021) DOI: 10.1016/j.str.2021.03.001

© 2021, Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International <http://creativecommons.org/licenses/by-nc-nd/4.0/>

Copyright © and Moral Rights are retained by the author(s) and/ or other copyright owners. A copy can be downloaded for personal non-commercial research or study, without prior permission or charge. This item cannot be reproduced or quoted extensively from without first obtaining permission in writing from the copyright holder(s). The content must not be changed in any way or sold commercially in any format or medium without the formal permission of the copyright holders.

This document is the author's post-print version, incorporating any revisions agreed during the peer-review process. Some differences between the published version and this version may remain and you are advised to consult the published version if you wish to cite from it.

Structure-based modelling and dynamics of MurM, a *Streptococcus pneumoniae* penicillin resistance determinant that functions at the cytoplasmic membrane interface

Anna York^{a*}, Adrian. J. Lloyd^{a*}, Charo I. del Genio^b, Jonathan Shearer^c, Karen. J. Hinxman^a, Konstantin Fritz^a, Vilmos Fulop^a, Christopher. G. Dowson^a, Syma Khalid^{c**}
David. I. Roper^{a**}

^aSchool of Life Science, University of Warwick, Coventry, CV4 7AL

^bCentre for Fluid and Complex Systems, School of Computing, Electronics and Mathematics, University of

^cSchool of Chemistry, University of Southampton, Southampton, SO17 1BJ

*Joint first author

**Corresponding author

Email addresses: anna.york@yale.edu (Anna York), Adrian.lloyd@warwick.ac.uk (Adrian. J. Lloyd), ad0364@coventry.ac.uk (Charo I. del Genio), Jrws1m15@soton.ac.uk (Jonathan Shearer), k.hinxman@imperial.ac.uk (Karen. J. Hinxman), k.fritz@warwick.ac.uk (Konstantin Fritz), Vilmos.fulop@warwick.ac.uk (Vilmos Fulop), S.Khalid@soton.ac.uk (Syma Khalid), c.g.dowson@warwick.ac.uk (Christopher. G. Dowson), David.roper@warwick.ac.uk (David. I. Roper)

Summary

MurM, an aminoacyl-tRNA dependent ligase, generates branched Lipid II required for indirect cross-linked peptidoglycan, and is essential for high-level penicillin resistance in the human pathogen *Streptococcus pneumoniae*. We have solved the X-ray crystal structure of *Staphylococcus aureus* FemX, an isofunctional homologue of MurM, and used this as a template to generate a MurM homology model. Using this model, we perform molecular docking and molecular dynamics to examine the interaction of MurM with the phospholipid bilayer and the membrane embedded Lipid II substrate. Our model suggests that MurM is associated with the major membrane phospholipid cardiolipin, and experimental evidence confirms that the activity of MurM is enhanced by this phospholipid and inhibited by its direct precursor phosphatidylglycerol. The spatial association of pneumococcal membrane phospholipids and their impact on MurM activity may therefore be critical to the final architecture of peptidoglycan and the expression of clinically relevant penicillin resistance in this pathogen.

Key Words

Streptococcus pneumoniae, MurM, penicillin resistance, peptidoglycan, indirect cross-links, lipid bilayer, homology modelling, molecular dynamics, molecular docking

Introduction

The peptidoglycan (PG) of the bacterial cell wall is a polymer consisting of alternating β -1,4 linked *N*-acetyl glucosamine (GlcNAc) and *N*-acetyl muramic acid (MurNAc) residues. Appended to the MurNAc sugar is a pentapeptide stem that can be

cross-linked directly or indirectly to form a rigid mesh-like structure (Bugg et al., 2011). PG biosynthesis begins with the cytoplasmic formation of a Peptidoglycan nucleotide, which is subsequently converted into a lipid-linked PG precursor known as Lipid II. Lipid II is then transported across the membrane, where it is polymerised and cross-linked by the penicillin-binding proteins (PBPs) (Figure 1). PG is an essential component of the cell wall, involved in cell growth and division, maintaining structural integrity, and resisting high osmotic pressures. Inhibition of cell wall biosynthesis is a key mechanism for many antibiotics, including β -lactams, glycopeptides and amino acid analogues (Schneider and Sahl, 2010).

In *S. pneumoniae* and other Gram-positive bacteria, the glutamate at the second position of the Lipid II pentapeptide is α -amidated to *iso*-glutamine by the essential GatT/MurD complex (Figueiredo et al., 2012, Münch et al., 2012, Zapun et al., 2013, Morlot et al., 2018). In addition, branched Lipid II, capable of generating indirect cross-links, can be formed by the non-essential MurM and MurN proteins. MurM and MurN are responsible for the sequential addition of amino acids to the third-position lysine of the pentapeptide stem (Filipe et al., 2000). MurM can append either L-serine or L-alanine at the first position of the dipeptide bridge, whilst MurN extends this modification by addition of an invariable L-alanyl moiety. Branched PG precursors are also found in several other Gram-positive bacterial pathogens, for example the glycyl-tRNA^{Gly} dependent enzymes FemX, A and B are responsible for the addition of a pentaglycyl bridge in *Staphylococcus aureus* (Schneider et al., 2004). In comparison with other Gram positive organisms, the PG of *S. pneumoniae* is highly heterogeneous: the predominant C-terminal amino acid at position 1 of the dipeptide, and the proportion of indirect cross-links throughout the PG, vary significantly between different strains (Severin and Tomasz, 1996, Garcia-Bustos et al., 1987, Garcia-Bustos and Tomasz, 1990). *In vitro* and *in vivo* studies indicate that MurM from the penicillin-resistant strain *S. pneumoniae*(159) preferentially incorporates L-alanine, whilst MurM from a penicillin-sensitive strain *S. pneumoniae*(Pn16) preferentially incorporates L-serine (Lloyd et al., 2008). In addition, penicillin resistant strains demonstrated higher levels of indirect cross-linking in the PG compared to penicillin susceptible isolates, however the overall degree of cross-linking remained constant (Garcia-Bustos and Tomasz, 1990).

Resistance to β -lactam antibiotics in *S. pneumoniae* is characterised by extensive interspecies recombination of PBP transpeptidase domains, which results in a mosaic PBP active site with a lower β -lactam binding affinity (Smith et al., 1991). This mechanism of resistance contrasts with that of many other bacteria that have acquired genes for β -lactamase enzymes which inactivate the antibiotic before it binds to and inhibits the PBPs. Interestingly, deletion of the *murM* gene in *S. pneumoniae* eliminates indirect cross-links from the PG and results in a complete loss of penicillin resistance (Filipe et al., 2001). It has been proposed that the changes to the PBP active site which prevent β -lactam binding, may also alter the Lipid II substrate specificity such that the PBPs bind branched Lipid II more tightly than unbranched Lipid II. MurM is therefore necessary, but not sufficient for resistance in clinical strains of *S. pneumoniae*, making

it an interesting target for the development of new inhibitors of antimicrobial resistance (Filipe and Tomasz, 2000).

The cytoplasmic membrane of *S. pneumoniae* contains two phospholipids, phosphatidylglycerol (PhG) and cardiolipin (CL) (Trombe et al., 1979, Pesakhov et al., 2007), where cardiolipin synthase is responsible for generating cardiolipin from two molecules of phosphatidylglycerol (Schlame, 2008). The proportion of cardiolipin and phosphatidylglycerol, as a percentage of the overall membrane lipids, varies in *S. pneumoniae* between anaerobic and aerobic growth conditions. Cardiolipin was found to decrease from 15.3 % to 8.3 %, whilst phosphatidylglycerol increased from 12.7 % to 16.3 % in anaerobic conditions compared to aerobic conditions (Pesakhov et al., 2007). The peptidoglycan precursor, Lipid II is tethered to the cell membrane by virtue of its C55 Lipid II tail. MurM then transfers a single amino acyl moiety (Seryl or Alanyl) from aminoacyl-tRNA to the third position lysine of the cytoplasmically oriented Lipid II pentapeptide stem.

Previously, MurM inhibitors have been identified; however, none have shown growth inhibition of effect on penicillin MIC, indicating that these compounds cannot effectively cross the cytoplasmic membrane of *S. pneumoniae* (Cressina et al., 2007, 2009). MurM has thus far resisted extensive crystallization in our laboratory, and consequently, its X-ray-solved structure is not available. However, in a related study we were able to solve the X-ray structure of the isofunctional homologue of MurM from *S. aureus* (FemX), which we have used here as a template for homology modelling of MurM. Using this MurM homology model we have successfully identified the Lipid II binding site, and used molecular dynamics (MD) simulations to investigate interactions between MurM and both membrane phospholipids and its Lipid II substrate (Witzke et al., 2016). We subsequently, studied the effects of these membrane embedded phospholipids, on the enzymatic activity of MurM *in vitro*, corroborating our *in silico* analysis. These studies provide new insights into the structure and activity of MurM, providing a link between phospholipid membrane composition and peptidoglycan architecture. This may be useful for the development of novel chemical probes for these proteins, and have important implications for future studies on penicillin resistance mechanisms in *S. pneumoniae*.

Results

X-ray crystallography and structure determination of *S. aureus* FemX

The crystal structure of *S. aureus* FemX was solved to a resolution of 1.62 Å and the structure was deposited in the PDB with accession number: 6SNR. A summary of the data collection and refinement statistics is given in Table 1.

The final solved structure of *S. aureus* FemX, contains two domains; a globular domain and a coiled-coil domain. Similarly to FemA (Benson et al., 2002), the globular domain can be divided into two subdomains. Each subdomain contains a central five-stranded mixed polarity β -sheet surrounded by four α -helices. Subdomain 1A

comprises residues 1-145 and 384-421 whilst subdomain 1B comprises residues 146-234 and 298-383. Unfortunately, residues 403-421 were not present in the density. The coiled-coil domain consists of two antiparallel α -helices, comprising residues 235-297. FemX and FemA can be superimposed onto each other with a root-mean-square deviation (RMSD) of ~ 2.7 Å over 384 residues. Similarly to FemA, FemX has a deep L-shaped channel of about 20 x 40 Å located along side the globular domain and mainly in subdomain 1B. This channel comprises a peptidoglycan precursor binding site which was previously identified in *S. aureus* FemA (Benson et al., 2002). The identity of this peptidoglycan precursor binding site has been further confirmed by the crystallography studies of *Weissella viridescens* FemX complexed with substrates (Biarrotte-Sorin et al., 2004).

Homology modelling of *S. pneumoniae* MurM

The structures of two MurM homologues, *S. aureus* FemA and *W. viridescens* FemX were solved previously by X-ray crystallography (Benson et al., 2002, Fonvielle et al., 2013, Biarrotte-Sorin et al., 2004). The *S. aureus* FemA structure was subsequently used as a template for homology modelling of MurM by Fiser et al. (2003). However, alignment of *S. aureus* FemA, *S. aureus* FemX, *S. pneumoniae* MurM and *W. viridescens* FemX (Supplemental Information: Figure S2) showed that *S. aureus* FemX possesses the highest sequence identity to MurM. *S. aureus* FemX is also more functionally homologous to *S. pneumoniae* MurM, and appends the first amino acid of the cross-bridge to the Lipid II precursor (Matsushashi et al., 1967, Schneider et al., 2004). In contrast *S. aureus* FemA appends the second and third amino acid residues of the cross-bridge to the α -amino group of a glycyl residue appended to the ϵ -amino group of the stem peptide L-lysyl residue of the Lipid II precursor. Therefore, given the difficulties in obtaining MurM crystals, we were motivated to solve the structure of its functional homologue (*S. aureus* FemX) by X-ray crystallography, to access by *in silico* approaches the structure of MurM.

Using the newly solved structure of *S. aureus* FemX we generated a new homology model for MurM which consists of a globular domain comprising two subdomains and a coiled-coil helical arm (Figure 2). Each subdomain comprises two twisted β -sheet cores surrounded by α -helices; subdomain 1A is formed of residues 1-153 and 382-401, whilst subdomain 1B is made up of residues 154-241 and 294-381. The coiled-coil domain comprises residues 242-293. Whilst the new MurM homology model is similar to the previous model (Fiser et al., 2003), the RMSD of the two models is 3.8 Å over 368 residues, indicating that there are some key structural differences, namely; loss of N-terminal $\beta 1$, and antiparallel $\beta 6/\beta 13$ from the previous model; addition of $\alpha 5$ and $\beta 11/\beta 12$; and presence of α -helical secondary structure at the C-terminal end of the new MurM model.

Identification of a possible Lipid II binding site of MurM

The new MurM model revealed a binding pocket which was not present in the previous model of MurM. Structural comparison between *W. viridescens* FemX co-crystallised with its UDP-MurNAc-pentapeptide substrate and the new MurM model, allowed identification of a Lipid II binding site that corresponds with those identified previously in

S. aureus FemA and FemX, as well as *W. viridescens* FemX. When the new MurM model and the *W. viridescens* FemX were aligned and overlaid, the newly identified MurM binding site appeared to easily accommodate the soluble UDP-MurNAc-pentapeptide substrate well (Figure 3A). The following 8 residues; Tyr103, Lys36, Asn38, Trp39, Thr209, Arg211, Try215 and Tyr256, were independently proposed to be involved in substrate binding in both *W. viridescens* FemX structures (Biarrotte-Sorin et al., 2004, Fonvielle et al., 2013). The corresponding MurM residues, defined as having residues which have similar properties, and occupying a similar location and orientation in physical space, with side chains facing the binding pocket, were identified in the MurM structure as Phe103, Lys35, Trp38, Arg215 and Tyr219, therefore these residues may also be important for substrate binding in MurM.

Next molecular docking using AutoDock Vina (Trott and Olson, 2010) was conducted to independently investigate docking of the Lipid II substrate to the new MurM model. Lipid II is a large molecule that is, in general, unsuitable for molecular docking studies. In addition, the lipid tail is embedded in the membrane, and so is not itself available for binding to MurM. Therefore, a truncated Lipid II molecule, comprised of a methyl capped diphosphate GlcNAc-MurNAc-pentapeptide, was used for these docking experiments (Supplemental Information: Figure S4).

When AutoDock Vina Trott and Olson, 2010) was allowed to search the entire protein surface of MurM, all docking results returned were within the identified binding site, indicating that there are no other suitable binding sites on the protein. The search was then restricted to the binding site and the top ten results were obtained. The top five results obtained all had identical binding affinities of $-7.3 \text{ kcal.mol}^{-1}$. Two docking orientations, whereby the phosphates are located deep within the binding pocket, would be physically impossible for the natural substrate (Lipid II) *in vivo*, since the membrane-embedded prenyl lipid tail is appended via the phosphate. The remaining 3 docking orientations, all orient the phosphates close to the opening of the binding site with the pentapeptide chain disappearing deep into the binding pocket. The exact orientation of the pentapeptide chain is variable, indicating that the binding site is spacious and that Lipid II may be accommodated in a number of different possible orientations. Figure 3B shows one conformation in which the docking of truncated Lipid II is similar to the orientation of the soluble UDP-MurNAc-pentapeptide from *W. viridescens* FemX overlaid with MurM (Figure 3A), the remaining four substrate orientations with binding affinities of $-7.3 \text{ kcal.mol}^{-1}$ are shown in the Supplementary Information (Figure S5). A key limitation of docking is that it considers the protein as rigid; therefore, multiple substrate orientations may indicate that conformational changes within the binding site may occur upon substrate binding or during catalysis.

Additional docking studies were performed to further explore the lipid II binding pose in the putative MurM binding site and establish whether this binding site shows specific affinity for lipid II or is indiscriminate between different lipid species. Lipid II, cardiolipin, phosphatidylglycerol and phosphatidylethanolamine lipids (full molecules) were docked into the putative binding site, where the binding site is defined by the residues Lys35, Trp38, Phe103, Arg215 and Tyr219. We found lipid II bound into this site with atoms of the headgroup within 3 Å of the residues Lys35, Trp38, Phe215 and Arg219, and with the lipid II tail outside of this site. In stark contrast the cardiolipin and phosphatidylethanolamine headgroups did not show any preference for this site,

instead the binding poses showed their tails inserted into the cavity. There was one binding pose for phosphatidylglycerol in which the lipid headgroup was in the putative binding site, however a portion of the tail was also inside it, other two poses of phosphatidylglycerol had the tails inserted into this cavity. For each lipid type the top cluster of binding poses produced by HADDOCK (Van Zundert et al., 2016) were analysed, and the top 3 binding poses for each lipid are shown in the Supplemental Information (Figure S6).

Interactions between MurM and the lipid bilayer

Given the docking studies are performed in the absence of the membrane environment, a set of coarse-grained MD simulations was next performed to establish the likelihood of MurM being orientated on the membrane such that the putative binding site is available to Lipid II. Molecular dynamics simulations were used to model the interactions of MurM with the lipid bilayer. Six independent coarse-grained simulations were conducted for each of three membrane systems described in the Supplemental Information (Table S1). In 16 of these simulation runs MurM readily associated with the membrane in $<3 \mu s$ and its orientation with respect to the membrane remained unchanged for the remaining $2 \mu s$ (Supplemental Information: Figure S7). In addition, Lipid II in the same leaflet as the peripheral MurM was found to cluster around the protein; more than 50 % of Lipid II molecules were located within 2 nm of the MurM. MurM associated with the membrane in a number of different orientations which can be classified into two groups, those in which the binding site predicted from docking studies is available for lipid binding and those in which it is not. Table 2 shows a breakdown of results from the 18 unbiased simulations; MurM adhered to the membrane in 16 of the simulations and of these, the binding site was available for Lipid II binding in 11 and unavailable in only 5. In 3 out of 18 unbiased simulations, the MurM was oriented such that the Lipid II molecule was located in the putative binding site, which demonstrates that the Lipid II is able to successfully enter this binding site even on the short timescale of a MD simulation (Figure 4). Back mapping of one of these systems to all-atom resolution allowed the binding site to be explored in more detail. Three independent atomistic simulations each of 250 ns duration found Lipid II located in the same binding site of MurM that was identified during molecular docking of the truncated Lipid II substrate, and in the co-crystal structure of UDP-MurNAc-pentapeptide with *W. viridescens* FemX (Figure 3A). In both the molecular docking and atomistic simulations, the lipid II headgroup forms stable interactions with Lys35, Trp38, Arg215 and Tyr219.

Similarly to the molecular docking findings, this MD simulation shows that the MurM binding site is flexible and allows the Lipid II molecule to adopt a wide variety of conformations (Figure 5). This may suggest that binding of a second substrate or a large conformational change may be required for catalysis.

Interactions between MurM and membrane phospholipids

Molecular dynamics simulations were used to investigate the effects of membrane phospholipids (cardiolipin and phosphatidylglycerol) on MurM at the cytoplasmic membrane interface. To investigate local lipid enrichment/depletion, we calculated 2D enrichment maps across the entire membrane and D-E indices within a 1.1 nm of MurM, as described by Corradi et al. (2018) (See STAR Methods). A D-E Index < 0 , or an enrichment percentage < 0 %, indicated that the specified lipid was depleted with

respect to the bulk membrane composition. Figure 6 shows that upon association of MurM with the cytoplasmic membrane, there was no effect on the distribution of phosphatidylglycerol or phosphatidylethanolamine. However, in membranes containing 8 % or 16 % cardiolipin, cardiolipin was enriched at the MurM:membrane interface, whilst the phosphatidylethanolamine and phosphatidylglycerol distributions remained largely unaffected. The importance of these observations were considered *in vitro* by measuring the enzymatic activity of MurM in the presence of varying concentrations of cardiolipin or phosphatidylglycerol. These enzymatic studies show that cardiolipin activates MurM, whilst phosphatidylglycerol inhibits MurM in a concentration dependent manner. Figure 6E shows the enzymatic activation of MurM with respect to cardiolipin concentration, a 9.1-fold activation of MurM was achieved, with 50 % activation occurring at 0.4 mM cardiolipin. Figure 6F shows that the activity of MurM could be completely inhibited by phosphatidylglycerol, with an IC_{50} of 0.2 mM. Furthermore, Hill coefficients of 2.7 ± 0.3 and 2.8 ± 0.2 for cardiolipin and phosphatidylglycerol respectively, indicate that both these phospholipids exhibit their effects on MurM in a cooperative manner. Phosphatidylethanolamine, used in the construction of the model pneumococcal membrane to which MurM bound, when tested at a concentration of 0.72 mM, only slightly activated MurM activity by 0.32-fold (duplicate determination with a difference of <10 %). In comparison, 0.72 mM cardiolipin activated MurM by 8-fold (Figure 6E). Therefore, the impact of phosphatidylethanolamine on the disposition of MurM relative to its interaction with Lipid II and the phospholipid bilayer could be neglected.

Discussion

The crystal structure of *S. aureus* FemX has allowed the us to generate an improved homology model of MurM leading to the identification of a putative Lipid II binding site. Fiser et al. (2003) proposed a different MurM model and speculated about an alternative binding site based upon structural and functional analogy between MurM and N-myristoyltransferase (NMT) proteins. However, whilst the substrates of both NMT proteins and MurM are lipids, they are contextually very different. The NMT proteins are cytoplasmic proteins that contain a deep, narrow pocket which is highly specific for the myristoyl fatty acyl chain (Wright et al., 2010, Heuckeroth et al., 1988). In contrast, MurM binds the disaccharide head group and pentapeptide side chain of Lipid II, and the undecaprenyl C55 lipid tail is embedded in the membrane. Despite similarities with NMT proteins, our newly identified substrate binding site more closely resembles those of *W. viridescens* FemX, *S. aureus* FemX and FemA.

The orientations of truncated Lipid II in docking studies and Lipid II in molecular dynamics simulations are strikingly similar to each other and also the orientation of UDP-MurNAc-pentapeptide substrate co-crystallised in *W. viridescens* FemX. This is consistent with the observation that, although inefficient compared to Lipid II, UDP-MurNAc pentapeptide is a MurM substrate (Lloyd et al., 2008). In all cases, the

diphosphates are near the surface of the protein, and the protruding pentapeptide reaches into the binding pocket with the third-position lysine on the left-hand side of the binding pocket. In addition, previous studies suggest that the height of the Lipid II head group is 19 Å (Ganchev et al., 2006), and the binding pocket of this model was measured to be 15 Å. Since the Lipid II head group is flexible, and the binding site provides enough room for the substrate to bend, these measurements are consistent. Together with our findings, this strongly supports the identification of this newly identified cavity as the Lipid II binding site and suggests that the Lipid II binds to MurM in an orientation similar to that of *W. viridescens* FemX binding to its substrate.

Alanylphosphatidylglycerol synthase (PDB ID 4v34) similarly to MurM, also utilises both lipid and alanyl-tRNA^{Ala} substrates (Hebecker et al., 2015), in order to successfully bring these two substrates together for catalysis, it possesses two binding sites, located on opposite sides of the protein which are connected by a channel. The protein itself provides a barrier between the hydrophobic lipid and the hydrophilic tRNA, such that they do not come into close proximity with each other. The negatively charged surface patch identified previously (Fiser et al., 2003) remains present on this new homology model of MurM, and is located on the opposite side of the protein with respect to the Lipid II binding site, which is located within a positively charged surface patch (Supplementary Material: Figure S8). This negative patch is unsuitable for the binding of negatively charged tRNA and so it is unlikely that MurM shares the same mechanism of action as alanylphosphatidylglycerol synthase. The negatively charged surface patch may however be important for protein:protein interactions occurring either at the cell surface or in the cytoplasm.

These modelling studies reveal that the stem peptide protrudes perpendicular to the surface of the membrane into the active site of MurM. Therefore, in order for alanyl-tRNA^{Ala} to simultaneously interact with MurM, whilst it is located over its lipid substrate, the highly negatively charged hydrophilic tRNA would have to be brought into close proximity with the negatively charged phospholipid head groups and/or the hydrophobic phospholipid tails below them. Given that this would be a highly unfavourable interaction, we propose an alternative 'ping-pong' mechanism of action for MurM whereby MurM is initially aminoacylated by alanyl- or seryl-tRNA in the cytoplasm before travelling to the cell membrane for transfer to Lipid II. Whilst the MurM is in the cytoplasm and not interacting with the membrane, the positively charged patch, located at the newly proposed Lipid II binding site, may facilitate interaction with a polyanionic substrate such as tRNA. Once this has occurred, subsequent interaction of the aminoacyl-MurM with the surface of the membrane could accommodate the correct and catalytically productive interaction of aminoacylated-MurM with Lipid II. Although this proposed mechanism is at variance with the sequential mechanism of catalysis proposed for *W. viridescens* FemX (Hegde and Blanchard, 2003), in this case, both substrates were highly hydrophilic nucleotide or polynucleotide derivatives in the same cellular sub-compartment and are therefore without biophysical impediment with regard to their proximity during catalysis. Here, with regard to MurM, the chemical properties and location of both substrates indicate an advantage to a mechanism which avoids their simultaneous binding.

The pneumococcal peptidoglycan is heterogeneous with respect to its composition of directly and indirectly cross-linked stem peptides. It remains unclear as to whether the activity of MurM, and therefore the generation of indirect cross-links is distributed equally around the entire cell surface, or whether it is localised to specific sites. Phospholipids are known to be involved in the spatial and temporal biochemistry of cells (Lin et al., 2019), and cardiolipin was shown to be enriched at the poles and septa of *E. coli* and *Bacillus subtilis*, localising specific membrane-associated proteins to these regions (Bramkamp and Lopez, 2015). Our simulations indicate that, whilst cardiolipin enrichment occurs within the membrane in the presence of MurM, this phospholipid is not essential for membrane association of MurM to occur. Therefore, it remains uncertain as to whether *in vivo* cardiolipin is highly concentrated in patches in the membrane and is used to recruit MurM to that location, or whether association of MurM with the membrane drives the enrichment of cardiolipin in the membrane.

Despite this uncertainty, we show that cardiolipin stimulates the enzymatic activity of MurM, and whilst it is not clear if this increased activity is as a result of a direct effect on the protein or the Lipid II substrate, or both, the spatial association of cardiolipin to the MurM protein suggests that at least some of this effect may be due to direct interactions with the MurM protein. Cardiolipin has previously been found to bind to and activate a wide range of proteins including MurG (Boots et al., 2003), rat liver protein kinase N (Morrice et al., 1994, Peng et al., 1996), porcine heart AMP deaminase (Purzycka-Preis and Zydowo, 1987), rat liver multi-catalytic proteinase (Ruiz de Mena et al., 1993), *E. coli* glycerol-3-phosphate acyltransferase (Scheideler and Bell, 1989), *E. coli* dnaA (Sekimizu and Kornberg, 1988) and streptococcal hyaluronan synthases (Tlapak-Simmons et al., 1999a,b, 2004, Weigel et al., 2006, Tlapak-Simmons et al., 1998). This further supports the contention that cardiolipin affects MurM activity by directly interacting with MurM. Similar cardiolipin-mediated sigmoidal stimulatory effects have been seen with other streptococcal membrane proteins such as the hyaluronan synthases from *Streptococcus pyogenes* and *Streptococcus equismilis* (Tlapak-Simmons et al., 1999a,b, 2004, Weigel et al., 2006). In these examples, up to sixteen cardiolipin molecules are believed to associate with single hyaluronan synthase molecule (Tlapak-Simmons et al., 1998).

We also show that phosphatidylglycerol inhibits the catalytic activity of MurM, and that the concentration of this lipid in the membrane environment surrounding the MurM changes very little. Therefore, the inhibitory effect of phosphatidylglycerol may be exerted by altering the presentation of the Lipid II substrate to MurM, rather than by having a direct effect on the protein itself. It is possible that in *S. pneumoniae*, as in *E. coli* and *B. subtilis*, cardiolipin gathers in specific regions of the membrane, where it localises and up-regulates the activity of MurM, resulting in higher levels of indirect cross-linking in these regions.

Whilst MurM alone is not sufficient for penicillin resistance, the enzyme is crucial together with mosaic *S. pneumoniae* PBPs for the generation of a highly resistant phenotype. Deletion of *murM* from resistant strains resulted in a virtual abolition of

penicillin resistance that could not be restored by *PBP* DNA. Indeed, additional *murM* DNA from a resistant strain was required for full expression of donor level penicillin resistance (Filipe and Tomasz, 2000, Smith and Klugman, 2001). Given the importance of MurM for penicillin resistance, the enrichment of cardiolipin at the MurM:membrane interface which activates MurM, and the inhibition of MurM activity by phosphatidylglycerol, may regulate the penicillin resistance phenotype imparted by MurM activity which may therefore be regulated by cardiolipin synthase activity. These findings have therefore revealed a crucial and hitherto unexplored area with regards to establishing a more complete understanding of penicillin resistance mechanisms and the influence on them by other areas of pneumococcal metabolism.

This new MurM structural model allowed identification of the Lipid II binding site and the contextual presentation of this substrate to MurM, characterised the impact of membrane phospholipids on MurM at the MurM:membrane interface, and may have spatial mechanistic implications for the catalytic activity of this protein. Molecular dynamics enabled the *in silico* investigation into MurM:membrane interactions, which are often overlooked when studying enzymes which act at the cytoplasmic membrane interface. The subsequent *in vitro* experiments on the importance of phospholipids for MurM activity, corroborate the *in silico* findings, supporting the role of phospholipids as an important contributor to the regulation of MurM at the membrane. These studies provide new insights into the structure of MurM which may guide future mutational studies, and allow a more detailed analysis of the structure-function relationship of this protein. This research contributes important findings towards achieving a more complete understanding of its role in pneumococcal penicillin resistance mechanisms.

Acknowledgements

This work was supported in part by the Midlands Integrative Biosciences Training Partnership (MIBTP) BBSRC grant BB/J014532/1, and the Centre for Doctoral Training in Theory and Modelling in Chemical Sciences (TMCS DTC) EPSRC grant EP/L015722/1, as well as MRC grants G1100127, G0400848, MR/N002679/1 and BBSRC grant BB/N003241/1. The authors would like to acknowledge the help of the Media Preparation Facility in the School of Life Sciences at the University of Warwick. We would also like to thank Dr Allister Crow for his help and support with using PyMOL.

Author Contributions

Conceptualization, A.Y., A.J.L., D.I.R., and S.K.; Investigation, K.F., K.J.H., V.F., A.J.L., A.Y., J.S., C.I.G.; Methodology/Software/Formal Analysis, C.I.G., J.S., A.Y., and S.K.; Writing – Original Draft, A.Y., J.S., and C.I.G.; Writing – Review & Editing, A.Y., A.J.L., S.K., D.I.R.; and C.G.D.; Funding Acquisition, A.Y., J.S., and C.G.D.; Supervision, A.J.L., D.I.R., S.K., and C.G.D.; Project Administration, A.Y.; Visualization, A.Y., A.J.L., J.S., and S.K.

Declaration of Interests

The authors declare no competing interests.

References

- M. J. Abraham, T. Murtola, R. Schulz, S. Páll, J. C. Smith, B. Hess, and E. Lindahl. GROMACS: High performance molecular simulations through multi-level parallelism from laptops to supercomputers. *SoftwareX*, 1:19–25, 2015.
- T. E. Benson, D. B. Prince, V. T. Mutchler, K. A. Curry, A. M. Ho, R. W. Sarver, J. C. Hagadorn, G. H. Choi, and R. L. Garlick. X-ray crystal structure of *Staphylococcus aureus* FemA. *Structure*, 10(8): 1107–1115, 2002.
- H. J. Berendsen, J. v. Postma, W. F. van Gunsteren, A. DiNola, and J. Haak. Molecular dynamics with coupling to an external bath. *The Journal of chemical physics*, 81(8):3684–3690, 1984.
- S. Biarrotte-Sorin, A. P. Maillard, J. Delettré, W. Sougakoff, M. Arthur, and C. Mayer. Crystal structures of *Weissella viridescens* FemX and its complex with UDP-MurNAc-pentapeptide: insights into FemABX family substrates recognition. *Structure*, 12(2):257–267, 2004.
- J.-W. P. Boots, R. E. Spelbrink, G. M. Kool, E. Breukink, J. A. Killian, B. de Kruijff, et al. Membrane interaction of the glycosyltransferase MurG: a special role for cardiolipin. *Journal of bacteriology*, 185 (13):3773–3779, 2003.
- M. Bramkamp and D. Lopez. Exploring the existence of lipid rafts in bacteria. *Microbiol. Mol. Biol. Rev.*, 79(1):81–100, 2015.
- A. T. Brünger. Free R value: a novel statistical quantity for assessing the accuracy of crystal structures. *Nature*, 355(6359):472–475, 1992.
- T. Bugg, D. Braddick, C. Dowson, and D. Roper. Bacterial cell wall assembly: still an attractive antibacterial target. *Trends Biotechnology*, 29(4):167–73, 2011.
- V. Corradi, E. Mendez-Villuendas, H. I. Ingólfsson, R.-X. Gu, I. Siuda, M. N. Melo, A. Moussatova, L. J. DeGagné, B. I. Sejdiu, G. Singh, et al. Lipid–protein interactions are unique fingerprints for membrane proteins. *ACS central science*, 4(6):709–717, 2018.
- E. Cressina, A. J. Lloyd, G. De Pascale, D. I. Roper, C. G. Dowson, and T. D. Bugg. Adenosine phosphonate inhibitors of lipid II: alanyl tRNA ligase MurM from *Streptococcus pneumoniae*. *Bioorganic & medicinal chemistry letters*, 17(16):4654–4656, 2007.
- E. Cressina, A. J. Lloyd, G. De Pascale, B. J. Mok, S. Caddick, D. I. Roper, C. G. Dowson, and T. D. Bugg. Inhibition of tRNA-dependent ligase MurM from *Streptococcus pneumoniae* by phosphonate and sulfonamide inhibitors. *Bioorganic & medicinal chemistry*, 17(9):3443–3455, 2009.
- D. Cruickshank. Remarks about protein structure precision. *Acta Crystallographica Section D: Biological Crystallography*, 55(3):583–601, 1999.
- T. Darden, D. York, and L. Pedersen. Particle mesh Ewald: An $N \log(N)$ method for Ewald sums in large systems. *The Journal of chemical physics*, 98(12):10089–10092, 1993.
- D. H. de Jong, G. Singh, W. D. Bennett, C. Arnarez, T. A. Wassenaar, L. V. Schäfer, X. Periole, D. P. Tieleman, and S. J. Marrink. Improved parameters for the martini coarse-grained protein force field. *Journal of Chemical Theory and Computation*, 9(1):687–697, 2012.

- D. H. de Jong, G. Singh, W. F. D. Bennett, C. Arnarez, T. A. Wassenaar, L. V. SchÄfer, X. Periole, D. P. Tieleman, and S. J. Marrink. Improved Parameters for the Martini Coarse-Grained Protein Force Field. *Journal of Chemical Theory and Computation*, 9(1):687–697, 2013. doi: 10.1021/ct300646g. URL <https://doi.org/10.1021/ct300646g>. PMID: 26589065.
- G. Q. Dong, H. Fan, D. Schneidman-Duhovny, B. Webb, and A. Sali. Optimized atomic statistical potentials: assessment of protein interfaces and loops. *Bioinformatics*, 29(24):3158–3166, 2013.
- S. Doubl  . [29] Preparation of selenomethionyl proteins for phase determination. In *Methods in enzymology*, volume 276, pages 523–530. Elsevier, 1997.
- P. Emsley, B. Lohkamp, W. G. Scott, and K. Cowtan. Features and development of Coot. *Acta Crystallographica Section D: Biological Crystallography*, 66(4):486–501, 2010.
- N. Eswar, B. Webb, M. A. Marti-Renom, M. Madhusudhan, D. Eramian, M.y. Shen, U. Pieper, and A. Sali. Comparative protein structure modeling using Modeller. *Current protocols in bioinformatics*, 15(1):5–6, 2006.
- T. A. Figueiredo, R. G. Sobral, A. M. Ludovice, J. M. F. de Almeida, N. K. Bui, W. Vollmer, H. de Lencastre, and A. Tomasz. Identification of genetic determinants and enzymes involved with the amidation of glutamic acid residues in the peptidoglycan of *Staphylococcus aureus*. *PLoS pathogens*, 8(1):e1002508, 2012.
- S. R. Filipe and A. Tomasz. Inhibition of the Expression of Penicillin Resistance in *Streptococcus Pneumoniae* by Inactivation of Cell Wall Muropeptide Branching Genes. *Proceedings of the National Academy of Sciences of the United States of America*, 97(9):4891–4896, 2000.
- S. R. Filipe, M. G. Pinho, and A. Tomasz. Characterization of the murMN operon involved in the synthesis of branched peptidoglycan peptides in *Streptococcus pneumoniae*. *Journal of Biological Chemistry*, 275 (36):27768–27774, 2000
- S. R. Filipe, E. Severina, and A. Tomasz. The role of murMN operon in penicillin resistance and antibiotic tolerance of *Streptococcus pneumoniae*. *Microbial Drug Resistance*, 7(4):303–316, 2001.
- A. Fiser, R. K. G. Do, et al. Modeling of loops in protein structures. *Protein science*, 9(9):1753–1773, 2000. A. Fiser, S. R. Filipe, and A. Tomasz. Cell wall branches, penicillin resistance and the secrets of the MurM protein. *Trends in Microbiology*, 11(12):547 – 553, 2003.
- M. Fonvielle, I. Li de La Sierra-Gallay, A. H. El-Sagheer, M. Lecerf, D. Patin, D. Mellal, C. Mayer, D. Blanot, N. Gale, T. Brown, et al. The Structure of FemXWv in Complex with a Peptidyl-RNA Conjugate: Mechanism of Aminoacyl Transfer from Ala-tRNAAla to Peptidoglycan Precursors. *Angewandte Chemie*, 125(28):7419–7422, 2013.
- D. Ganchev, H. Hasper, E. Breukink, and B. de Kruijff. Size and orientation of the lipid II headgroup as revealed by AFM imaging. *Biochemistry*, 45(19):6195–6202, 2006.
- J. Garcia-Bustos and A. Tomasz. A Biological Price of Antibiotic Resistance: Major Changes in the Peptidoglycan Structure of Penicillin-Resistant pneumococci. *Proceedings of the National Academy of Sciences of the United States of America*, 87(14):5415–5419, 1990.
- J. Garcia-Bustos, B. T. Chait, and A. Tomasz. Structure of the peptide network of pneumococcal peptidoglycan. *Journal of Biological Chemistry*, 262(32):15400–15405, 1987.
- R. J. Gowers, M. Linke, J. Barnoud, T. J. Reddy, M. N. Melo, S. L. Seyler, J. Domański, D. L. Dotson, S. Buchoux, I. M. Kenney, et al. MDAnalysis: a Python package for the

- rapid analysis of molecular dynamics simulations. In Proceedings of the 15th Python in Science Conference, volume 98. SciPy Austin, TX, 2016.
- J. A. Graham, J. W. Essex, and S. Khalid. PyCGTOOL: automated generation of coarse-grained molecular dynamics models from atomistic trajectories. *Journal of chemical information and modeling*, 57(4):650–656, 2017.
- M. D. Hanwell, D. E. Curtis, D. C. Lonie, T. Vandermeersch, E. Zurek, and G. R. Hutchison. Avogadro: an advanced semantic chemical editor, visualization, and analysis platform. *Journal of cheminformatics*, 4(1):17, 2012.
- S. Hebecker, J. Krausze, T. Hasenkampf, J. Schneider, M. Groenewold, J. Reichelt, D. Jahn, D. W. Heinz, and J. Moser. Structures of two bacterial resistance factors mediating tRNA-dependent aminoacylation of phosphatidylglycerol with lysine or alanine. *Proceedings of the National Academy of Sciences*, 112(34): 10691–10696, 2015.
- S. S. Hegde and J. S. Blanchard. Kinetic and mechanistic characterization of recombinant *Lactobacillus viridescens* FemX (UDP-N-acetylmuramoyl pentapeptide-lysine N6-alanyltransferase). *Journal of Biological Chemistry*, 278(25):22861–22867, 2003.
- R. O. Heuckeroth, L. Glaser, and J. I. Gordon. Heteroatom-substituted fatty acid analogs as substrates for N-myristoyltransferase: an approach for studying both the enzymology and function of protein acylation. *Proceedings of the National Academy of Sciences*, 85(23):8795–8799, 1988.
- J. Huang, S. Rauscher, G. Nawrocki, T. Ran, M. Feig, B. L. de Groot, H. Grubmüller, and A. D. MacKerell Jr. CHARMM36m: an improved force field for folded and intrinsically disordered proteins. *Nature methods*, 14(1):71, 2017.
- S. Jo, T. Kim, V. G. Iyer, and W. Im. CHARMM-GUI: a web-based graphical user interface for CHARMM. *Journal of computational chemistry*, 29(11):1859–1865, 2008.
- S. Jo, X. Cheng, J. Lee, S. Kim, S.J. Park, D. S. Patel, A. H. Beaven, K. I. Lee, H. Rui, S. Park, et al. CHARMM-GUI 10 years for biomolecular modeling and simulation. *Journal of computational chemistry*, 38(15):1114–1124, 2017.
- W. Kabsch. Xds. *Acta Crystallographica Section D: Biological Crystallography*, 66(2):125–132, 2010.
- G. Langer, S. X. Cohen, V. S. Lamzin, and A. Perrakis. Automated macromolecular model building for X-ray crystallography using ARP/wARP version 7. *Nature protocols*, 3(7):1171, 2008.
- T.Y. Lin, W. S. Gross, G. K. Auer, and D. B. Weibel. Cardiolipin Alters *Rhodobacter sphaeroides* Cell Shape by Affecting Peptidoglycan Precursor Biosynthesis. *mBio*, 10(1):e02401–18, 2019.
- A. J. Lloyd, A. M. Gilbey, A. M. Blewett, G. De Pascale, A. El Zoeiby, R. C. Levesque, A. C. Catherwood, A. Tomasz, T. D. Bugg, D. I. Roper, et al. Characterization of tRNA-dependent peptide bond formation by MurM in the synthesis of *Streptococcus pneumoniae* peptidoglycan. *Journal of Biological Chemistry*, 283(10):6402–6417, 2008.
- F. Madeira, J. Lee, N. Buso, T. Gur, N. Madhusoodanan, P. Basutkar, A. Tivey, S. C. Potter, R. D. Finn, R. Lopez, et al. The EMBL-EBI search and sequence analysis tools APIs in 2019. *Nucleic acids research*, 2019.

- M. A. Martí-Renom, A. C. Stuart, A. Fiser, R. Sánchez, F. Melo, and A. Šali. Comparative protein structure modeling of genes and genomes. *Annual review of biophysics and biomolecular structure*, 29(1):291–325, 2000.
- M. Matsushashi, C. P. Dietrich, and J. L. Strominger. Biosynthesis of the Peptidoglycan of Bacterial Cell Walls III. THE ROLE OF SOLUBLE RIBONUCLEIC ACID AND OF LIPID INTERMEDIATES IN GLYCINE INCORPORATION IN STAPHYLOCOCCUS AUREUS. *Journal of Biological Chemistry*, 242(13):3191–3206, 1967.
- C. Morlot, D. Straume, K. Peters, O. A. Hegnar, N. Simon, A.M. Villard, C. Contreras-Martel, F. Leisico, E. Breukink, C. Gravier-Pelletier, et al. Structure of the essential peptidoglycan amidotransferase MurT/GatD complex from *Streptococcus pneumoniae*. *Nature communications*, 9(1):3180, 2018.
- N. A. Morrice, B. Gabrielli, B. E. Kemp, and R. Wettenhall. A cardiolipin-activated protein kinase from rat liver structurally distinct from the protein kinases C. *Journal of Biological Chemistry*, 269(31):20040–20046, 1994.
- G. M. Morris, R. Huey, W. Lindstrom, M. F. Sanner, R. K. Belew, D. S. Goodsell, and A. J. Olson. Autodock4 and AutoDockTools4: automated docking with selective receptor flexibility. *Journal of Computational Chemistry*, 16:2785, 2009.
- D. Münch, T. Roemer, S. H. Lee, M. Engeser, H. G. Sahl, and T. Schneider. Identification and in vitro analysis of the GatD/MurT enzyme-complex catalyzing lipid II amidation in *Staphylococcus aureus*. *PLoS pathogens*, 8(1):e1002509, 2012.
- M. Parrinello and A. Rahman. Polymorphic transitions in single crystals: A new molecular dynamics method. *Journal of Applied physics*, 52(12):7182–7190, 1981.
- B. Peng, N. A. Morrice, L. C. Groenen, and R. E. Wettenhall. Phosphorylation Events Associated with Different States of Activation of a Hepatic Cardiolipin/Protease-activated Protein Kinase STRUCTURAL IDENTITY TO THE PROTEIN KINASE N-TYPE PROTEIN KINASES. *Journal of Biological Chemistry*, 271(50):32233–32240, 1996.
- X. Periole, M. Cavalli, S.J. Marrink, and M. A. Ceruso. Combining an Elastic Network With a Coarse-Grained Molecular Force Field: Structure, Dynamics, and Intermolecular Recognition. *Journal of Chemical Theory and Computation*, 5(9):2531–2543, 2009. doi: 10.1021/ct9002114. URL <https://doi.org/10.1021/ct9002114>. PMID: 26616630.
- S. Pesakhov, R. Benisty, N. Sikron, Z. Cohen, P. Gomelsky, I. Khozin-Goldberg, R. Dagan, and N. Porat. Effect of hydrogen peroxide production and the Fenton reaction on membrane composition of *Streptococcus pneumoniae*. *Biochimica et Biophysica Acta (BBA)-Biomembranes*, 1768(3):590–597, 2007.
- J. Purzycka-Preis and M. Zydowo. Regulatory effect of pig heart phospholipids on heart muscle AMP-deaminase. *The International journal of biochemistry*, 19(6):565–568, 1987.
- I. Ruiz de Mena, E. Mahillo, J. Arribas, and J. Castano. Kinetic mechanism of activation by cardiolipin (diphosphatidylglycerol) of the rat liver multicatalytic proteinase. *Biochemical Journal*, 296(1):93–97, 1993.
- A. Šali and T. L. Blundell. Comparative protein modelling by satisfaction of spatial restraints. *Journal of molecular biology*, 234(3):779–815, 1993.
- M. Scheideler and R. M. Bell. Phospholipid dependence of homogeneous, reconstituted sn-glycerol-3-phosphate acyltransferase of *Escherichia coli*. *Journal of Biological Chemistry*, 264(21):12455–12461, 1989.

- M. Schlame. Thematic Review Series: Glycerolipids. Cardiolipin synthesis for the assembly of bacterial and mitochondrial membranes. *Journal of lipid research*, 49(8):1607–1620, 2008.
- T. Schneider and H.G. Sahl. An oldie but a goodie – cell wall biosynthesis as antibiotic target pathway. *International Journal of Medical Microbiology*, 300(2):161 – 169, 2010.
- T. Schneider, M. M. Senn, B. Berger-Bächi, A. Tossi, H.G. Sahl, and I. Wiedemann. In vitro assembly of a complete, pentaglycine interpeptide bridge containing cell wall precursor (lipid II-Gly5) of *Staphylococcus aureus*. *Molecular microbiology*, 53(2):675–685, 2004.
- K. Sekimizu and A. Kornberg. Cardiolipin activation of dnaA protein, the initiation protein of replication in *Escherichia coli*. *Journal of Biological Chemistry*, 263(15):7131–7135, 1988.
- A. Severin and A. Tomasz. Naturally occurring peptidoglycan variants of *Streptococcus pneumoniae*. *Journal of bacteriology*, 178(1):168–174, 1996.
- J. Shearer, D. Jefferies, and S. Khalid. Outer membrane proteins ompa, fhua, ompf, esta, btub, and ompx have unique lipopolysaccharide fingerprints. *Journal of chemical theory and computation*, 15(4):2608–2619, 2019.
- G. M. Sheldrick. Experimental phasing with SHELXC/D/E: combining chain tracing with density modification. *Acta Crystallographica Section D: Biological Crystallography*, 66(4):479–485, 2010.
- M.y. Shen and A. Sali. Statistical potential for assessment and prediction of protein structures. *Protein science*, 15(11):2507–2524, 2006.
- F. Sievers, A. Wilm, D. Dineen, T. J. Gibson, K. Karplus, W. Li, R. Lopez, H. McWilliam, M. Remmert, J. Söding, et al. Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Molecular systems biology*, 7(1):539, 2011.
- A. M. Smith and K. P. Klugman. Alterations in MurM, a cell wall muropeptide branching enzyme, increase high-level penicillin and cephalosporin resistance in *Streptococcus pneumoniae*. *Antimicrobial agents and chemotherapy*, 45(8):2393–2396, 2001.
- J. M. Smith, C. G. Dowson, and B. G. Spratt. Localized sex in bacteria. *Nature*, 349(6304):29–31, 1991.
- I. G. Tironi, R. Sperb, P. E. Smith, and W. F. van Gunsteren. A generalized reaction field method for molecular dynamics simulations. *The Journal of chemical physics*, 102(13):5451–5459, 1995.
- V. L. Tlapak-Simmons, E. S. Kempner, B. A. Baggenstoss, and P. H. Weigel. The active streptococcal hyaluronan synthases (HASs) contain a single HAS monomer and multiple cardiolipin molecules. *Journal of Biological Chemistry*, 273(40):26100–26109, 1998.
- V. L. Tlapak-Simmons, B. A. Baggenstoss, T. Clyne, and P. H. Weigel. Purification and lipid dependence of the recombinant hyaluronan synthases from *Streptococcus pyogenes* and *Streptococcus equisimilis*. *Journal of Biological Chemistry*, 274(7):4239–4245, 1999a.
- V. L. Tlapak-Simmons, B. A. Baggenstoss, K. Kumari, C. Heldermon, and P. H. Weigel. Kinetic characterization of the recombinant hyaluronan synthases from *Streptococcus pyogenes* and *Streptococcus equisimilis*. *Journal of Biological Chemistry*, 274(7):4246–4253, 1999b.

- V. L. Tlapak-Simmons, C. A. Baron, and P. H. Weigel. Characterization of the purified hyaluronan synthase from *Streptococcus equisimilis*. *Biochemistry*, 43(28):9234–9242, 2004.
- M.C. Trombe, M.A. Lan  elle, and G. Lan  elle. Lipid composition of aminopterin-resistant and sensitive strains of *Streptococcus pneumoniae*. Effect of aminopterin inhibition. *Biochimica et Biophysica Acta (BBA)-Lipids and Lipid Metabolism*, 574(2):290–300, 1979.
- O. Trott and A. J. Olson. AutoDock Vina: improving the speed and accuracy of docking with a new scoring function, efficient optimization, and multithreading. *Journal of computational chemistry*, 31(2):455–461, 2010.
- A. A. Vagin, R. A. Steiner, A. A. Lebedev, L. Potterton, S. McNicholas, F. Long, and G. N. Murshudov. REFMAC5 dictionary: organization of prior chemical knowledge and guidelines for its use. *Acta Crystallographica Section D: Biological Crystallography*, 60(12):2184–2195, 2004.
- G. Van Zundert, J. Rodrigues, M. Trellet, C. Schmitz, P. Kastiris, E. Karaca, A. Melquiond, M. vanDijk, S. De Vries, and A. Bonvin. The haddock2. 2 web server: user-friendly integrative modeling of biomolecular complexes. *Journal of molecular biology*, 428(4):720–725, 2016.
- J. Wang, R. M. Wolf, J. W. Caldwell, P. A. Kollman, and D. A. Case. Development and testing of a general amber force field. *Journal of computational chemistry*, 25(9):1157–1174, 2004.
- T. A. Wassenaar, K. Pluhackova, R. A. Bockmann, S. J. Marrink, and D. P. Tieleman. Going Backward: A Flexible Geometric Approach to Reverse Transformation from Coarse Grained to Atomistic Models. *Journal of Chemical Theory and Computation*, 10(2):676–690, 2014. doi: 10.1021/ct400617g. URL <https://doi.org/10.1021/ct400617g>. PMID: 26580045.
- P. H. Weigel, Z. Kyosseff, and L. C. Torres. Phospholipid dependence and liposome reconstitution of purified hyaluronan synthase. *Journal of Biological Chemistry*, 281(48):36542–36551, 2006.
- C. J. Williams, J. J. Headd, N. W. Moriarty, M. G. Prisant, L. L. Videau, L. N. Deis, V. Verma, D. A. Keedy, B. J. Hintze, V. B. Chen, et al. MolProbity: More and better reference data for improved all-atom structure validation. *Protein Science*, 27(1):293–315, 2018.
- G. Winter. xia2: an expert system for macromolecular crystallography data reduction. *Journal of applied crystallography*, 43(1):186–190, 2010.
- S. Witzke, M. Petersen, T. S. Carpenter, and S. Khalid. Molecular Dynamics Simulations Reveal the Conformational Flexibility of Lipid II and Its Loose Association with the Defensin Plectasin in the *Staphylococcus aureus* Membrane. *Biochemistry*, 55(23):3303–3314, 2016.
- M. H. Wright, W. P. Heal, D. J. Mann, and E. W. Tate. Protein myristoylation in health and disease. *Journal of chemical biology*, 3(1):19–35, 2010.
- A. Zapun, J. Philippe, K. A. Abrahams, L. Signor, D. I. Roper, E. Breukink, and T. Vernet. In vitro reconstitution of peptidoglycan assembly from the Gram-positive pathogen *Streptococcus pneumoniae*. *ACS chemical biology*, 8(12):2688–2696, 2013.

Figure Titles and Legends

Figure 1. Diagram illustrating stages of the PG biosynthesis pathway. 1) The cytoplasmic stage is characterised by the formation of UDP-MurNAc-pentapeptide (UDP-MurNAc-5P) by the Mur ligases. The pentapeptide (5P) stem peptide usually comprises L-Ala- γ -D-Glu-L-Lys-D-Ala-D-Ala in Gram-positive organisms. 2) At the internal face of the cytoplasmic membrane *MraY* catalyses the addition of UDP-MurNAc-5P to undecaprenyl-pyrophosphate forming Lipid I, which is then converted to Lipid II by *MurG*. In *S. pneumoniae*, the second position D-glutamate is α -amidated to D-iso-glutamine (iGln) by the *MurT/GatD* complex, and in some cases a dipeptide branch of either L-Ser/L-Ala or L-Ala/L-Ala may be appended at the ϵ -amino group of the third position lysine by *MurM* and *MurN*, respectively. The exact order of the cytoplasmic membrane steps remains uncertain, but for clarity, in this figure, they appear in a linear format, with conversion to Lipid II occurring before peptide stem modifications, and amidation occurring before branching. Lipid II is translocated across the membrane by *MurJ*. 3) At the external face of the cytoplasmic membrane, PBPs form glycan chains by transglycosylation (TG), with the concomitant release of undecaprenyl-pyrophosphate, and form either direct or indirect cross-links throughout the PG layer via transpeptidation (TP). Nucleotide sugars UDP-GlcNAc and UDP-MurNAc and the sugars GlcNAc and MurNAc are signified by blue, violet, dark blue and purple elongated hexagons respectively. Figure created using BioRender.com.

Figure 2. Cartoon representation of MurM predicted structure. 14 α -helices (red), 12 β -sheet (yellow) and unstructured regions (green). Best model obtained based on SOAP and DOPE scores following homology modelling using MODELLER with *S. aureus* FemX as a template.

Figure 3. Surface representation of MurM binding site. A) *MurM*₁₅₉ model aligned and overlaid with the UDP-MurNAc-pentapeptide substrate which was co-crystallised with *W. viridescens* FemX B) *MurM*₁₅₉ model with truncated Lipid II docked in the binding site, using AutoDock Vina C) *MurM*₁₅₉ model with Lipid II in the binding site, from membrane simulations. Figures were created with PyMOL (Version 2.2.0) and Chimera (Version 1.13.1).

Figure 4. Lipid II binding to the putative MurM binding site. Panels A, B and C show the 3 simulations where Lipid II was found to bind in the putative *MurM* binding site. Each panel shows the *MurM* binding to Lipid II with respect to the membrane (top), and an enlarged image of the *MurM* binding to Lipid II, with the membrane removed (bottom). *MurM* binding site residues F103, K35, W38, R215 and Y219 (yellow), Lipid II head group (red) and Lipid II prenyl chain (blue).

Figure 5. Different conformations of Lipid II inside MurM binding site. *MurM* (grey) with Lipid II binding, coloured on a BWR scale with respect to simulation time, in system 5.

Figure 6. Interactions between MurM and membrane phospholipids. Depletion-enrichment (D-E) indices for phosphatidylethanolamine (PE), phosphatidylglycerol

(PhG) and cardiolipin (CL) occurring within a 1.1 nm perimeter of the MurM protein for A) Systems 4-5 (molar ratio of 75 % phosphatidylethanolamine and 25 % phosphatidylglycerol), B) System 6-7 (molar ratio of 76 % phosphatidylethanolamine, 16% phosphatidylglycerol and 8 % cardiolipin) and C) System 8-9 (molar ratio of 72 % phosphatidylethanolamine, 12 % phosphatidylglycerol and 16 % cardiolipin). The D-E index was determined from 150-250 ns in 50 ns blocks for all repeats for a total of 8 values per plot. D) Example of a depletion-enrichment map with MurM at the membrane. White dots represent the center of geometry of each protein residue, and the percentage enrichment of phospholipid is indicated by the colour. E) Activation of MurM was calculated as the product of subtraction of MurM velocity in the absence of cardiolipin ($v_{0(-C)}$) from MurM velocity in the presence of cardiolipin ($v_{0(+C)}$) divided by $v_{0(-C)}$ and was plotted versus cardiolipin concentration. F) Inhibition of MurM was calculated as $((v_{0(-PhG)}) - (v_{0(+PhG)}))/v_{0(-PhG)} \times 100$ (where PhG denotes phosphatidylglycerol) and was plotted versus phosphatidylglycerol concentration. Data were fitted as described in the text. GraphPad Prism (Version 8.4.1) and Matplotlib (Version 3.0.3) were used for data analysis and figure preparation.

Tables with Titles and Legends

Data collection	FemX
Synchrotron radiation, detector and wavelength (Å)	Pilatus 6M-F, 0.920
Unit cell (a, b, c (Å), α , β , γ (°))	45.01, 83.62, 133.93, 90.0
Space group	P2 ₁ 2 ₁ 2 ₁
Resolution (Å)	52.27-1.62 [1.66-1.62]
Observations	422,822 [29,596]
Unique reflections	65,058 [4,782]
$I/s(I)$	15.7 [2.6]
R_{sym}^a	0.065 [0.567]
R_{meas}	0.078 [0.690]
$R_{p.i.m}$	0.031 [0.273]
Completeness (%)	99.7 [99.8]
Refinement	
Non-hydrogen atoms	3,397 (including 177 waters)
R_{cryst}^b	0.221 [0.262]
Reflections used	61,691 [4,531]
R_{free}^c	0.262 [0.296]
Reflections used	3,294 [244]
R_{cryst} (all data) ^b	0.222
Average temperature factor (Å ²)	26
Rmsds from ideal values	
Bonds (Å)	0.013
Angles (°)	1.5
DPI coordinate error (Å) ^d	0.098
Ramagandran Plot ^e	
Favoured (%)	98.0
Outliers (%)	0.0

Table 1. Summary of crystallographic data collection and refinement statistics from the *S. aureus* FemX structure. The highest resolution bin of data is indicated by square parentheses. Numbers in square parentheses refer to values in the highest resolution shell. ^a $R_{sym} = S_j S_h |I_{h,j} - \langle I_h \rangle| / S_j S_h < I_h \rangle$ where $I_{h,j}$ is the j th observation of reflection h , and I_h is the mean intensity of that reflection. ^b $R_{cryst} = S ||F_{obs}| - |F_{calc}|| / S |F_{obs}|$ where F_{obs} and F_{calc} are the observed and calculated structure factor amplitudes, respectively. ^c R_{free} is equivalent to R_{cryst} for a 4% subset of reflections not used in the refinement (Brünger, 1992). ^d DPI refers to the diffraction component precision index (Cruickshank, 1999). ^e As calculated by Molprobit (Williams et al., 2018).

Membrane	Available	Unavailable	Non-adherence	Lipid II in binding site
1 (0 % CL)	4	1	1	1
2 (12 % CL)	4	1	1	2
3 (16 % CL)	3	3	0	0

Table 2. A breakdown of MurM adherence to, and orientation on, the membrane in three different systems.

STAR Methods

RESOURCE AVAILABILITY

Lead Contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, David Roper (David.Roper@warwick.ac.uk).

Materials Availability

This study did not generate any new unique reagents. Plasmids will be available by Materials Transfer Agreement (MTA) request in line with University of Warwick IP requirements.

Data and Code Availability

No novel code was generated during this work. Modelling scripts and raw data are available from the authors upon request.

METHOD DETAILS

Cloning, overexpression and purification of *S. aureus* FemX

The *S. aureus* Mu50 FemX gene was amplified from chromosomal DNA using Oligonucleotides FemX forward: TTTGCGGGTGGTCTCCCATGGAAAAGATGCATATC ACTAATCAGG and FemX Reverse: TTTGCGCTCGAGGCCCTGAAAATACAGGTTTC TTTTCGTTTTAATTTACGAGATATTTAATTTTAGC. The resulting PCR fragment was cleaved with *BsaI* and *XhoI* and cloned into pET28 between the *NcoI* and *XhoI* restriction sites to create pET28::FemX, containing a Tobacco Etch Virus (TEV) protease cleavable C-terminal hexa-histidine tag. *E. coli* B834 (DE3) harbouring plasmid pRARE2 (which supplies seven rare tRNAs to support expression of genes in *E. coli*) were transformed with pET28::FemX. Transformed *E. coli* B834 (DE3) pRARE2 were used to inoculate M9 media supplemented with all 19 L-amino acids and 40 mM L-selenomethionine in place of L-methionine (Doublié, 1997). Transformants were cultured at 37 °C at 180 rpm until an optical density at 600 nm (OD₆₀₀) of 0.4-0.6 was

reached. Protein expression was induced by 1 mM isopropyl-D-1-thiogalactopyranoside at 25 °C for 4 hours. Cells were harvested by centrifugation at 6,000 xg for 15 minutes and cell pellets containing 4-6 g of cells were resuspended in 20 mL of 50 mM sodium phosphate pH 7.0, 1M NaCl and 2.5 mg.mL⁻¹ lysozyme to which one tablet of Pierce EDTA free Protease Inhibitor was added. The cell suspension was incubated with slow rotation for 30 minutes at 4 °C before disruption using a Bandelin Sonopuls sonicator with 3 x 30 second bursts at 70 % power. The lysate was clarified by centrifugation at 50,000 xg at 4 °C for 30 minutes. FemX was then purified by immobilised metal affinity chromatography (IMAC) using a 5 mL gravity fed column of cobalt Talon resin equilibrated with 50 mL of 50 mM sodium phosphate pH 7.0, 500 mM NaCl, 10 mM imidazole and 20 % (v/v) glycerol (equilibration buffer). Once the 50,000 xg supernatant was loaded onto the column, it was eluted sequentially with 50 mL of equilibration buffer, 30 mL of equilibration buffer with 50 mM imidazole and 30 mL of equilibration buffer with 200 mM imidazole. 10 mL Fractions were analysed by SDS-PAGE and those containing FemX were pooled and concentrated, using a vivaspin 20 centrifugal concentrator column (10,000 molecular weight cut off (MWCO), as required. Size exclusion chromatography in 50 mM sodium phosphate pH 7.0, 500 mM NaCl and 20 % (v/v) glycerol was used to further purify FemX on a Superdex 75 10/300 column. The histidine tag was then removed from the FemX protein by digestion with histidine-tagged TEV protease at a molar ratio of 100:1 FemX : TEV protease at 4 °C overnight. Cleaved and uncleaved protein were separated by a reverse IMAC following the procedure described above.

FemX crystallisation and data collection

FemX was exchanged into 50 mM ethanolamine pH 10.0, 100 mM NaCl and 20 % (v/v) glycerol, concentrated to 15 mg.mL⁻¹ using a vivaspin 20 centrifugal concentrator column with a 10,000 MWCO and screened for suitable crystallisation conditions using a honeybee 963 crystallisation robot against JCSG plus, PACT primer and Morpheus crystallisation screens. Crystals obtained from the Morpheus screen were used directly for data collection experiments, although crystallization conditions were further refined to 0.12 M Ethylene Glycol, 0.1 M MES/imidazole pH 6.3 and 28 % (w/v) Ethylene Glycol-PEG 8000. Crystals were frozen directly for X-ray diffraction data experiments on the I04-1 beamline at the Diamond synchrotron (Didcot, UK) using a Pilatus 6M-F detector. Data were processed automatically using Xia2 (Winter, 2010) to 1.62 Å. Molecular replacement was not successful so selenomethionine containing FemX crystals (FemX-SeMet) were produced as described above and the structure was solved by the single anomalous diffraction method. X-ray data from the FemX-SeMet crystal were collected on the I02 beamline at the Diamond synchrotron (Didcot, UK) using a Pilatus 6M detector. All data were indexed, integrated and scaled using the XDS package (Kabsch, 2010). All 10 of expected selenium atoms in the asymmetric unit were located and refined by the SHELX suite (Sheldrick, 2010). These sites were used to obtain preliminary phases. The starting model was built by ARP/wARP (Langer

et al., 2008). This model was used to refine the higher resolution data. The structure was refined using iterative cycles of REFMAC (Vagin et al., 2004) and model building/solvent addition with COOT (Emsley et al., 2010).

Cloning, Overexpression and Purification of *S. pneumoniae*(159) MurM

As described in Lloyd et al. (2008), the MurM allele of *S. pneumoniae*(159) was cloned with a C-terminal histidine tag into pET21b and over-expressed in *E. coli* C41 (DE3)/pRIL. Cells were harvested by centrifugation and treated with 2.5 mg.mL⁻¹ hen egg white lysozyme prior sonication. MurM was solubilised with 1 M NaCl and fractionated between 25 % and 50 % of saturation ammonium sulfate followed by purification by size exclusion chromatography with Sephacryl S-200 and by immobilized metal affinity chromatography (IMAC) using cobalt Talon resin. The purity and identity of the final products of these purifications were assessed by SDS-PAGE.

Cloning, Overexpression and Purification of *S. pneumoniae* (Pn16) AlaRS

As described in Lloyd et al. (2008), the AlaRS allele of *S. pneumoniae*(Pn16) was cloned into pET26a and over-expressed in *E. coli* BL21(DE3) star/pRARE. The soluble protein was purified using nickel-chelated Chelating sepharose, desalted and further purified by anion exchange chromatography on a 0.98 ml MonoQ™, column. The purity and identity of the final products of these purifications were assessed by SDS-PAGE.

Preparation of MurM substrates

The substrates used for assays of MurM were prepared as follows:

Lipid II(Lys): The peptidoglycan intermediate Lipid II(Lys) (undecaprenyl pyrophosphoryl N-N-acetyl muramyl (N-acetyl glucosaminy) L-alanyl-γ-D-glutamyl-L-lysyl-D-alanyl-D-alanine) was prepared by re-capitulation of the peptidoglycan synthesis pathway as described (Lloyd et al. 2008).

[³H]-Alanyl-tRNA^{Ala}: *Micrococcus flavus* tRNA was isolated from cell pellets of *M. flavus* cultures grown to late exponential phase by phenol extraction followed by isopropanol precipitation, anion exchange chromatography and ethanol precipitation as described by Zubay (1962) as adapted by Lloyd et al. (2008). tRNAs were renatured in 2 mM MgCl₂ at 60°C and aminoacylated with [2,3-³H]-L-alanine as described by Lloyd et al. (2008) and quantitated by liquid scintillation counting.

***S. pneumoniae* MurM enzymology**

MurM was assayed as described by Lloyd et al. (2008) in duplicate in a final volume of 35 µl of 50 mM 3-(N-morpholino)-propane sulphonic acid adjusted to pH 6.8, 30 mM KCl, 10 mM MgCl₂, 1.5 % (w/v) CHAPS (Assay Buffer), 1 mM DTT, 1 mM L-alanine, 10 µM Lipid II-Lys and 24.3 nM MurM. Reactions were initiated by the addition of 0.45 M [³H]-alanyl-tRNA^{Ala} (1000 cpm.pmol⁻¹) and were incubated at 37 °C for two minutes, over which time frame, product accumulation was linear with respect to time. Where the impact of cardiolipin or phosphatidylglycerol on MurM activity was assessed, the required amounts of 10 mg/mL stocks of each phospholipid in ethanol or

chloroform/methanol (49:1) were dried down in the reaction vials the assays were to be performed in, and solubilised by addition of assay buffer. Reactions were terminated by the addition of 35 µl of ice-cold 6 M pyridinium acetate pH 4.5 and 70 µl ice-cold n-butanol. The incubations were rapidly mixed and centrifuged for 5 minutes at 1 °C at 13,000 *xg*, after which time the n-butanol phase was washed with 70 µl of water and then assayed for [³H]-Lipid II-L-Ala by liquid scintillation counting. Tritium counts accumulated in control reactions performed without Lipid II(Lys) were subtracted from corresponding data acquired in the presence of this substrate. MurM activities in the presence of phospholipid were related to the activity of the enzyme in the absence of phospholipid and plotted as fold activation or percentage inhibition vs phospholipid concentration. The data were then fitted using GraphPad Prism (Version 8.4.1) to either of equations 1 or 2 as appropriate:

Equation 1

$$Fold\ Activation = \frac{Maximum\ Activation \cdot [Cardiolipin]^h}{S_{0.5(Activation)}^h + [Cardiolipin]^h}$$

Equation 2

$$\% Inhibition = \frac{100 \cdot [Phosphatidylglycerol]^h}{IC_{50}^h + [Phosphatidylglycerol]^h}$$

Maximum activation and $S_{0.5}$ (Activation) (Equation 1) corresponded to the degree of activation at infinite cardiolipin concentration and the cardiolipin concentration required to elicit half maximal activation respectively. IC_{50} (Equation 2) corresponds to the phosphatidylglycerol concentration that elicited half maximal inhibition. For both equations, h denoted the Hill coefficient.

Computational Studies Overview

A number of computational techniques were used in this study, to assist the reader in understanding the logistics of these methods, we have provided a summary flowchart (Supplemental Information: Figure S1).

Homology Modelling of MurM

Due to the natural ability of streptococci to undergo homologous recombination, *S. pneumoniae* MurM genes are highly mosaic, and so, in line with the enzymology studies, the MurM sequence used for homology modelling was that of *S. pneumoniae* MurM₁₅₉. *S. pneumoniae* MurM, *S. aureus* FemX (PDB ID: 6SNR) and FemA (PDB ID: 1LRZ), and *Weissella viridescens* FemX (PDB ID: 3GKR) were aligned (Supplemental

Information: Figure S2) by multiple sequence alignment using Clustal Omega to determine sequence identity (Sievers et al., 2011).

The structure of *S. aureus* FemX was used as the template for homology modelling due to its high relatedness with MurM. *S. aureus* FemX and MurM₁₅₉ sequences were aligned, and using MODELLER (Eswar et al., 2006, Martí-Renom et al., 2000, Šali and Blundell, 1993, Fiser et al., 2000) a test model was generated to verify the validity of the template and the alignment. This model was evaluated by computing its energy profile according to the DOPE-HR (high-resolution version of the Discrete Optimized Protein Energy) (Shen and Sali, 2006), smoothed via window averaging with a size of 15 residues. The profiles of template and model were compared (Supplemental Information: Figure S3), and further refinement was conducted in the region between Lys230 and Pro299, as well as in all loop regions. This optimisation was conducted by performing a very slow MD annealing on the selected regions, whilst maintaining the structure of other ones. The scoring of the resulting conformations was obtained via a function built specifically to evaluate the geometry of loops. For this step, 64 different base models were created and their secondary structure was refined independently 16 times. The resulting 1024 models were evaluated and ranked using DOPE-HR as well as the SOAP (Statistically Optimized Atomic Potentials) (Dong et al., 2013). The 10 best scoring models for each score were selected and evaluated based on the number of physical constraint violations present.

The best model of MurM₁₅₉ was aligned with the previous MurM model (Fiser et al., 2003) or *W. viridescens* Femx homologues (Fonvielle et al., 2013, Biarrotte-Sorin et al., 2004) for visualisation and analysis in PyMOL (Version 2.1.0).

Molecular docking of truncated Lipid II to MurM

A truncated Lipid II substrate (Supplemental Information: Figure S4) was created for initial molecular docking simulations. The truncated Lipid II was drawn in ChemDraw Professional (Version 17.1) and converted to a pdb file using Avogadro (Version 1.2.0). To prepare the ligand file for docking, the protonation state in H₂O at pH 7.4 was computed. Subsequently the equilibrium geometry minimizing the potential energy was computed using the general amber force field (GAFF) (Wang et al., 2004) from within the Avogadro2 software (Hanwell et al., 2012). Molecular docking was conducted using AutoDock Vina (Trott and Olson, 2010), for which pdbqt files were generated from the pdb files of receptor model and ligands using AutoDock Tools (Morris et al., 2009). Initially the location of the binding site was verified by providing the algorithm with a search space that included the entire protein. Docking was then repeated by restricting the search space to the identified binding site, in order to obtain the final docked conformation.

Molecular docking of Lipid II, Cardiolipin, Phosphatidylglycerol and Phosphatidylethanolamine to the putative MurM binding site.

Using the HADDOCK web server (Van Zundert et al., 2016), full length Lipid II, Cardiolipin and Phosphatidylglycerol and Phosphatidylethanolamine were docked into the putative MurM binding site, where the binding site is defined by the residues Lys35, Trp38, Phe103, Arg215 and Tyr21 (Van Zundert et al., 2016).

Coarse-grained molecular dynamics simulations

All coarse-grained simulations were carried out with the GROMACS package (Version 2018) and the Martini (Version 2.2) forcefield (Abraham et al., 2015, de Jong et al., 2012). Simulations at the coarse-grained and atomistic resolutions were carried out at 313 K. For coarse-grained simulations, a stochastic velocity rescale thermostat with a coupling constant of 1:0 ps controlled the temperature.

The coordinates of the MurM homology model were used to generate a coarse-grained model using the 'martinise.py' script (de Jong et al., 2013). The protein was coarse grained to the ElnDyn model (Periole et al., 2009) with an elastic network strength and cutoff of 500 kJmol⁻¹nm⁻² and 0.9 nm, respectively. The Lipid II model for inclusion in the membrane was parameterised using a united atom model (Gromos 53a6) generated by the Auto-mated topology builder (ATB) web-interface. Following this, the coarse-grained mapping was decided iteratively, and the bonded terms fitted with PyCGTOOL (Graham et al., 2017).

Since the pneumococcal membrane comprises a complex mixture of lipids, a simplified membrane composition was required for the simulations. In order to elucidate the effects of phosphatidylglycerol and cardiolipin on MurM, a non-pneumococcal lipid, phosphatidylethanolamine, was used as the majority lipid. Simulations were conducted with three different membrane systems (Supplemental Information: Table S1). System 1 comprised phosphatidylethanolamine and phosphatidylglycerol in a molar ratio of 75 % and 25 % respectively, system 2 contained phosphatidylethanolamine, phosphatidylglycerol and cardiolipin in a molar ratio of 76 %, 16 % and 8 % respectively and system 3 comprised phosphatidylethanolamine, phosphatidylglycerol and cardiolipin at a molar ratio of 72 %, 12 % and 16 % respectively. The membrane systems of size ~16x16x11.5 nm were generated with the CHARMM-GUI web interface (Jo et al., 2017). Each system was relaxed with a series of minimisation and equilibration steps with timesteps of 5-20 fs, for up to 30 ns. The equilibration steps utilised a semi-isotropic Berendsen barostat, with a 4:0 ps coupling constant (Berendsen et al., 1984). Following equilibration, Lipid II molecules (10 in total) were added to each membrane. The systems were then minimised and equilibrated (for 10 ns), followed by a 2 μ s production run to ensure sufficient mixing of all the lipid components. All production runs were carried out using a 10 fs timestep and a Parrinello-Rahman semi-isotropic barostat with a 12 ps coupling constant (Parrinello and Rahman, 1981). The Lennard-Jones potential was cutoff using the Potential shift

Verlet scheme at long ranges. The reaction field method (Tironi et al., 1995) was used for electrostatics calculations, with dielectric constants of 15 and infinity for charge screening in the short- and long-range regimes, respectively. The short-range cutoff for non-bonded and electrostatic interactions was 1:2 nm. Once lipid mixing was ensured, the size of each system was increased to ~32 nm in the dimension perpendicular to the membrane and MurM was added in a random orientation around 8 nm above each membrane. Biologically relevant salt concentrations (0.15 M NaCl) were added and 10 % of the water molecules were changed to antifreeze particles to prevent localised freezing during simulations. After minimisation and 1 ns of equilibration, during which the protein backbone was restrained with 1000 kJmol⁻¹nm⁻² harmonic restraints, 6x5 μ s production runs were generated per membrane composition (Supplemental Information: Table S1).

All-atom molecular dynamics simulations

Atomistic simulations were conducted using the CHARMM36m forcefield (Huang et al., 2017). The Lipid II model used here was also used in previous work (Witzke et al., 2016), while all other lipid models were obtained from the CHARMM-GUI membrane builder module (Jo et al., 2008). For each coarse-grained membrane system, two repeats were chosen where: 1) the last frame of the production run had a distinct orientation of MurM, relative to the membrane 2) MurM adhered to the membrane surface (Supplemental Information: Table S2). The last frame of the chosen coarse-grained repeats were then backmapped to the all-atom model, using the backward script (Wassenaar et al., 2014). Unfavourable ring conformers were corrected by carrying out minimisation and equilibration steps with dihedral restraints of 25000 kJmol⁻¹rad⁻² on key ring torsions. After the transformation was carried out, each system was cropped in the z dimension to a height of 16:5 nm, to remove unnecessary H₂O molecules.

Each system was minimised and equilibrated for a total of 1 ns, while the backbone of the protein was restrained with 1000 kJmol⁻¹nm⁻² harmonic restraints. Two production runs of 250 ns were carried out for each system. During the production runs a timestep of 2 fs was used, and the pressure (1 bar) regulated with a semi-isotropic Parrinello-Rahman barostat, with a coupling constant of 5:0 ps. The Lennard-Jones potential was cutoff with the Force-switch modifier from 1:0 to 1:2 nm. The short range cutoff for the electrostatic interaction was also 1:2 nm and the Particle mesh Ewald (PME) algorithm (Darden et al., 1993) was used for the long-range regime.

Analysis was carried out over the final 100 ns of each simulation, unless stated otherwise. All simulations were visualised using Visual Molecular Dynamics (VMD) or PyMOL (Version 2.2.0). Other analysis tools were written with a combination of GROMACS tools and in house scripts, that utilised the python module MDAnalysis

(Gowers et al., 2016). The depletion/enrichment (D-E) indices were determined by first counting the number of lipids with a centre of geometry within 1.1 nm of the protein and then comparing this number to the number expected in the bulk of the membrane, using the procedure described by Corradi et al. (2018). The D-E index was obtained by dividing the lipid composition in the 1.1 nm shell around the protein by the bulk membrane composition. Thus a D-E index >1 indicates enrichment, while a D-E index <1 indicates depletion. The D-E index was determined for the last 100 ns of each simulation in 50 ns blocks for all repeats. For a given membrane composition, 8 D-E indices were obtained for each lipid, from which the average and standard errors were calculated. The enrichment maps were generated by first determining the 2D density map of the membrane using the GROMACS tool densmap. Following this, the enrichment percentage was determined using the procedure described by Corradi et al. (2018). An enrichment percentage $<0\%$ indicated that local membrane composition was depleted with respect to the bulk membrane composition. The code for the 2D enrichment maps and D-E indices was reported in Shearer et al. (2019).

QUATIFICATION AND STATISTICAL ANALYSIS

Crystallographic statistics were calculated using software/programs as described in the methods, and values are reported in Table 1. Enzyme assays were performed in duplicate, generating data that differed by no more than 10%. Average values were then plotted. The standard errors of the fits of constants defining the relationships between the response of MurM to phospholipid and phospholipid concentration to equations 1 and 2 were calculated by GraphPad Prism (Version 8.4.1).

Supplemental Information titles and legends

Table S1. Summary of course grained simulations

Table S2. Summary of atomistic simulations. CG label refers to the coarse grained system (Table S1) from which the simulations was constructed.

Figure S1. Flow chart showing the logistics of the computational studies

Figure S2. Sequence alignment showing putative MurM binding site residues.

Alignment of *Weissella viridescens* FemX, *Streptococcus pneumoniae* MurM, *Staphylococcus aureus* FemA and *Staphylococcus aureus* FemX using CLUSTAL Omega (1.2.4). The sequence identity between MurM and *Staphylococcus aureus* FemA, *Staphylococcus aureus* FemX and *Weissella viridescens* FemX was 20.25 %, 26.93 % and 24.38 % respectively. Residues of the putative MurM binding site, proposed to interact with the Lipid II substrate are indicated by red boxes.

Figure S3. Discrete Optimized Protein Energy Profile for MurM and FemX.

Comparison of DOPE-HR profiles for MurM model (red) and FemX template (green).

Figure S4: Structures of lipids investigated in these studies. A) UDP-MurNAc-pentapeptide (Lysine variant), B) Lipid II, C) Truncated Lipid II structure where the C55 prenyl chain has been replaced with a methyl group, D) Cardiolipin, E) Phosphatidylglycerol and F) Phosphotidylethanolamine. All Lipid II precursors and variants contain L-Lysine at the third position of the pentapeptide chain. Structures produced in ChemDraw (Version 19.1).

Figure S5. The remaining four highest scoring poses from molecular docking of truncated Lipid II to MurM using AutoDock Vina. All possessed identical binding affinities of $-7.3 \text{ kcal.mol}^{-1}$. A) and B) show the phosphate group located near the entrance of the cavity, with the pentapeptide located deeper into the pocket. C) and D) show orientations that are not considered possible, since the phosphate group would be linked to the membrane embedded Lipid II, and this would prevent the phosphate from being located deep in the binding site as shown.

Figure S6. Top 3 binding poses for the docking of different lipids to the MurM binding site. Docking of A) Lipid II, B) Cardiolipin, C) Phosphatidylglycerol and D) Phosphotidylethanolamine to the MurM binding site, where residues F103, K35, W38, R215 and Y219 are shown in yellow.

Figure S7. The association of coarse-grained MurM to the surface of the membrane. A) The minimum distance between MurM and the membrane surface for System 1 (top), System 2 (middle) and System 3 (bottom). Snapshots taken of the B) first and C) last frame of repeat 2 (r2), for System 1 (Supplemental Information: Table S1) Colour key: red = Lipid II, blue = protein, and grey = membrane.

Figure S8. Electrostatic surface representation of MurM. A) MurM showing the proposed Lipid II binding site to be positively charged (blue) B) MurM rotated 180° , showing a negatively charged surface patch. Figure prepared in PyMOL (Version 2.2.0) using the APBS Electrostatics Pluggin.

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6

