# A Randomized Controlled Trial of a Psycho-Education Intervention by Midwives in Reducing Childbirth Fear in Pregnant Women

Toohill, J., Fenwick, J., Gamble, J., Creedy, D. K., Buist, A., Turkstra, E. & Ryding, E. L.

Published PDF deposited in Coventry University's Repository

### **Original citation:**

Toohill, J, Fenwick, J, Gamble, J, Creedy, DK, Buist, A, Turkstra, E & Ryding, EL 2014, 'A Randomized Controlled Trial of a Psycho-Education Intervention by Midwives in Reducing Childbirth Fear in Pregnant Women', Birth, vol. 41, no. 4, pp. 384-394. https://dx.doi.org/10.1111/birt.12136

DOI 10.1111/birt.12136 ISSN 0730-7659 ESSN 1523-536X

Publisher: Wiley

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

## A Randomized Controlled Trial of a Psycho-Education Intervention by Midwives in Reducing Childbirth Fear in Pregnant Women

Jocelyn Toohill, RM, MMid, Jennifer Fenwick, RM, PhD, Jenny Gamble, RM, PhD, Debra K. Creedy, RN, PhD, Anne Buist, MD, FRANZCP, Erika Turkstra, PhD, and Elsa-Lena Ryding, MD, PhD

ABSTRACT: Background: Childbirth fear is associated with increased obstetric interventions and poor emotional and psychological health for women. The purpose of this study is to test an antenatal psycho-education intervention by midwives in reducing women's childbirth fear. **Methods:** Women (n = 1,410) attending three hospitals in South East Queensland, Australia, were recruited into the BELIEF trial. Participants reporting high fear were randomly allocated to intervention (n = 170) or control (n = 169) groups. All women received a decision-aid booklet on childbirth choices. The telephone counseling intervention was offered at 24 and 34 weeks of pregnancy. The control group received usual care offered by public maternity services. Primary outcome was reduction in childbirth fear (WDEQ-A) from second trimester to 36 weeks' gestation. Secondary outcomes were improved childbirth self-efficacy, and reduced decisional conflict and depressive symptoms. Demographic, obstetric & psychometric measures were administered at recruitment, and 36 weeks of pregnancy. Results: There were significant differences between groups on postintervention scores for fear of birth (p < 0.001) and childbirth self-efficacy (p = 0.002). Decisional conflict and depressive symptoms reduced but were not significant. Conclusion: Psycho-education by trained midwives was effective in reducing high childbirth fear levels and increasing childbirth confidence in pregnant women. Improving antenatal emotional well-being may have wider positive social and maternity care implications for optimal childbirth experiences. (BIRTH 41:4 December 2014)

**Key words:** childbirth fear, midwife counseling, psycho-education, childbirth self-efficacy, depression, decisional conflict, RCT

Trial Registration: Australian New Zealand Controlled Trials Registry ACTRN12612000526875.

Address correspondence to Jocelyn Toohill, School of Nursing & Midwifery, Griffith University, Logan Campus, University Drive, Meadowbrook. Qld 4131, Australia.

Accepted August 23, 2014

© 2014 The Authors. Birth Published by Wiley Periodicals, Inc. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Jocelyn Toohill is a PhD candidate at Griffith Health Institute, School of Nursing & Midwifery, Griffith University, Brisbane, Qld, Australia; Jennifer Fenwick is a Professor of Midwifery and Clinical Chair at Griffith Health Institute, School of Nursing & Midwifery, Griffith University, Brisbane, Old, Australia, Gold Coast University Hospital, Parkwood, Qld, Australia; Jenny Gamble is Acting Head of School & Professor of Midwifery at Griffith Health Institute, School of Nursing & Midwifery, Griffith University, Brisbane, Qld, Australia; Debra K. Creedy is Professor of Perinatal Mental Health at Griffith Health Institute, School of Nursing & Midwifery, Griffith University, Brisbane, Qld, Australia; Anne Buist is a Professor & Director at University of Melbourne, Melbourne, Vic, Australia; Erika Turkstra is a Senior Research Fellow at Griffith Health Institute, School of Medicine, Griffith University, Brisbane, Qld, Australia; Elsa-Lena Ryding is an Associate Professor at Department of Women's and Children's Health, Division of Obstetrics and Gynecology, Karolinska Institutet, Stockholm, Sweden.

High rates of childbirth fear have been reported in Scandinavian countries (1-5), Australia (6-8), and the United Kingdom (9). Childbirth fear has been linked to adverse maternal outcomes including poor postpartum mental health and high rates of cesarean (10-15). Outside Sweden, no reported systematic approach was seen to identify and care for fearful women (16,17). Without appropriate intervention, women reporting high fear may perceive surgery as their only birth option (18). Although intervention studies have focused on improving vaginal birth rates, no trials have reported on the effectiveness of an antenatal intervention to reduce fear levels before birth (19,20). This paper reports on the effects of an intervention to reduce fear in pregnant women and assist their preparation for a positive birth.

#### Aim

To test an antenatal psycho-education counseling intervention by midwives in reducing women's childbirth fear.

#### **Hypotheses**

Relative to women in the control group, women receiving the psycho-education intervention will report lower levels of childbirth fear at 36 weeks; improved birth confidence, less decisional conflict; and less depressive symptoms.

#### Method

Participants were drawn from a multisite randomized controlled trial. The protocol for the study (known as BELIEF: Birth Emotions: Looking to Improve Expectant Fear) has been published (21). A two arm nonblinded parallel trial design was used. Participants were stratified by hospital site and parity and allocated to study groups using a web-based randomization service to generate blocks for groups of ten. A research assistant accessed the service after receiving a participant's written consent and completed baseline measures. A midwife providing the intervention was notified of the woman's contact details. Ethical approval was obtained from the university and participating hospitals.

#### Participants and Setting

Women in their second trimester attending antenatal clinics of three hospitals in South East Queensland, 385

Australia, able to communicate sufficiently in English, and aged 16 years or older were recruited by research assistants. Participants were screened for high childbirth fear using the Wijma Delivery Expectancy/Experience Questionnaire Version A (W-DEQ A) (22). Women requiring an interpreter, younger than 16 years, or more than 24 weeks pregnant, and anticipating or experiencing a perinatal death (e.g., congenital abnormality incompatible) or stillbirth were excluded. Participants (n = 1,410) were recruited from May 2012 to June 2013. Three hundred and thirty-nine women (n = 339)reporting high fear (defined as a score of  $\geq 66$  on the W-DEO A) were allocated to the intervention (n = 170)or control (n = 169) groups. Two women were incorrectly randomized (W-DEO A scores < 66) and removed from the analysis. This paper reports on primary and secondary antenatal outcomes for women who returned data at 36 weeks' gestation. Baseline characteristics of participants were similar to the national Australian birthing population and have been reported (8).

#### Sample Size

The sample size was determined on a reduction in level of fear between the intervention group and the control group. The statistical package R-Project version 2.14.2 (23) was used to calculate a meaningful reduction in level of childbirth fear based on the study of Fenwick, Gamble, Nathan, Bayes, and Hauck (6) who found one in four women experienced childbirth fear. The calculation also included a standard error of measure to determine how an individual's own score would change over time. A standard deviation of 20 and reliability of 0.87 based on the W-DEQ A (2,22) provided a standard error of measurement to be 7.211. For an 80 percent chance to detect a 10-point difference in WDEQ scores from baseline to around 36 weeks' gestation in the intervention compared with the control group, 140 women plus 30 percent allowance for attrition were needed. The estimate was based on a two-tailed  $\alpha = 0.05$ .

#### Measures

Demographic, obstetric details, and birth preference were collected. The primary outcome measure was a reduction in childbirth fear as measured by the W-DEO A. Secondary outcomes measures included the Childbirth Self-Efficacy Inventory (CBSEI) (24); Edinburgh Postnatal Depression Scale (EPDS) (25), which has been validated for use in pregnancy (26); and the Decisional Conflict Scale (DCS) (27) (Fig. 1). Data were collected at recruitment and 36 weeks of pregnancy.

#### Intervention

BELIEF is a telephone psycho-education counseling intervention offered by midwives. The intervention aims to review women's current expectations and feelings around fear of childbirth, support the expression of feelings, and provide a framework for women to identify and work through distressing elements of childbirth. The intervention was adapted from a midwifery counseling framework for distressed postpartum

WDEQ	The W-DEQ A measures fear of childbirth.
	Respondents rate 33 expectations and experiences
	before birth (version A) and after birth (version B)
	(22). Questions are presented in positive and
	negative formats on a six-point Likert scale from 0
	to 5 requiring reverse scoring of positively
	formulated questions. A score equal to or lower
	than 37 is considered low fear, a score between 38
	and 65 equates to moderate fear, and a score equal
	to or higher than 66 represents a high level of fear
	(2). In this current study, the W-DEQ A had a
	Cronbach's alpha coefficient of 0.94.

- CBSEI The 62-item CBSEI is used to screen childbirth self-efficacy and comprises two subscales in both first stage of labor (total 30 questions) and during the pushing stage of labor (total 32 questions). Within these two stages of labor women's belief that a certain behavior would lead to a given outcome (expectancy outcome), and women's conviction that she could indeed perform that behavior (self-efficacy) are measured to indicate levels of high or low childbirth confidence (24). In this current study, Cronbach's alpha coefficient was 0.98.
- EPDS The 10-item EPDS is widely used to screen for probable antenatal and postnatal depression (25,53). Women selected one of four possible responses to each question. Each item is scored from 0–3 and summed to produce a total score. The range of scores is from 0–30 with higher scores indicating more negative feelings. EPDS scores of > 12 in the antenatal or postnatal period have been recommended as an indicator of probable depression but not diagnostic of depression (54). In this current study, the Cronbach's alpha coefficient was 0.87.
- DCS The DCS is a validated 16-item scale, which investigates concepts that can compromise or facilitate effective decision making (27). A score of 25 or lower (out of a possible score of 100) on the DCS indicates an ability to make decisions (27). Scores ≥ 37.5 are associated with decision delay or feeling unsure about implementing decisions. Participants identify on a 5-point Likert scale of 1 = strongly disagree to 5 = strongly agree their certainty or uncertainty about decisions when there are trade-offs of outcomes in a health-related decision. High internal consistency for the DCS was found in the present study with a Cronbach's alpha coefficient of 0.98.

Fig. 1. Validated measures used in the study.

women developed by Gamble and Creedy (28). The BELIEF intervention aimed to help women develop individual situational supports for the present and near future, affirming that negative things can be managed with a simple plan developed to achieve this. Women in the control group received usual care offered by publicly funded maternity services in Australia. All participants received a copy of a decision-aid booklet titled *Having a baby in Queensland* (29).

#### Statistical Methods

Data were analyzed using the Statistical Package for the Social Sciences (SPSS) version 21.0 for Windows (30). An alpha level of 0.05 was used for all statistical tests. A one-way between-groups analysis of covariance compared the effectiveness of the intervention in reducing the primary outcome of childbirth fear (W-DEQ A). The independent variable was group allocation, and dependent variable was participants' fear score. Women's baseline fear score at recruitment was used as the covariate.

Secondary outcomes were conducted using independent samples t-test. Change scores for fear (W-DEQ A), decisional conflict (DCS), and depression (EPDS) were calculated (Time 1-Time 2) for both groups comparing posttreatment scores for the intervention and control groups with the baseline score as the covariate. Calculation of change scores for childbirth confidence (CBSEI) were Time 2-Time 1 because of inverse scoring compared with the other measures. Effect size statistics (Cohen's d) were determined by subtracting the mean change score for the two groups and dividing by the pooled baseline SD (31). Furthermore, the chi square statistic compared percentages of cases in the intervention and control groups who improved or showed no improvement/deteriorated for secondary outcomes to illustrate individual response to the treatment rather than mean scores for each group. Bonferroni adjustment was used to determine significance levels for group comparisons (32).

#### Results

No significant differences were reported between women returning all data (n = 198) and women lost to follow-up (n = 141) for age, country of birth, parity, previous mode of birth, marital status, DCS, EPDS, CBSEI, and W-DEQ A scores. However, statistically significant differences were found for education and household income (Table 1). Women lost to follow-up were less likely to be educated beyond Year 12, and less likely to have a household income of more than \$AUD 52,000 per annum in comparison to women who completed the study.

One hundred and ninety-eight (n = 198, 58%) eligible women completed the follow-up questionnaire at 36 weeks. Baseline comparison of women randomized to intervention and control arms of the study are presented in Table 2. One hundred and thirty-nine (n = 139) women did not complete the second questionnaire; of these, 45 withdrew (2 because of late pregnancy loss), and 94 were lost to follow-up (including four who gave birth prematurely). The Consolidated Standards of Reporting Trials (CONSORT) participant flow diagram is presented in Fig. 2.

Of the 170 women allocated to the intervention group, 101 (59.4%) returned data. Of these, three women (2.9%) did not receive the intervention, six women (5.9%) completed one session, and 92 (91%) received both sessions of the intervention as planned. The mean

#### Internal Reliability of Scales

Reliability of measures was assessed using baseline scores. Cronbach's alpha coefficient for each scale indicated good internal consistency (as shown in Table 3).

#### Effect of the Intervention on Childbirth Fear

After adjusting for preintervention scores, a significant difference resulted between groups on postintervention W-DEQ A scores for fear of birth, (F(1, 191) = 11.6, p = 0.001, partial eta squared = 0.06) with medium

	Participants returning data at 36 weeks $n = 198$	Participants not returning data at 36 weeks $n = 141$	
Characteristics	No. (%)	No. (%)	р
Age (years) Mean [SD, range]	29.1 [5.44, 17–51]	27.9 [5.89, 17–40]	0.06
No partner	11 (5.6)	15 (10.6)	0.13
Education Year 12 or less	84 (42.4)	82 (58.2)	0.006
Income			
\$0-\$51,999	53 (26.8)	55 (39)	0.04
\$52,000-\$77,999	55 (27.8)	34 (24.1)	
> \$78,000	85 (42.9)	46 (32.6)	
Missing	5 (2.5)	6 (4.3)	
Foreign born	56 (28.3)	35 (24.8)	0.56
Aboriginal	4 (2)	4 (2.8)	0.90
Nulliparous	116 (58.6)	75 (53.2)	0.38
Cesarean last birth			
No labor/Cesarean	9 (31)	10 (43.5)	0.52
Labor/Cesarean	20 (69)	13 (56.5)	
Preferred mode of birth			
Cesarean	31 (16.1)	23 (16.4)	1.00
Missing	4 (2)	1 (< 1)	
W-DEQ Mean [SD, range]	78.3 [11.8, 66–127]	77.8 [11.5, 63–128]	0.69
Missing	3 (1.5)	4 (2.8)	
CBSEI Mean [SD, range]	377 [113.4, 62–620]	372 [126.5, 62–620]	0.71
Missing	6 (3)	9 (6.4)	
DCS Mean [SD, range]	40.23 [22.11, 0–100]	40.32 [23.29, 0–100]	0.89
Missing	2 (1)	3 (2)	
EPDS Mean [SD, range]	8 [5.35, 0–24]	7.7 [5.22, 0–23]	0.65

Table 1. Comparison of Randomized Participants Who Returned Data Versus Those Who Did Not Return Data

*W-DEQ* = Wijma Delivery Expectancy Scale; CBSEI = Childbirth Self-Efficacy Inventory; DCS = Decisional Conflict Scale; EPDS = Edinburgh Postnatal Depression Scale.

effect (33). Women receiving the intervention reported reduced childbirth fear at 36 weeks compared to women in the control group.

#### **Outcome Measurement Change Scores**

Change scores for the primary outcome of fear (W-DEQ A) and the secondary outcomes of childbirth self-efficacy (CBSEI), decisional conflict (DCS), and depression (EPDS) were calculated for both groups (Table 4). Statistically significant differences in fear (p < 0.001) and childbirth confidence (p = 0.002) were found between groups (Bonferroni significance levels) (32). Although there was a reduction in decisional conflict and depression symptoms, there was no statisti-

cally significant difference (p = 0.09; p = 0.38, respectively) between groups (Table 4).

To date, there is insufficient evidence to determine if fear naturally increases or decreases across pregnancy; however, a reduction in childbirth fear scores were found for women in the control group at 36 weeks compared to baseline scores in this study (Control Group: Mean W-DEQ A: 75.7 to 66.5). A 20-point difference in W-DEQ scores delineates high (W-DEQ  $\geq$  66), severe (W-DEQ  $\geq$  85), and extreme childbirth fear (W-DEQ  $\geq$  100) (2,10). Therefore, a change in 20 points would indicate a clinically meaningful change because of decreasing severity of childbirth fear, and was chosen for chi square analysis to determine the percentage of participants who improved versus no improvement. More women in the intervention group

Table 2. Baseline Participant Characteristics for T	Freatment and	Control	Groups
---	---------------	---------	--------

Characteristic	Randomized to treatment group N = 101 No. (%)	Randomized to control group N = 97 No. (%)
Age (years) Mean [SD, range]	29 [5.9, 17–51]	29.2 [4.98, 18–42]
Education Year 12 or less	49 (48.5)	35 (36.1)
Income		
0-\$51,999 per annum	24 (23.8)	29 (29.9)
> \$52,000-\$77,999	35 (34.7)	20 (20.6)
> \$78,000 per annum	41 (40.6)	44 (45.4)
Missing	1 (1)	4 (4.1)
Nulliparous	58 (57.4)	58 (59.8)
No partner	9 (8.9)	2 (2.1)
Foreign born	25 (24.8)	31 (32)
Gestation at recruitment Mean [SD, range]	18.2 [3.17, 11–25]	17.9 [2.8, 13–24]
Preferred birth mode		
Cesarean	15 (14.9)	16 (16.5)
Uncertain	2 (2)	3 (3.1)
Cesarean last birth		
No labor cesarean	5 (5)	4 (4.1)
Labor/Cesarean	10 (9.9)	10 (10.3)
History depression/anxiety	52 (51.5)	42 (43.3)
History tobacco smoking	52 (51.5)	58 (59.8)
W-DEQ Median {IQR}	77 {72,86}	73 {69, 78.5}
W-DEQ Mean [SD, range]	80.9 [13.1, 66–127]	75.7 [9.7, 66–115]
CBSEI Median {IQR}	372 {301, 465.5}	371.5 {302, 464}
CBSEI Mean [SD, range]	368.5 [122.5, 62–620]	385.9 [102.9, 62–591]
Missing	3 (3)	3 (3)
EPDS Median {IQR}	9 {4, 13}	7 {3.5, 10}
EPDS Mean [SD, range]	8.72 [5.8, 0-22]	7.33 [4.7, 0–24]
DCS Median {IQR}	43.7 {25, 56.2}	37.5 {26.6, 50}
DCS Mean [SD, range]	41.3 [24.3, 0–100]	39.1 [19.5, 0–94]
Missing	_	2 (2)

W-DEQ = Wijma Delivery Expectancy Scale; CBSEI = Childbirth Self-Efficacy Inventory; DCS = Decisional Conflict Scale; EPDS = Edinburgh Postnatal Depression Scale; IQR = Interquartile range: 25th percentile, 75th percentile.

A 12-point or more improvement in DCS score was adjusted for, being the minimal score difference between making or delaying decision making. No adjusted changes were made to CBSEI or EPDS scores. A higher percentage of women who received the intervention had higher levels of childbirth confidence (CBSEI) (n = 74/97, 76.3%) compared to controls (n = 55/91, 60.4%), ( $\chi^2(1, n = 188) = 4.8, p = 0.03$ , phi = -3) with moderate effect. Although not significant, reduced levels of decisional conflict were found

for women who received the intervention (n = 66/101, 65.3%) compared to controls (n = 50/95, 52.6%),  $(\chi^2(1, 96) = 2.8, p = 0.09, phi = -.2)$  with small to moderate effect (33). Similarly, a higher percentage of women in the intervention group showed improvement for depression (n = 56/101, 55.4%) compared to controls (n = 47/97, 48.5%); however, this improvement was not statistically significant,  $(\chi^2(1, n = 198) = .71, p = 0.40, phi = 0.15$  (Table 5).

#### Discussion

This brief psycho-education telephone counseling intervention offered by midwives during pregnancy was



Fig. 2. The Consolidated Standards of Reporting Trials (CONSORT) flow diagram.

effective in reducing women's fears, and improved childbirth confidence. Two previous studies of antenatal interventions by midwives to reduce childbirth fear showed reduction in requests for cesarean. Neither study, however, used a randomized controlled trial (RCT) design to evaluate the intervention, nor did they use a standardized measure of fear (18,34).

The closest comparative studies are two RCTs conducted in Finland. Saisto et al randomized fearful nulliparous and multiparous women deemed at low obstetric risk to receive either six counseling sessions or two sessions provided within conventional care (19). The intervention was delivered by an obstetrician trained in cognitive psychotherapy. A significant reduction in women's birth concerns was found in the intervention group, with decreased requests for cesarean in both study groups. A more recent RCT by the same team offered women with severe fear (WDEQ  $\geq 100$ ) six antenatal and one postpartum group sessions with a psychologist (14 hours overall). The intervention reduced fear levels, and improved vaginal birth rates (20) and confidence for women receiving group psycho-education (35). Although 56 percent of women in

Table 3		Internal	Reliability	of	Scales
---------	--	----------	-------------	----	--------

Scale	Cronbach's alpha	Mean [SD, Range]
Wijma Delivery Expectancy/Experience Questionnaire (W-DEQ-A)	0.93	49.5 [21.9, 0–128]
ChildBirth Self-Efficacy Inventory (CBSEI)	0.98	443 [112.2, 62–620]
Edinburgh Postnatal Depression Scale (EPDS)	0.86	5.1 [4.6, 0–24]
Decisional Conflict Scale (DCS)	0.97	29.7 [23, 0–100]

their control group also received specialized support for fear from other services, women allocated to the intervention group fared better.

The interventions by the Finnish researchers had similar intent to the current study to increase women's knowledge and confidence by identifying women's feelings that affect choices for their pending births. However, Saisto et al did not measure fear using the WDEQ-A, and Rouhe et al did not measure fear in the third trimester (19,20). Furthermore, women in control

 
 Table 5. Percentage of Participants in Intervention and Control Groups Who Showed Improvement on Primary and Secondary Outcomes

	Intervention n = 101	$\begin{array}{l} Control\\ n = 97\\ N_{T} \left( \mathcal{O} \right) \end{array}$	
Outcome variables	NO. (%)	NO. (%)	р
Fear (WDEQ-A)			
Improved > 20 score	48 (47.5)	25 (25.8)	0.002
No Improvement	50 (59.5)	71 (73.2)	
Missing	3 (3)	1 (1)	
Childbirth confidence	(CBSEI)		
Improved	74 (73.3)	55 (56.7)	0.03
No improvement	23 (22.8)	36 (37.1)	
Missing	4 (4)	6 (6.2)	
Decisional Conflict (D	DCS)		
Improved > 12 score	66 (65.3)	50 (51.5)	0.09
No Improvement	35 (34.7)	45 (46.4)	
Missing	_	2 (2.1)	
Depression (EPDS)			
Improved	56 (55.4)	47 (48.5)	0.40
No Improvement	45 (44.6)	50 (51.5)	
Missing	_	_	

*W-DEQ* = *Wijma Delivery Expectancy Scale; CBSEI* = *Childbirth Self-Efficacy Inventory; DCS* = *Decisional Conflict Scale; EPDS* = *Edinburgh Postnatal Depression Scale.* 

Table 4.	Change	in 🖁	Scores	for	Intervention a	and (	Control	Groups f	for l	Primary	and	Secondary	Outcomes
----------	--------	------	--------	-----	----------------	-------	---------	----------	-------	---------	-----	-----------	----------

Variables	1	ntervention N = 101		$\begin{array}{l} Control \\ N = 97 \end{array}$		Change in sco	res	
Primary and secondary outcomes	n	Mean change score [SD]	n	Mean change score [SD]	Mean change difference	95% CI for mean change difference	р	Effect size
WDEQ	98	19.52 [18.59]	96	9.28 [16.32]	10.24	5.29-15.19	< 0.001*	0.59 <sup>†</sup>
CBSEI	97	61.10 [87.45]	91	19.70 [92.61]	41.40	15.48-67.30	0.002*	$0.46^{\dagger}$
DCS	101	21.74 [25.69]	95	16.13 [21.24]	5.60	-1.06-12.26	0.09	n/a
EPDS	101	1.26 [4.98]	97	.61 [5.30]	.65	-0.79 - 2.09	0.38	n/a

\*Bonferonni indicates significance at p = 0.0125. Change scores were calculated by subtracting time 2 from time 1 (WDEQ, DCS, EPDS) and for CBSEI subtracting time 1 from time 2. <sup>†</sup>Moderate effect size (32). W-DEQ = Wijma Delivery Expectancy Scale; CBSEI = Childbirth Self-Efficacy Inventory; DCS = Decisional Conflict Scale; EPDS = Edinburgh Postnatal Depression Scale.

groups of both studies had access to other treatment options for fear. In the current Australian maternity system, no specific treatment for childbirth fear is available, and this allows for a clear determination that differences were a result of the midwife intervention and not confounded by women accessing a similar service elsewhere.

#### Childbirth Confidence

Low self-efficacy is associated with childbirth fear, increased perception of pain in labor, and obstetric interventions (36.37). We found that a brief midwife telephone counseling intervention for women with high childbirth fear could significantly improve confidence in women of any parity during pregnancy, and confirmed that childbirth self-efficacy is modifiable. A recent Australian pilot study also found significant improvement in women's (n = 18) childbirth self-efficacy and fear levels following a mindfulness antenatal education program with first-time mothers (38). That program included 20 contact hours with an experienced yoga and childbirth educator and required participants to complete a precourse homework package. An efficacy enhancing antenatal education program conducted in China also reported improved childbirth confidence and lower perception of pain and anxiety during labor (39). Two studies in Taiwan improved childbirth confidence through the use of birth balls in labor (40) and a prenatal yoga program (41). These three studies, however, did not include parous women or women with childbirth fear. Our study is the first RCT to test childbirth self-efficacy within an Australian population of fearful women.

Improving women's belief in their ability to cope with normal physiological and emotional challenges of labor is fundamental to birth preparation. The task is even more challenging in women who are fearful of birth and at higher risk of requesting a cesarean to avoid the experience of labor.

Women in our cohort who had lower incomes and education levels were less likely to continue in the study. Further investigation is required to determine the best methods for engagement and providing support. Women's lack of confidence may have impacted their ability to continue participation, or challenging women's health beliefs may have contributed to attrition and would be best assessed and monitored through a trusted continuity of caregiver relationship.

#### Decisional Conflict

Women receiving the BELIEF intervention reported lower levels of decisional conflict than controls at 36 weeks of pregnancy but the change was not significant. This finding may be an important one. High rates of decisional conflict are associated with regret and ambivalence (42) and may have consequences for women's birth choices and satisfaction. Decisional conflict is commonly measured in studies investigating women's decision making in a subsequent pregnancy after a previous cesarean and has not been applied to childbirth fear (43,44). Our participants with childbirth fear had higher baseline decisional conflict scores compared to women in other studies who were possibly less distressed and making choices about birth mode alone. The reduction in decisional conflict in both the intervention and control groups may indicate not only the benefit of the decision aid for women with childbirth fear but also that psychoeducation with a midwife enhances the effectiveness of the decision aid.

#### **Depression** Symptoms

Poor emotional health is associated with increased childbirth fear (13,15,45). In our study, 21 percent of women reported EPDS scores of greater than 12, which is higher than the 8.1 percent identified in fearful Norwe gian women (WDEQ  $\geq 85$ ) (46) but lower than the 30 percent rate of probable depression in fearful Swedish women (47). Our telephone counseling intervention reduced women's EPDS scores compared to baseline, but not significantly. Positive effects may be attributed to listening and facilitating women's concerns, where their worries were addressed and alleviated. The reduction in depressive symptoms following the intervention in the current study aligns with previous advice to explore women's unhappiness in the preparation for birth (48). This finding also gives support for midwifery continuity of care models where psychosocial issues are more readily addressed than within biomedical models of care (11,49,50).

#### Limitations

Recruitment occurred within the public health system; however, one-third of all women in Australia receive private health care. At baseline, demographics were similar to the national birthing population, but of those randomized, women who were less educated and poorer were less likely to continue in the study. Women requiring an interpreter were also not included; therefore, caution is needed in drawing generalizations. Although there may be differences in childbirth fear by parity, the sample size was not sufficient to support subgroup analysis.

However, the robustness of findings is strengthened because of the RCT design with participants stratified by site and parity. The study was the first to include a measure of decisional conflict and therefore no comparison to other childbirth fear studies could be made. Investigating decisional conflict in this group of vulnerable women can shed light on how and why they make health decisions, and increase awareness about how informed consent for procedures might be determined. Although conducting the intervention over the telephone offered a great deal of flexibility and accessibility for participants, there were also the disadvantages. A small proportion of women were multitasking (for example doing dishes, washing clothes) when receiving the psycho-education. Outcomes may have been further improved had all women been able to find a quiet space for these conversations.

Few women scored extremely high on the W-DEQ ( $\geq 85$ , n = 42), indicating severe fear with probable clinical symptoms (51). We cannot know whether the BELIEF intervention would be enough in these cases. On the other hand, perhaps reducing moderate fear in one pregnancy may prevent a development of a more serious childbirth anxiety or phobia later, in the same or in the next pregnancy. Furthermore, a proportion of women did not proceed with the intervention. Engagement of fearful women may have been enhanced by using other modes such as the Internet, which would allow 24-hour access to educational materials and chat options for questions and support.

#### Conclusions

This was the first reported RCT of an antenatal intervention for childbirth fear by midwives. The BELIEF psycho-education intervention facilitated a communicative and caring partnership with women to reduce or heal childbirth fear from mid to late pregnancy. Our results indicate fear can be modified. Assisting women to reframe their perceptions about their ability to birth is a critical strategy in preserving the normality of birth. Provision of educational and emotional support has the potential to reduce interventions such as cesarean delivery and the emotional or psychological consequences of what could be otherwise experienced as a disappointing or traumatic birth (52). Assisting women to achieve a normal birth will improve women's quality of reproductive life, reduce health care costs, and improve postpartum maternal and child health outcomes (21,28).

Asking women explicitly about their fears and concerns significantly lowered fear and improved childbirth confidence compared to women receiving standard care. Consideration needs to be given to routine screening around childbirth fear, and application of existing BIRTH 41:4 December 2014

best evidence for promoting normal birth. There was a tendency toward improved women's decision making with the provision of a decision aid and access to individualized midwifery telephone support; however, the difference was not statistically significant. This tendency has important implications for how informed consent is secured in clinical practice and within the context of a maternity culture of high obstetric interventions where women's ability for ready decision making is shown to be vulnerable. Furthermore, the intervention provided a midwife confidante with whom women could share and discuss their concerns, and have their concerns heard. This level of information and support may be particularly important where women are not offered continuity of care or where social support is lacking.

The BELIEF telephone intervention is brief (particularly in comparison to other reported interventions), reproducible (given it has been adapted from use with postnatal women) (28), effective, and could be easily introduced into routine midwifery practice with women reporting high fear. BELIEF involves listening and responding to women's feelings (a factor previously identified as important) and provides consistent and accurate information. The current intervention is adaptable for individual sessions or group work, and could be delivered in person or using other media. Given the geographical remoteness of some health services in Australia and other countries, flexibility in delivery is important. Where trusting relationships are built, women are more likely to stay engaged in care.

#### Acknowledgments

The BELIEF study was funded by NHMRC (grant ID APP1025099). The decision-aid booklet was developed by the Queensland Centre for Mothers & Babies, The University of Queensland. Associate Professor Julie Pallant provided statistical advice.

#### References

- Salomonsson B, Bertero C, Alehagen S. Self-efficacy in pregnant women with severe fear of childbirth. J Obstet Gynecol Neonatal Nurs 2013;42(2):191–202.
- Zar M, Wijma K, Wijma B. Pre- and postpartum fear of childbirth in nulliparous and parous women. *Scandinavian J Behav Therapy* 2001;30(2):75–84.
- Adams S, Eberhard-Gran M, Eskild A. Fear of childbirth and duration of labour: A study of 2206 women with intended vaginal delivery. *BJOG* 2012;119:1238–46. doi:10.1111/j1471-0528201203433x.
- Kjaergaard H, Wijma K, Dykes A, Alehagen S. Fear of childbirth in obstetrically low-risk nulliparous women in Sweden & Denmark. J Reprod Infant Psyc 2008;26(4):340–350.

- Rouhe H, Salmela-Aro K, Halmesmaki E, Saisto T. Fear of childbirth according to parity, gestational age, and obstetric history. *BJOG* 2009;116(1):67–73.
- Fenwick J, Gamble J, Nathan E, et al. Pre- and postpartum levels of childbirth fear and the relationship to birth outcomes in a cohort of Australian women. J Clin Nurs 2009;18:667–677.
- Haines H, Pallant J, Karlstrom A, Hildingsson I. Cross-cultural comparison of levels of childbirth-related fear in an Australian and Swedish sample. *Midwifery* 2011;27(4):560–567.
- Toohill J, Fenwick J, Gamble J, Creedy DK. Prevalence of childbirth fear in an Australian sample of pregnant women. *BMC Pregnancy Childbirth*. 2014;14: 275. DOI: 10.1186/1471-2393-14-275.
- Johnson R, Slade P. Does fear of childbirth during pregnancy predict emergency caesarean section? *BJOG* 2002;109(11):1213– 1221.
- Ryding E, Wijma B, Wijma K, Rydhstrom H. Fear of childbirth during pregnancy may increase the risk of emergency cesarean section. *Acta Obstet Gynecol Scand* 1998;77:542–547.
- Cumberland S. Cause for concern: Maternal fear around childbirth and caesarean section as a mode of delivery. *MIDIRS* 2010;20(3):297–304.
- Saisto T, Halmesmaki E. Fear of childbirth: A neglected dilemma. Acta Obstet Gynecol Scand 2003;82:201–208.
- Laursen M, Hedegaard M, Johansen C. Fear of childbirth: Predictors and temporal changes among nulliparous women in the Danish National Birth Cohort. *BJOG* 2008;115(3):354–360.
- Rouhe H, Salmela-Aro K, Gissler M, et al. Mental health problems common in women with fear of childbirth. *BJOG* 2011;118 (9):1104–1111.
- Storksen H, Eberhard-Gran M, Garthus-Niegel S, Eskild A. Fear of childbirth; the relation to anxiety and depression. *Acta Obstet Gynecol Scand* 2012;91:237–242.
- Bewley S, Cockburn J. Responding to fear of childbirth. Lancet 2002;359(9324):2128–2129.
- Otley H. Fear of childbirth: Understanding the causes, impact and treatment. Br J Midwifery 2011;19(4):215–220.
- Halvorsen L, Nerum H, Sorlie T, Oian P. Does counsellor's attitude influence change in a request for a caesarean in women with fear of birth? *Midwifery* 2010;26:45–52.
- Saisto T, Salmela-Aro K, Nurmi J, et al. A randomized controlled trial of intervention in fear of childbirth. *Obstet Gynecol* 2001;98(5):820–826.
- Rouhe H, Salmela-Aro K, Toivanen R, et al. Obstetric outcome after intervention for severe fear of childbirth in nulliparous women – randomised trial. *BJOG* 2013;120(1):75–84.
- Fenwick J, Gamble J, Creedy D, et al. Study protocol for reducing childbirth fear: A midwife-led psycho-education intervention. *BMC Pregnancy and Childbirth* [Internet]. 2013;13(190):[Published online October 20th, 2013 pp.]. Available at: http://www. biomedcentral.com/1471-2393/13/190.
- Wijma K, Wijma B, Zar M. Psychometric aspects of the W-DEQ; a new questionnaire for the measurement of fear of childbirth. J Psychosom Obstet Gynecol 1998;19(2):84–97.
- Institute for Statistics and Mathematics. The R Project for Statistical Computing 2012 Accessed January 19, 2012. Available at: http://www.r-project.org/.
- 24. Lowe N. Maternal confidence for labor: Development of the childbirth self-efficacy inventory. *Res Nurs Health* 1993;16(2): 141–149.
- Cox J, Holden J, Sagousky R. Detection of postnatal depression development of the 10 item Edinburgh Depression Scale. *BMJ* 1987;150:782–786.
- Downe S, Butler E, Hinder S. Screening tools for depressed mood after childbirth in UK-based South Asian women: A systematic review. J Adv Nurs 2007;57:565–583.

- O'Connor A. User Manual Decisional Conflict Scale (16 item statement format). Ottawa, Ontario, Canada: Ottawa Hospital Research Institute; 1993, updated 2010. Available at: http://decisionaid.ohri.ca/docs/develop/User\_Manuals/UM\_Decisional\_Conflict.pdf.
- Gamble J, Creedy D. A counselling model for postpartum women after distressing birth experiences. *Midwifery* 2009;25: e21–e30.
- 29. Queensland Centre for Mothers and Babies. The Having a Baby in Queensland Book - Your choices during pregnancy and birth. Accessed November 6, 2011. Available at: http://www.qcmb.org. au/the\_having\_a\_baby\_in\_queensland\_book\_project.
- IBM Corp. IBM SPSS Statistics for Windows, Version 21.0. Armonk, NY: IBM Corp. 2012.
- Kazis J, Anderson J, Meenan R. Effect sizes for interpreting changes in health status. *Med Care* 1989;27:s178–s188.
- 32. Streiner D, Norman G *Biostatistics: The Bare Essentials* (2nd ed). Hamilton, Ontario, Canada: B.C. Decker Inc., 2000.
- Cohen J. Statistical Power Analysis for the Behavioural Sciences, Vol. 2nd. Hillsdale, NJ: Erlbaum Associates, 1988.
- Nerum H, Halvorsen L, Sorlie T, Oian P. Maternal request for cesarean section due to fear of birth: Can It be changed through crisis-oriented counseling? *Birth* 2006;33 (3):221–228.
- Salmela-Aro K, Read S, Rouhe H, et al. Promoting positive motherhood among nulliparous pregnant women with an intense fear of childbirth: RCT intervention. *J Health Psychol* 2012;17 (4):520–534.
- Drummond J, Rickwood D. Childbirth confidence: Validating the childbirth self-efficacy inventory (CBSEI) in an Australian sample. J Adv Nurs 1997;26(3):613–622.
- Berentson-Shaw J, Scott K, Jose P. Do self-efficacy beliefs predict the primiparous labour and birth experience? A longitudinal study. J Reprod Infant Psychol 2009;27(4):357–373.
- Byrne J, Hauck Y, Fisher C, et al. Effectiveness of a Mindfulness-Based Childbirth Education pilot study on maternal self-efficacy and fear of childbirth. J Midwifery Women's Health 2014; 59(2):192–197.
- Ip W, Tang C, Goggins W. An educational intervention to improve women's ability to cope with childbirth. J Clin Nurs 2009;18:2125–2135.
- Gau M, Chang C, Tian S, Lin K. Effects of birth ball exercise on pain and self-efficacy during childbirth: A randomised controlled trial in Taiwan. *Midwifery* 2011;27:e293–e300.
- Sun Y, Hung Y, Chang Y, Kuo S. Effects of a prenatal yoga programme on the discomforts of pregnancy and maternal childbirth self-efficacy in Taiwan. *Midwifery* 2010;26(6):e31– e36.
- Gattelari M, Ward J. Decisional conflict forgoing PSA testing for prostate cancer. J Med Screen 2004;11:165–9.
- Shorten A, Shorten B, Keogh J, et al. Making choices for childbirth: A randomized controlled trial of a decision-aid for informed birth after cesarean. *Birth*: Issues in Perinatal Care 2005;32(4):252–261.
- Montgomery A, Emmett C, Fahey T, et al. Two decision aids for mode of delivery among women with previous caesarean section: Randomised controlled trial. *BMJ* 2007;334:1305. doi:10. 1136/bmj.39217.671019.55.
- Toohill J, Fenwick J, Gamble J, et al. Psycho-social predictors of childbirth fear in pregnant women: An Australian study. *Open J Obstet Gynecol.* 2014;4(9):531–543. doi.org/10.4236/ojog.2014. 49075
- Nordeng H, Hansen C, Garthus-Niegel S, Eberhard-Gran M. Fear of childbirth, mental health, and medication use during pregnancy. Arch Womens Ment Health 2012;15:203–209.

- 47. Waldenstrom U, Hildingsson I, Ryding E. Antenatal fear of childbirth and its association with subsequent caesarean section and experience of childbirth. *BJOG* 2006;113(6):638–646.
- Barclay L, Lloyd B. The misery of motherhood: Alternative approaches to maternal distress. *Midwifery* 1996;12(3):136–139.
- Lyberg A, Severinsson E. Midwives' supervisory styles and leadership role as experienced by Norwegian mothers in the context of a fear of childbirth. *J Nurs Manag* 2010;18(4):391–399.
- Kirkham M. Emotion work around reproduction: Supportive or constraining. In: Hunter B, Deery R, eds. *Emotions in Midwifery* and Reproduction. New York: Palgrave Macmillan, 2009.
- Nieminen K, Stephansson O, Ryding EL. Women's fear of childbirth and preference for cesarean section – a cross-sectional study at various stages of pregnancy in Sweden. *Acta Obstet Gynecol Scand* 2009;88(7):807–813.
- Garthus-Niegel S, Knoph C, von Soest T, et al. The role of labor pain and overall birth experience in the development of posttraumatic stress symptoms: A longitudinal cohort study. *Birth* 2014;41(1):108–115.
- 53. Murray D, Cox J. Screening for depression during pregnancy with the Edinburgh Depression Scale (EPDS). Special Issue: Psychiatric disorders associated with childbearing. *J Reprod Infant Psychol.* 1990;8:99–107.
- 54. Beyondblue. Clinical practice guidelines for depression and related disorders – anxiety, bipolar disorder and puerperal psychosis – in the perinatal period. A guideline for primary care health professionals. Melbourne, Australia: BeyondBlue: The National Depression Initiative, 2011. Accessed January 7, 2014. Available at: https://www.bspg.com.au/dam/bsg/product?client=BEYONDBLUE &prodid=BL/0891&type=file.