

Red, Blue or Mix: Choice of Optimal Light Qualities for Enhanced Plant Growth and Development through *in silico* Analysis

Andrew M.H. Chan,¹ Miao Lin Pay,² Jesper Christensen,² Fei He,³ Laura C. Roden,⁴ Hafiz Ahmed⁵ and Mathias Foo^{1,*}

¹School of Engineering, University of Warwick, CV4 7AL, Coventry, UK, ²Centre for Future Transports and Cities, Coventry University, CV1 5FB, Coventry, UK, ³Centre for Computational Science and Mathematical Modelling, Coventry University, CV1 5FB, Coventry, UK, ⁴Centre for Health and Life Sciences, Faculty of Health and Life Sciences, School of Life Sciences, Coventry University, CV1 5FB, Coventry, UK and ⁵Nuclear Advanced Manufacturing Research Centre, The University of Sheffield, S60 5WG, Derby, UK

*Corresponding author. M.Foo@warwick.ac.uk

Accepted Manuscript

Abstract

In smart greenhouse farming, the impact of light qualities on plant growth and development is crucial but lacks systematic identification of optimal combinations. This study addresses this gap by analysing various light properties' effects (photoperiod, intensity, ratio, light-dark order) on *Arabidopsis thaliana* growth using days-to-flower (DTF) and hypocotyl length as proxies to measure plant growth and development. After establishing suitable ranges through comprehensive literature review, these properties were varied within those ranges. Compared to white light, a 16-hour cycle of blue light reduces DTF and hypocotyl length by 12% and 3%, respectively. Interestingly, similar results can be achieved using a shorter photoperiod of 14-hour light (composed of 8 hours of a mixture of $66.7 \mu\text{mol}/\text{m}^2\text{s}^{-1}$ red and $800 \mu\text{mol}/\text{m}^2\text{s}^{-1}$ blue lights (i.e., blue:red ratio of 12:1) followed by 6 hours of monochromatic red light and 10-hour dark. These findings offer potential for efficient growth light recipes in smart greenhouse farming, optimising productivity while minimising energy consumption.

Key words: light quality, light order, photomorphogenesis, flowering time, hypocotyl elongation, plant growth and development, smart greenhouse farming

Accepted Manuscript

Introduction

In 2017, The Food and Agriculture Organization of the United Nations published a report, stating that by the year 2100, global population is expected to grow to around 11 billion of which most population growth is expected to be in Asia and Africa (United Nation, 2017). This rapid growth inevitably brings forward rapid urbanisation, leading to a reduction in agricultural farmland if no further deforestation is considered. The same report also stated that the population demographic is also likely to get much older and this would hit farming in rural areas especially hard, as the younger workforce migrates to cities, thereby reducing the farming workforce. The reduction in the workforce coupled with the ongoing climate changes that bring in more floods and drought would make it even more challenging to meet the growing food demand.

Controlled environment agriculture - a technology-oriented approach that provide crop with consistent optimal growing condition and shielding them from external elements – such as smart greenhouse farming, has been identified as one of the viable solutions to achieve food security (Walter et al., 2017; Mohamed et al., 2021; O'Shaughnessy et al., 2021; Karanisa et al., 2022). By artificially controlling plant exposure to important growth variables, such as light and water, smart greenhouse farming has the potential to be more sustainable than conventional farming practices with the “more food with less” concept through better efficiency in managing the use of those variables (Aune et al., 2017; Kumar et al., 2019).

One of the important variables that notably influence plant development in smart greenhouse farming is light (Kami et al., 2010; Neo et al., 2022). Thus, it is not surprising to see many studies are dedicated in finding the most efficient lighting strategy for smart greenhouse farming usage either through experimentation (Olvera-Gonzalez et al., 2013) or optimisation theory (Mosharafian et al., 2021).

The speed breeding protocol pioneered in (Watson et al., 2018) represents the pinnacle of the role of manipulating light in enhancing crop yield. Through experimentation, this approach has demonstrated that the growth of crops such as wheat and barley can be accelerated by at least two times when exposed to extended periods of light - up to 22 hours. More importantly, this protocol can be easily adapted for use in growth chambers and greenhouses (Ghosh et al., 2018), making it an appealing option for incorporating effective lighting management into intelligent greenhouse agriculture to boost crop productivity.

To establish a framework of artificial light management capable for speed breeding protocol in smart greenhouse farming, six different light properties of white light were analysed in a systematic manner to determine which property is the most influential on plant growth (Pereira et al., 2021). The study confirmed the conclusions of literature reviews articles by (Bian et al., 2015) and (Ahmed et al., 2020). They concluded photoperiod and intensity are the most influential properties and suggested that similar DTF and hypocotyl length can be achieved with light exposure of up to 18 hours as opposed to using photoperiod of 22 hours following the speed breeding protocol. Interestingly, the review by (Bian et al., 2015) and (Ahmed et al., 2020) indicated that light quality (colour) is a key property for plant development, which is also highlighted by (Pereira et al., 2021) as a future research work.

In this study, following a similar approach to (Pereira et al., 2021), we extend the use of systematic approach to investigate the effect of light qualities and its associated properties on plant development as a possible extension to the speed breeding protocol. As detailed in the next section, there are numerous experimental studies with different suggestions on the preferred light qualities for plant growth and development. To reconcile this discordance, we explore the use of *in silico* analysis to provide a more representative quantitative range of the light qualities. Another advantage of *in silico* experiments is the provision of benchmarking using one or multiple plant species that presents a viable alternative to the conventional experimental approaches in determining the best conditions for plant growth and development.

The main contributions of this study are as follows: a comprehensive literature search compiling the range of light quality relevant for plant growth and development across various types of plants followed by performing a systematic analysis on the effect of different blue and red light properties, i.e., intensity, ratio, photoperiod, and wavelength of light order on plant growth and development. For the light order, a total of 28 combinations of light-dark order are considered, which to our best knowledge is the first time being analysed in an *in silico* setting. Such exhaustive light combinations are considered to allow us to find the suitable light recipe that optimises plant growth and development and eventually leads to enhanced yield and energy consumption, thereby further improving the sustainability of smart greenhouse farming.

Methods

Identifying experimental lighting conditions through literature review

There are many experimental studies available in the literature that investigate the effect of different light qualities on different plants. To enable us to obtain a practical range of these different light properties for our analysis, a comprehensive literature search is carried out to identify all those possible ranges that have been considered. Specifically, in our literature search, we focused on studies that use either blue or red lights to investigate plant growth or developmental characteristic, and we narrowed this down to 34 publications that are relevant to our study. Supplementary Table 1 summarises our findings from the 34 publications covering 14 different genera of plants used in experimental studies encompassing agriculturally important plants, as well as the model plant *Arabidopsis thaliana*. All these plants respond differently to the effect of light qualities and any agricultural benefits that can be acquired from these light qualities depend on whether the crop favours vegetative growth (development of roots and shoots) or reproductive growth (flowering and fruit or seed formation). For instance, fruit- and seed-based crops may benefit, if the flowering were accelerated; vegetative crops benefit from light regimes which delay flowering.

Considering the varieties of plant genera, this naturally leads to the question of the relevance of the light experiments from other plant genera to *in silico* light analysis using a well-established *Arabidopsis thaliana* mathematical model that relates input light to plant growth and development related pathways (Seaton et al., 2015; De Caluwé et al., 2016; Chew et al., 2022). In (Song et al., 2010), the authors found that the circadian gene expression of *Arabidopsis thaliana* is shared in other eudicots in the plant kingdom, such as in legumes, rice, duckweed, moss, alga and many more long-day plants within and even outside of the angiosperms (flowering producing and fruit seed bearing plants) clade. Moreover, the study in (McClung, 2013) concludes that in angiosperms many of the circadian clock characteristics and functions of *Arabidopsis thaliana* are retained. These two studies provide evidence that the circadian clock components that regulate growth and other characteristics of *Arabidopsis thaliana* are shared by a plethora of other plants. Given this rich knowledge we have on *Arabidopsis thaliana*,

we aim to understand plant growth behaviour in relation to various light qualities through mathematical modelling and to use these findings to provide inferences on the effect of light qualities on the growth behaviour of other plants.

It has been demonstrated in *Arabidopsis thaliana* that the spectral quality of light is more important in regulating flowering time (Eskins, 1992; Guo et al., 1998) and hypocotyl length (Spaninks et al., 2020) under different photoperiods, i.e., 14 hours (Eskins, 1992), 16 hours (Spaninks et al., 2020) and 18 hours Guo et al. (1998) and light intensities, i.e., 25 to 164 $\mu\text{mol}/\text{m}^2\text{s}^{-1}$ (Eskins, 1992), $120 \pm 10 \mu\text{mol}/\text{m}^2\text{s}^{-1}$ (Spaninks et al., 2020), and 75 to 85 $\mu\text{mol}/\text{m}^2\text{s}^{-1}$ (Guo et al., 1998), with blue light being most effective at promoting flowering. Conversely, red light, with the red: blue ratio ranging from 2:1 to 10:1, has been demonstrated to promote vegetative growth, important in leafy plants such as lettuce, cabbage, spinach, and basil (Hoenecke et al., 1992; Matsuda et al., 2007; Stutte et al., 2009; Lee et al., 2010; Vaštakaitė et al., 2015; Zhang et al., 2018; Zou et al., 2020). However, studies that have considered a mixture of red and blue on a 1:1 ratio, or a pure blue instead, have demonstrated greater biomass accumulation than with red light alone (Muneer et al., 2014; Kwack et al., 2015; Piovene et al., 2015). It is thus key to optimise spectral quality to balance vegetative and reproductive growth dependent on the requirements of the crop plant, while at the same time promoting carbon fixation and plant growth.

Literature on fruit-, root- and shoot-based plants, also have similar arguments, suggesting that the large variation from drawn conclusions is not unique to literature investigating leafy plants alone, with some studies concluding that red alone or with relatively small blue light as the supplement is the optimal for plant growth (Brown et al., 1995; Hogewoning et al., 2007; Folta and Childers, 2008; Gangadhar et al., 2012; Hernández and Kubota, 2014; Paradiso et al., 2019) while others conclude that blue light alone if not blue light with red light as the supplement is better (Wang et al., 2009; Yoshida et al., 2012; Cope and Bugbee, 2013; Kwack et al., 2015; Piovene et al., 2015). All these mentioned studies do not use the same intensity nor share a similar intensity range in their respective studies. The Photosynthetic Photon Flux Density (PPFD) values utilised in those studies range from around 50 to 300 $\mu\text{mol}/\text{m}^2\text{s}^{-1}$.

On the effect of PPFD on plants, a study in (Hidaka et al., 2013) concludes that lighting with a PPFD of 400 to 1200 $\mu\text{mol}/\text{m}^2\text{s}^{-1}$ is far more effective than with PPFD ranging from 0 to 200 $\mu\text{mol}/\text{m}^2\text{s}^{-1}$ for strawberry. In another study in (Fu et al., 2012), the authors discover that a range of PPFD 400 to 600 $\mu\text{mol}/\text{m}^2\text{s}^{-1}$ is the optimal intensity for lettuce depending on the altitude where the lettuce is from and grown. Both studies also discovered a plateau in the photosynthetic rate and suggest the reason for a lack of increase in crop yield even when light intensity is increased limitlessly.

One key takeaway point from this literature review is that a cohesive conclusion on the effect of light qualities is difficult to be established even when examining the plant from the same family with the same mutation and the same measuring indices. We thus wish to investigate the effect of all the ranges of light properties: spectrum, photoperiod, and intensity, to establish a more cohesive conclusion on their effects on plant growth and development in a systematic manner using *Arabidopsis thaliana* as a model. More importantly, it provides us the opportunity to fill any gaps on different combination of light properties that have not been previously considered.

Arabidopsis thaliana Simulation Model

To investigate the optimal lighting strategy, the simulation model used in this study is from our previous works (Pay et al., 2022b,a). It comprises 27 Ordinary Differential Equations (ODEs), where the input is the light, and the outputs are the growth-related circadian genes that are used to compute days-to-flower (DTF) and hypocotyl length as shown in Figure 1.

Fig. 1. Overview of the interactions in *Arabidopsis thaliana* mathematical model relating input light to output hypocotyl and flowering pathways.

These 27 ODEs describe the expressions of genes and proteins that form the plant circadian system and two downstream pathways related to flowering and hypocotyl elongation. These ODEs are solved using MATLAB function `ode15s`. As a note, the notations used for all the ODEs follows (Pay et al., 2022b,a) and their respective parameter values are found in Supplementary Table 2.

Core Plant Circadian System

Eight ODEs form the core plant circadian system, which describes the dynamics of the CIRCADIAN CLOCK ASSO-CIATED 1/LATE ELONGATED HYPOCOTYL, (CL) gene and protein, the PSEUDO RESPONSE REGULATOR 9/7, (P97) gene and protein, EARLY FLOWERING 4/LUX ARRHYTHMO, (EL) gene and protein and PSEUDO RESPONSE REGULATOR 5/TIMING OF CAB EXPRESSION 1, (P51) gene and protein.

$$\begin{aligned}\frac{d[CL]_m}{dt} &= (v_1 + L_a) \left(\frac{1}{1 + \left(\frac{[P97]_p}{K_1}\right)^2 + \left(\frac{[P51]_p}{K_2}\right)^2} + \right) \\ &= (k_{1L}\Theta_{PhyA} + k_{1D}(1 - \Theta_{PhyA}))[CL]_m \\ \frac{d[CL]_p}{dt} &= (p_1 + p_{1L}\Theta_{PhyA})[CL]_m - d_1[CL]_p \\ \frac{d[P97]_m}{dt} &= (v_2 + L_b) \left(\frac{1}{1 + \left(\frac{[CL]_p}{K_3}\right)^2 + \left(\frac{[P51]_p}{K_4}\right)^2 + \left(\frac{[EL]_p}{K_5}\right)^2} \right) \\ &\quad - k_2[P97]_m \\ \frac{d[P97]_p}{dt} &= p_2[P97]_m - (d_{2D}(1 - \Theta_{PhyA}) + d_{2L}\Theta_{PhyA})[P97]_p\end{aligned}$$

$$\begin{aligned}
\frac{d[P51]_m}{dt} &= \frac{v_3}{1 + \left(\frac{[CL]_p}{K_6}\right)^2 + \left(\frac{[P51]_p}{K_7}\right)^2} - k_3[P51]_m \\
\frac{d[P51]_p}{dt} &= p_3[P51]_m - (d_{3D}(1 - \Theta_{PhyA}) + d_{3L}\Theta_{PhyA})[P51]_p \\
\frac{d[EL]_m}{dt} &= \frac{\Theta_{PhyA}v_4}{1 + \left(\frac{[CL]_p}{K_6}\right)^2 + \left(\frac{[P51]_p}{K_9}\right)^2 + \left(\frac{[EL]_p}{K_5}\right)^2} - k_4[EL]_m \\
\frac{d[EL]_p}{dt} &= p_4 - d_{e1}[EL]_p \\
&\quad - \left(\frac{d_{e2}[COP1] + d_{e3}[COP1 : PhyA]}{C_{tot}}\right) \\
&\quad - \left(\frac{d_{e4}[COP1 : PhyB] + d_{e5}[COP1 : Cry1]}{C_{tot}}\right)[EL]_p
\end{aligned} \tag{1}$$

where $C_{tot} = [COP1] + [COP1 : PhyA] + [COP1 : PhyB] + [COP1 : Cry1]$ is the total concentration of COP1.

Photoreceptors

Three ODEs are used to describe the three photoreceptors namely PHYTOCHROME A (PhyA), PHYTOCHROME B (PhyB) and CRYPTOCHROME (Cry1), which respond to red and blue lights.

$$\begin{aligned}
\frac{d[PhyA]}{dt} &= (1 - \Theta_{PhyA})A_{p3} - \frac{A_{m7}[PhyA]}{A_{k7} + [PhyA]} \\
&\quad - q_2\Theta_{PhyA}[PhyA] + k_d[COP1 : PhyA] \\
&\quad - k_{mpac}\Theta_{PhyA}[PhyA][COP1] \\
\frac{d[PhyB]}{dt} &= B_{p4} - \frac{B_{m8}[PhyB]}{B_{k8} + [PhyB]} \\
&\quad - k_{mpbc}\Theta_{PhyB}[PhyB][COP1] \\
&\quad + k_d[COP1 : PhyB] \\
\frac{d[Cry1]}{dt} &= C_{p5} - \frac{C_{m9}[Cry1]}{C_{k9} + [Cry1]} \\
&\quad - k_{mpcc}\Theta_{Cry1}[Cry1][COP1] \\
&\quad + k_d[COP1 : Cry1]
\end{aligned} \tag{2}$$

COP1 Interactions

The interaction of these photoreceptors with CONSTITUTIVE PHOTOMORPHOGENIC 1 (COP1) protein, which acts as a key regulator of photomorphogenesis (biological response to light) are described using four ODEs.

$$\begin{aligned}
\frac{d[COP1]}{dt} &= k_d[COP1 : PhyA] + k_d[COP1 : PhyB] \\
&\quad + k_d[COP1 : Cry1] + \frac{A_{m7}[COP1 : PhyA]}{A_{k7} + [COP1 : PhyA]} \\
&\quad + \frac{B_{m8}[COP1 : PhyB]}{B_{k8} + [COP1 : PhyB]} + \frac{C_{m9}[COP1 : Cry1]}{C_{k9} + [COP1 : Cry1]} \\
&\quad + q_2\Theta_{PhyA}[COP1 : PhyA] \\
&\quad - k_{mpac}\Theta_{PhyA}[PhyA][COP1] \\
&\quad - k_{mpbc}\Theta_{PhyB}[PhyB][COP1] \\
&\quad - k_{mpcc}\Theta_{Cry1}[Cry1][COP1]
\end{aligned}$$

$$\begin{aligned}
\frac{d[COP1 : Phya]}{dt} &= k_{mpac}\Theta_{Phya}[Phya][COP1] \\
&\quad - k_d[COP1 : Phya] \\
&\quad - \frac{A_{m7}[COP1 : Phya]}{A_{k7} + [COP1 : Phya]} \\
&\quad - q_2\Theta_{Phya}[COP1 : Phya] \\
\frac{d[COP1 : Phyb]}{dt} &= k_{mpbc}\Theta_{Phyb}[Phyb][COP1] \\
&\quad - k_d[COP1 : Phyb] \\
&\quad - \frac{B_{m8}[COP1 : Phyb]}{B_{k8} + [COP1 : Phyb]} \\
\frac{d[COP1 : Cry1]}{dt} &= k_{mpcc}\Theta_{Cry1}[Cry1][COP1] \\
&\quad - k_d[COP : Cry1] \\
&\quad - \frac{C_{m9}[COP1 : Cry1]}{C_{k9} + [COP1 : Cry1]}
\end{aligned} \tag{3}$$

Hypocotyl Pathway

Three ODEs are used to describe the pathway related to hypocotyl elongation. They are PHYTOCHROME INTER-ACTING FACTOR (PIF) genes and protein and HYPOCOTYL (HYP) that is used to calculate the hypocotyl length.

$$\begin{aligned}
\frac{d[PIF]_m}{dt} &= \frac{v_5}{1 + \left(\frac{[EL]_p}{K_{11}}\right)^2} - k_5[PIF]_m \\
\frac{d[PIF]_p}{dt} &= p_5[PIF]_m - d_{5D}(1 - \Theta_{Phya})[PIF]_p \\
&\quad + d_{5L}\Theta_{Phya}[PIF]_p \\
\frac{d[HYP]_p}{dt} &= g_1 + \frac{g_2[PIF]_p^2}{K_{12}^2 + [PIF]_p^2}
\end{aligned} \tag{4}$$

Flowering Pathway

The remaining nine ODEs are used to describe the genes and proteins in the flowering related pathway. They are GIGANTEA (GI) gene and protein, CYCLING DOF FACTOR 1 (CDF1) gene and protein, FLAVIN-BINDING, KELCH REPEAT, F-BOX 1 (FKF1) gene and protein, CONSTANS (CO) gene and protein and FLOWERING LOCUS T (FT) gene.

$$\begin{aligned}
\frac{d[GI]_m}{dt} &= \frac{N_{v1}}{(NK_1 + [CL]_p^2)(NK_2^2 + [P51]_p^2)(NK_3^2 + [EL]_p^2)} \\
&\quad + L_a - Nk_1[GI]_m \\
\frac{d[GI]_p}{dt} &= Np_1[GI]_m - Nd_1[C_{tot}][EL]_p[GI]_p - Nd_2[GI]_p \\
\frac{d[CDF1]_m}{dt} &= \left(Nv_2 + Nv_3 \frac{[CL]_p^2}{NK_4^2 + [CL]_p^2} \right) \\
&\quad \times \left(\frac{NK_5^2}{NK_5^2 + ([P97]_p + [P51]_p)^2} \right) - Nk_2[CDF1]_m \\
\frac{d[CDF1]_p}{dt} &= Np_2[CDF1]_m - Nd_3Np_3[GI]_p[FKF1]_p[CDF1]_p \\
&\quad - Nd_3Np_4[GI]_p[CDF1]_p - Nd_3[CDF1]_p
\end{aligned}$$

$$\begin{aligned}
\frac{d[FKF1]_m}{dt} &= Nv_4 \left(\frac{NK_6^2}{NK_6^2 + [CL]_p^2} \right) \left(\frac{NK_7}{NK_7 + [EL]_p} \right) \\
&\quad + L_b - NK_3[FKF1]_m \\
\frac{d[FKF1]_p}{dt} &= Np_5[FKF1]_m \\
&\quad - Np_6 \left(Nd_4 - \Theta_{PhyA} \left(\frac{[GI]_p}{Ng_1 + [GI]_p} \right) \right) [FKF1]_p \\
\frac{d[CO]_m}{dt} &= B_{CO} + \left(\frac{NK_8^2}{NK_8^2 + [CDF1]_p^2} \right) \\
&\quad \times \left(Nv_5 + Nv_6(1 - \Theta_{PhyA}) \frac{[COP1]}{NK_9^2 + [COP1]} \right) \\
&\quad - Nk_4[CO]_m \\
\frac{d[CO]_p}{dt} &= Np_7[CO]_m \\
&\quad - Np_8(Nd_5 + Nd_6(1 - \Theta_{PhyA})[COP1])[CO]_p \\
&\quad + Np_8 \left(\Theta_{PhyA} \frac{[FKF1]_p}{Ng_2 + [FKF1]_p} \right) [CO]_p \\
\frac{d[FT]_m}{dt} &= \left(Nv_7 + Nv_8 \frac{[PIF]_p}{NK_{10} + [PIF]_p} \right) \\
&\quad \times \left(Nv_9 + Nv_{10} \frac{NK_{11}}{NK_{11} + [CDF1]_p} \right) \\
&\quad \times \left(\frac{[CO]_p^2}{NK_{12}^2 + [CO]_p^2} \right) - Nk_5[FT]_m
\end{aligned} \tag{5}$$

The equation of light input is given by,

$$\begin{aligned}
L_u &= q_{1u}[T_{PhyA}]\Theta_{PhyA} \\
&\quad + q_{3u}[T_{PhyB}]\log(\eta_1 I_{red} + 1)\Theta_{PhyB} \\
&\quad + q_{4u}[T_{Cry1}]\log(\eta_2 I_{blue} + 1)\Theta_{Cry1}
\end{aligned} \tag{6}$$

where $u \in a$ or b is the effect of light through PhyA or PhyB, q_{1u} , q_{3u} and q_{4u} are the light-induced synthesis rate through photoreceptors PhyA, PhyB and Cry1, respectively, $[T_{PhyA}]$, $[T_{PhyB}]$ and $[T_{Cry1}]$ are the total concentration of the three photoreceptors, (see Supplementary Table 2) I_{red} and I_{blue} are the light intensities for red and blue light, respectively, η_1 and η_2 are the normalisation parameter of light intensity, and Θ_{PhyA} , Θ_{PhyB} and Θ_{Cry1} are the light modes that take either 0 or 1, where 0 indicates dark while 1 indicates light. We used the initial condition of 1 for all the ODEs except [HYP], [COP1:PhyA], [COP1:PhyB] and [COP1:Cry], where zero initial conditions are used.

In (Pay et al., 2022b,a) only one light colour (either blue, red or mix of red and blue) is considered within the duration of the photoperiod. For example, if the photoperiod is 10 hours, only monochromatic red or blue lights or a mixture of red and blue can be present in the entire duration of the 10 hours. In this study, modifications have been made to the model to enable the insertion of multiple light colours within the duration of the photoperiod. For example, within this 10 hour, we can now have monochromatic red light for 4 hours followed by monochromatic blue light for 6 hours or vice versa. This modification allows us to investigate the effect of multiple light colours and their order within the photoperiod on plant growth.

Plant Growth Indices

The output of the *Arabidopsis thaliana* mathematical model are the hypocotyl and flowering pathways (see Figure 1). The two indices that are used as proxy for plant growth are Days-to-Flower (DTF) and hypocotyl length, which can be calculated from the two circadian genes namely FLOWERING TIME (FT) and PHYTOCHROME INTERACTING FACTOR (PIF) as follow.

Days-to-Flower (DTF) Calculation

The calculation of DTF is given by,

$$DTF \text{ (days)} = d_0 + \frac{a_0}{1 - \left(\frac{FT_{area}}{a_1} \right)} \tag{7}$$

Equation (7) is an empirical equation where FT_{area} is the area under the curve for gene FT with unit (a.u.).day. a_0 is the function multiplier, a_1 is the lower limit of the area under the curve for gene FT and d_0 is the minimum flowering time. These three parameters are the tuneable parameters used to fit the experimental data (Corbesier et al., 1996; Salazar et al., 2009; Seaton et al., 2015). In this study, $d_0=16.55$ days, $a_0=1355.22$ days and $a_1=0.02$ (a.u.).days are considered, which have been estimated and calibrated in our previous work (Pay et al., 2022a).

Hypocotyl Length Calculation

The calculation of the hypocotyl length is given by,

$$\text{Hypocotyl length (mm)} = \int_0^D g_1 + \frac{g_2 [PIF]_p^2}{K_{12}^2 + [PIF]_p^2} dt \quad (8)$$

Equation (8) is taken from (De Caluwé et al., 2016), where a Hill function parameterised by g_2 and K_{12} with a basal level g_1 are used to relate the hypocotyl length to the accumulation of protein PIF across the number of days, D . In this study, we consider $D = 10$ days, while $g_1 = 0.01 \text{ mmh}^{-1}$, $g_2 = 0.1 \text{ mmh}^{-1}$ and $K_{12} = 0.56 \text{ nM}$, which have been obtained from our previous work (Pay et al., 2022b). The calculation of DTF is rounded to the closest integer, while the hypocotyl length is rounded to two decimal places.

The simulation is carried over 10 days where the DTF and hypocotyl length calculated in this study are compared against the nominal values taken from (Pereira et al., 2021) when simulated under white light condition. Simulation length of 10 days is chosen following the same number of days used in (Pereira et al., 2021). Moreover, the model obtained in (Pay et al., 2022b,a) are calibrated against experiment data when *Arabidopsis thaliana* is at 10-day old.

The nominal DTF and hypocotyl length of *Arabidopsis thaliana* taken from (Pereira et al., 2021) are 21.62 days and 1.18 mm, respectively and a shorter DTF and a hypocotyl length that is at least 1.00 mm are considered as improvement to its corresponding nominal values. The minimum acceptable hypocotyl length is set to 1.00 mm as further reduction of hypocotyl length can be a sign or cause of stunted growth (Derbyshire et al., 2007). Recall that these two nominal values were obtained using purely white light and, in this study, it is of interest to investigate whether the use of different light qualities at their associated properties could further improve plant growth compared to using white light.

Model Fidelity

The simulation model used in this study has been previously calibrated and validated against experimental data from *Arabidopsis thaliana* in our previous studies (Pay et al., 2022b,a). Specifically, the hypocotyl elongation and flowering time predicted from the simulation model under different light conditions are in agreement with experimental findings, which have been summarised in Table 3 of (Pay et al., 2022b) and Figure 2 of (Pay et al., 2022a), respectively.

To further evaluate the predictive capability of our simulation model, we compare our simulated outputs with experimental findings using other plants from published literature. The comparison is summarised in Table 1 and shows a good agreement with *Arabidopsis thaliana*. For other plant types, the simulated outputs show good agreement qualitatively, where similar trends are observed across different plant types and phenotypes, considering that our model is calibrated using *Arabidopsis thaliana*. Taken altogether, the comparison shown in Table 1 indicates the good predictive capability of our model, thereby warranting its use for our analysis in this study.

Light Properties for Simulation

Four light properties namely light ratio, light intensity, photoperiod, and light order are analysed using the *Arabidopsis thaliana* mathematical model as these light properties are often being considered whenever plants are grown in smart greenhouse farming as obtaining the optimal combination of those light properties can improve growth and lighting efficiency (Ahmed et al., 2020). The input light has a period of 24 hours, where the pattern of light is repeated across 10 days. A brief description on how we set up each of the light property for simulation is provided.

Light Ratio

For our simulation, the red: blue and blue: red ratios are varied from 1:1 to 12:1 to cover all the possible light ratios listed in Supplementary Table 1. This is to allow us to identify possible different red and blue light ratios that are able to produce better results in terms of DTF and hypocotyl length than using either monochromatic blue or red light. The variation of the light ratios is reflected through light intensity for red or blue lights, i.e., I_{red} and I_{blue} in Equation (6). For example, to realise a light ratio of blue: red of 3:1, we set $I_{blue} = 600 \mu\text{mol}/\text{m}^2\text{s}^{-1}$ and $I_{red} = 200 \mu\text{mol}/\text{m}^2\text{s}^{-1}$.

Light Intensity

While the daily light integral (DLI) is often used as a more representative measure of the rate of photosynthetically active radiation over a 24-hour period in plant (Faust et al., 2005; Faust and Logan, 2018), in this study, we consider the light intensity (in terms of PPFD) instead. This is to ensure our simulation analysis is comparable with the experimental studies from the literature given in Supplementary Table 1, which mostly consider light intensity rather than DLI.

The light intensity for red and blue lights, i.e., I_{red} and I_{blue} in Equation (6) used in the simulation is ranged from 100 to 1200 $\mu\text{mol}/\text{m}^2\text{s}^{-1}$, with a 100 $\mu\text{mol}/\text{m}^2\text{s}^{-1}$ increment, totalling up to a total of 12 different intensities, which is similar to the light intensity range considered in (Hidaka et al., 2013).

The light intensity of 100 to 400 $\mu\text{mol}/\text{m}^2\text{s}^{-1}$, is categorised as lower range due to its relatively significant change of photosynthetic rate as identified in (Hidaka et al., 2013). This range allows improvements to be easily observed due to the relatively larger change to DTF and hypocotyl length due to the increase in photosynthetic rate, and still retains the possibility of producing a light combination that provides improvement over the nominal values. The light combination that can provide improvement to the nominal value within the lower range of light intensity is more likely to be selected as the optimal one, as lower intensities can reduce the amount of energy needed per plant grown.

The light intensity of 500 to 800 $\mu\text{mol}/\text{m}^2\text{s}^{-1}$ is categorised as medium range as this range is the limit between the intensities used in (Fu et al., 2012) and the light saturation point observed for strawberries in (Hidaka et al., 2013). Any improvements to the plant growth from the nominal values in the medium range are likely to be attributed to the plant yet to experience stronger light stress, as the photosynthetic rate is still increasing, albeit at a slower rate compared to the photosynthetic range in the lower range.

Results and Discussion

Effect of Only Monochromatic Blue and Red Lights

We begin our analysis by considering only the effect of monochromatic light, (i.e., light conditions B1, Reverse-B1, R1 and Reverse-R1) on DTF and hypocotyl length. These light conditions are first considered given their wide usage in experiments as shown in Supplementary Table 1. Due to the monochromatic nature, only the variation in photoperiods and light intensities are considered and the results are shown in Figure 4 while Table 2 presents the best improvement in DTF and hypocotyl length that can be achieved under these light conditions.

From Figure 4, we observe improvements in DTF and hypocotyl length when light conditions B1 and Reverse-B1 are used. Across different light intensities, the DTF portrays a decreasing trend as photoperiods increase. The smallest DTF of 19 days, which is a 12% reduction compared to the nominal DTF value of 21.62 days can be observed with light intensity of at least $900 \mu\text{mol}/\text{m}^2\text{s}^{-1}$, at photoperiod of 16 hours. The hypocotyl length is also showing a decreasing trend as photoperiods increase but it is not significantly affected by light intensity given the plots at different intensities almost overlap each other. In the B1 light condition, the hypocotyl length has reduced to the threshold of 1.00 mm at photoperiod of 14 hours but the DTF is still not at its minimum value. On the other hand, in Reverse-B1 light condition, the hypocotyl length reaches the 1.00 mm threshold at photoperiod of 16 hours that at the same time the DTF is at its minimum.

Fig. 4. DTF and hypocotyl length under light conditions B1, Reverse-B1, R1 and Reverse-R1 across different photoperiod and light intensities. The grey dashed-lines in both DTF and hypocotyl length plots represent the nominal values and the black dashed-line in hypocotyl length plots represent the 1.00 mm threshold.

For R1 and Reverse-R1 light conditions, we observe no light intensity nor photoperiod can improve the DTF below the nominal value despite the hypocotyl length displaying similar behaviour as the B1 and Reverse-B1 light conditions. The minimum DTF that both these light conditions is approximately 30 days at light intensity of $1200 \mu\text{mol}/\text{m}^2\text{s}^{-1}$ and photoperiod of 16 hours, which is an increase of 39% from the nominal value. We also note the oscillatory trend in DTF for R1 and Reverse-R1 light conditions as photoperiod changes, which can be attributed to the following reasons. Flowering is accelerated under very short day (photoperiod ≤ 3 hours) conditions before decelerating under short day conditions (Luccioni et al., 2019) leading to the decrease and increase in DTF from photoperiod 2 to 10 hours. Furthermore, (Mockler et al., 2003) found that the difference in DTF between very short day to short day with long day conditions is small, leading to the decrease in DTF as photoperiod increases, resulting in the observed oscillatory trend. This oscillatory trend is also occurring in B1 and Reverse-B1 light conditions albeit not as apparent as the R1 and Reverse-R1 light conditions as the difference in DTF between very short day to short day with long day conditions is large (Mockler et al., 2003).

Taken altogether, these results suggest that monochromatic blue light plays a vital role in improving flowering compared to monochromatic red light. Furthermore, the Reverse-B1 light with photoperiod of 16 hours is more preferred as this light condition can improve both the plant growth indices. More encouragingly, our results align with previous experimental studies (Eskins, 1992; Guo et al., 1998; Spaninks et al., 2020), where long day monochromatic blue light has been shown to decrease the days required for flowering and produces hypocotyl length within a desirable limit.

Effect of Light Order

Previously, our analysis on monochromatic light indicates that Reverse-B1 light condition with a photoperiod of 16 hours produces the best improvement for plant growth and development. Here, we extend our analysis by considering the effect of different light orders on DTF and hypocotyl length. Specifically, we are interested to know whether any further improvement can be achieved when light ratios and light orders are now considered. We introduce light conditions B2 to R5 with their reverse (i.e., the order of dark and light reversed) and repeat the simulation to obtain the DTF and hypocotyl length. These light conditions have either the blue or red lights first in the order. Note that our light order analysis is an extended version of the commonly used alternating light considered in experimental studies (Ohtake et al., 2018; Lanoue et al., 2019; Viršilė et al., 2020; Huang et al., 2021), where in those studies, the equivalent of our B2, B3, R2 and R3 with photoperiods of 24 are used. In addition, we would like to remark that the improvement observed through simulation in the two plant growth indices under different light orders is consistent with the findings of (Ohtake et al., 2018). The best results from these light conditions are given in Tables 3 to 5. For the details of all the simulation results, see the Data Availability section.

From Tables 3 to 5, we observe that largest improvement to the two plant growth indices is obtained with photoperiod of 16 hours with an 8-hour dark precedes them i.e., Reverse-B2, Reverse-B3, Reverse-B4, Reverse-B5, Reverse-R4 and Reverse-R5, which is similar to the monochromatic blue light Reverse-B1 light condition. Within this 16-hour photoperiod, the duration of blue or mix blue and red lights is at least 12 hours suggesting prolonged blue light exposure is essential. All these lighting conditions produce DTF of 19 days and hypocotyl length between 1.14 mm to 1.17 mm. For the remaining light conditions, we observe DTF between 20 to 30 days and hypocotyl length between 1.00 to 1.75 mm. An exception to this trend is observed for light condition B2, where a photoperiod of 14 hours (8-hour blue and 6-hour red) followed by 10 hours of dark can produce the same DTF of 19 days and hypocotyl length of 1.00 mm.

Next, we introduce additional four light conditions where the mix of blue and red light is the first in order, i.e., M1 to M4 with their reverse counterparts (third column of Figure 3). With the introduction of these light conditions, we obtain DTF ranges between 18 to 24 days, while the hypocotyl length ranges between 1.00 to 1.17 mm. While it is compelling to conclude that the light condition that achieve DTF of 18 days is of interest; all these light conditions produce minimal improvement to the hypocotyl length from the nominal value. If a balance of improvement to both DTF and hypocotyl length are required, then M3 and M4 light conditions with photoperiod of 14 hours followed by 10 hours dark fit this requirement. Again, if we consider from the perspective of energy savings, M3, M4 and B2 light conditions are the top lighting candidates as similar DTF and hypocotyl length can be achieved with another further two hours reduction of photoperiod compared to using monochromatic light.

One probable explanation for the difference in the hypocotyl length when the light order is reversed is as follow. Given that the initial conditions of the ODEs are the same, in the reverse light condition, the circadian clock may perceive it is as day even though it is night,

and begin employing the 10-day adaptation resulting in a longer hypocotyl length. Conversely, when the light condition is not reversed, the circadian clock may have achieved synchrony with light conditions from the onset of light thereby resulting in a shorter hypocotyl length.

We note an interesting observation related to hypocotyl length in the presence of red regardless of its duration and intensity. As long as red light is available, the hypocotyl length is sufficiently long to avert stunted growth and is not affected by photoperiod compared to the presence of blue light, which is influenced by photoperiod and order of dark to attain minimum hypocotyl length. In addition, the presence of dominant intensity of blue light being first in the light order (e.g., M4 or B2 light conditions) provide the most improvement particularly to DTF. These two observations, which are also consistent with previous studies suggest that the presence of red light ensures the hypocotyl grow sufficiently (Takase et al., 2003; Sellaro et al., 2009) while the blue light helps with the promotion of flowering (Costine et al., 2022).

Effect of Light Ratio and Light Intensity

After establishing that the light conditions M3, M4 and B2 with photoperiod of 14 hours being the desired light condition for plant growth, where no further improvement in DTF or hypocotyl can be made, we turn our attention to investigating what light ratio and light intensity entail, focussing on energy efficiency. While these three light conditions yield identical DTF and hypocotyl length, their light ratio and light intensity are different. For M3, the light ratio B:R of 1:1 with intensity of $900 \mu\text{mol}/\text{m}^2\text{s}^{-1}$ is used. For M4, the light ratio B:R of 12:1 with blue intensity of $800 \mu\text{mol}/\text{m}^2\text{s}^{-1}$ and red intensity of $66.67 \mu\text{mol}/\text{m}^2\text{s}^{-1}$ are used. Lastly, for B2, the light ratio B:R of 12:1 but with blue intensity of $1000 \mu\text{mol}/\text{m}^2\text{s}^{-1}$ and red intensity of $83.33 \mu\text{mol}/\text{m}^2\text{s}^{-1}$ are used. Consider that energy efficiency is inversely proportional to light intensity (Yan et al., 2019), light conditions with smaller light intensity would be preferred. If we consider aggregated use of these lights, we have a total intensities of $1800 \mu\text{mol}/\text{m}^2\text{s}^{-1}$, $866.67 \mu\text{mol}/\text{m}^2\text{s}^{-1}$ and $1083.33 \mu\text{mol}/\text{m}^2\text{s}^{-1}$, for M3, M4 and B2 light conditions, respectively. This indicates that among the three light conditions, M3 uses the most energy, while M4 uses the least energy, thereby making M4 the preferred light conditions among the three.

Effect of the Order of Dark

In general, when implemented in smart greenhouse environment, the light-dark cycle that the plant is exposed to is repeating over 24 hours period. Here, we are interested to investigate whether does the order of dark (i.e., dark precedes or succeeds light) matter. In other words, when the plant is transferred to the smart greenhouse environment, should the plant be exposed to dark that precedes or succeeds light.

In Tables 2 to 5, we have provided the best DTF and hypocotyl length measurements from each of the light conditions. We also collate the DTF and hypocotyl length where the order of dark is reversed for each of the light conditions shown in Tables 2 to 5. Note that the best improvement in plant growth indices in each light condition shown in Tables 2 to 5 may not necessarily means its reverse light conditions would provide the best improvement in plant growth indices as well. For example, in Table 2, the best improvement of plant growth indices for light condition B1 occurs at 12B12D with DTF of 20 days and hypocotyl length of 1.81 mm. Under the same photoperiod but with its order of dark reverse i.e., Reverse-B1 with 12D12B, the DTF is 20 days and the hypocotyl length is 2.65 mm. For convenience of analysis, we call this scenario, 'Best Light First'. Similarly, in Table 2, the best improvement of plant growth indices for Reverse-B1 occurs at 8D16B with DTF of 19 days and hypocotyl of 1.15 mm. Under the same photoperiod but with now dark succeeds light i.e., B1 with 16B8D, we have DTF of 19 days and hypocotyl length of 0.46 mm and we call this scenario 'Best Dark First'. With this we end up with collating DTF and hypocotyl length for 56 different light conditions.

Our analysis (Figure S1 in the Supplementary Information) suggest the order of dark has no strong influence on DTF but has significant influence on hypocotyl length. The substantial variation of hypocotyl length between order of dark is not surprising given that short or long hypocotyl have been observed when plant is either grown in darkness (skotomorphogenesis) or in the presence of light (photomorphogenesis) (Vandenbussche et al., 2005). Our results indicate that the order of dark can be potentially exploited to provide competitive advantage to plants depends on the germination type of the plant, i.e., between hypogeal (germination below ground) or epigeal (germination above ground) germination.

Power Consumption Calculation

In this section, we provide simple calculation of power consumption of the suggested light conditions for comparison of their energy usage given that lighting power consumption has often been seen as the impediment to realise sustainability in smart farming (Ahmed et al., 2023). The power consumption model, which is based on (Ahn et al., 2017; Lan and Tan, 2015) is given by,

$$\text{Power, P[kW]} = \frac{I_{PPFD} \times C \times A}{\eta \times 1000} \quad (9)$$

where I_{PPFD} is the light intensity in $\mu\text{mol}/\text{m}^2\text{s}^{-1}$, C is the illuminance to PPFD conversion factor, A is the grow area in m^2 , 1000 is the conversion to kW and η is the luminous efficacy in lm/W .

The Light Emitting Diode (LED) lights are typically powered by an LED driver, which is an AC/DC converter that converts AC grid power into DC power suitable for the LED lights. As such, we can rewrite the combined luminous efficacy as $\eta_{comb} = \eta \times \eta_{PCE}$, where η_{PCE} is the AC/DC power conversion efficiency that takes a value between 0 and 1. We can then rewrite Equation (9) as

$$\text{Power, P[kW]} = \frac{I_{PPFD} \times C \times A}{\underbrace{\eta \times \eta_{PCE}}_{\eta_{comb}} \times 1000} \quad (10)$$

As the typical electricity billing unit is kWh, to convert kW from Equation (10) to kWh, we multiply Equation (10) with the photoperiod and the DTF, i.e.,

$$\text{Power, P[kW]} = \frac{I_{PPFD} \times C \times A}{\eta \times \eta_{PCE} \times 1000} \times \text{Photoperiod} \times \text{DTF} \quad (11)$$

In our calculation, we assume that a 1.2m by 1.2m grow area is used resulting in $A = 1.44 \text{ m}^2$. Conventionally, an LED driver is considered efficient if $\eta_{PCE} = 0.9$, i.e., the LED driver is able to convert 90% of the input power into useful electricity (Esteki et al., 2023). The typical values of C and η for red and blue lights are adopted from (Ahn et al., 2017), while for white light, they are adopted from (Ahn et al., 2017; Lee and Kim, 2012; Masoud and Murnick, 2013). The parameters associated with white light are included for comparison with the power consumption based on the light condition suggested in (Pereira et al., 2021). All these parameters are summarised in Table 6.

Using Equation (11) and the parameters given in Table 6, we computed the power consumption for three light conditions that produces the best growth indices, under monochromatic light (Reverse-B1), mixed light (M4) and the white light conditions of 18L6D from (Pereira et al., 2021) and they are given in Table 7. For the details of the calculation, see Supplementary Table 3.

From Table 7 we observe that among the three light conditions, the power consumption is the least for M4 light condition. Interestingly, despite Reverse-B1 uses two hours less in photoperiod compared to white light, the power consumption is almost double primarily due to the larger I_{PPFD} in blue light compared to white light. Nevertheless, it is encouraging to note that by using different light colour combination such as M4 instead of only monochromatic light, the power consumption can be reduced while attaining good growth indices.

Conclusions

There have been extensive studies on the effect of light qualities on plant development as evidenced by the numerous studies summarised in Supplementary Table 1. Despite the extensiveness, most of these studies often consider only a small portion of the light properties primarily due to the arduous experimental task and different scope of study. The community would benefit tremendously if there were a methodical approach in narrowing down the range of light properties to be investigated. In this study, we have carried out a systematic analysis in identifying the combination of the most suitable blue and red light properties for enhanced plant growth that can be used in smart greenhouse farming. The *in silico* analyses are carried out using the *Arabidopsis thaliana* mathematical model that relates light input to flowering and hypocotyl length output pathways, where the comparison is made against a nominal value obtained from a previous study using only white light (Pereira et al., 2021). To ensure all the blue and red light properties are considered, we conducted a comprehensive literature search on experimental studies that investigate the relationship between light quality and plant growth (Supplementary Table 1). From that search, we are able to narrow down to four key blue and red light properties namely light ratio, light intensity, light photoperiod and light order (see also Figure 1) that have significant effect on plant development, which are also highlighted in a review paper in (Ahmed et al., 2020).

For monochromatic light, our analyses show that a photoperiod of 8-hour darkness followed by 16-hour blue light with intensity of $900 \mu\text{mol}/\text{m}^2\text{s}^{-1}$ (Reverse-B1) can reduce the DTF from the nominal value of 21.62 days to 19 days and hypocotyl length from the nominal value of 1.18 mm to 1.15 mm, which is an improvement by 12% and 3%, respectively.

We further our investigation by considering the combination involving different light colours as depicted in Figure 3. Interestingly, when combinations of different light colours are considered, we observe similar improvement of DTF and hypocotyl length can be achieved with another further 2 hours reduction in photoperiod. The M4 light condition, which has a 14-hour light that is made up of 8-hour mix blue and red with intensity of $800 \mu\text{mol}/\text{m}^2\text{s}^{-1}$ and $66.67 \mu\text{mol}/\text{m}^2\text{s}^{-1}$, respectively, 6-hour red light, followed by 10-hour dark yields DTF of 19 days and hypocotyl length of 1.00 mm. This further reduction of 2 hours from using monochromatic light is a welcoming finding as this could further reduce energy usage while achieving similar outcome.

In terms of light ratio and light intensity, the ratio of blue: red ranging from 10:1 to 12:1 with intensity of red ranging $66.67 \mu\text{mol}/\text{m}^2\text{s}^{-1}$ to $100 \mu\text{mol}/\text{m}^2\text{s}^{-1}$ and blue ranging from $800 \mu\text{mol}/\text{m}^2\text{s}^{-1}$ to $1000 \mu\text{mol}/\text{m}^2\text{s}^{-1}$ could yield comparable improvements in DTF and hypocotyl length compared to the nominal values. These light ranges concur with the finding from (Piovene et al., 2015) where strawberries and basil seem to grow well under these light ranges. If energy efficiency is to be taken into account, lower light intensity is preferred, and our analysis tips the balance to M4 light condition being the most energy efficient as shown through the power consumption calculation given in Table 7.

Finally, we found that the order of dark has marginal effect on flowering but has a significant effect on hypocotyl length. Our analysis, however, can help practitioners strategise ways to take advantage of the order of darkness and give plants that germinate through either epigeal or hypogeal processes their particular competitive advantage.

The findings from our study have a great prospect in aiding smart greenhouse farming practitioners particularly those employing speed breeding protocol in setting the appropriate lighting system in their farming practices. Our model can be used to provide an optimal combination of the use of different light qualities to promote the desired plant growth and development pathways. Moreover, the resulting optimal combination of the light qualities has the potential to reduce energy consumption. Energy consumption has always been an issue in speed breeding-based farming practice (Jahne et al., 2020), where energy consumption from light alone accounts for more than 30% of the total production cost (Eaves and Eaves, 2018). As illustrated in Table 7, the use of optimal light combinations such as M4 light condition can reduce energy consumption by 13.3% compared to white light, which is often used in speed breeding-based farming practices.

Competing interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data Availability

All the MATLAB simulation codes, and simulated data are available at <https://github.com/mathiasfoo/lightquality>.

Author contributions statement

A.M.H.C.: Data curation, Formal analysis, Investigation, Methodology, Writing – original draft, Writing – review and editing, M.L.P.: Methodology, Resources, Software, Validation, Writing – original draft, Writing – review and editing, J.C.: Supervision, Validation, Writing – review and editing, F.H.: Funding acquisition, Supervision, Validation, Writing – review and editing, L.C.R.: Supervision, Resources, Writing – original draft, Writing – review and editing, H.A.: Funding acquisition, Supervision, Validation, Methodology, Writing – review and editing, M.F.: Conceptualisation, Formal analysis, Supervision, Project administration, Funding acquisition, Writing – original draft, Writing – review and editing.

Acknowledgments

A.M.H.C. and M.F. thank the researchers from the School of Life Science, University of Warwick for the useful discussion on light and plant growth. M.L.P. acknowledges the financial support from Coventry University through Global Challenge Research Fund (GCRF), Fully Funded Doctoral Studentship Scheme.

Accepted Manuscript

References

- M. S. Ahamed, M. Sultan, D. Monfet, M. S. Rahman, Y. Zhang, A. Zahid, M. Bilal, T. A. Ahsan, and Y. Achour. A critical review on efficient thermal environment controls in indoor vertical farming. *J. Clean. Prod.*, page 138923, 2023.
- H. A. Ahmed, T. Yu-Xin, and Y. Qi-Chang. Optimal control of environmental conditions affecting lettuce plant growth in a controlled environment with artificial lighting: A review. *S. Afr. J. Bot.*, 130:75–89, 2020.
- Y. D. Ahn, S. Bae, and S.-J. Kang. Power controllable led system with increased energy efficiency using multi-sensors for plant cultivation. *Energies*, 10(10):1607, 2017.
- J. B. Aune, A. Coulibaly, and K. E. Giller. Precision farming for increased land and labour productivity in semi-arid west africa. a review. *Agron Sustain Dev.*, 37:1–10, 2017.
- Z. H. Bian, Q. C. Yang, and W. K. Liu. Effects of light quality on the accumulation of phytochemicals in vegetables produced in controlled environments: a review. *J. Sci. Food Agric.*, 95(5):869–877, 2015.
- C. S. Brown, A. C. Schuerger, and J. C. Sager. Growth and photomorphogenesis of pepper plants under red light-emitting diodes with supplemental blue or far-red lighting. *J. Am. Soc. Hortic. Sci.*, 120(5):808–813, 1995.
- Y. H. Chew, D. D. Seaton, V. Mengin, A. Flis, S. T. Mugford, G. M. George, M. Moulin, A. Hume, S. C. Zeeman, T. B. Fitzpatrick, et al. The arabidopsis framework model version 2 predicts the organism-level effects of circadian clock gene mis-regulation. *in silico Plants*, 4(2):diac010, 2022.
- K. R. Cope and B. Bugbee. Spectral effects of three types of white light-emitting diodes on plant growth and development: absolute versus relative amounts of blue light. *HortScience*, 48(4):504–509, 2013.
- L. Corbesier, I. Gadisseur, G. Silvestre, A. Jacqard, and G. Bernier. Design in arabidopsis thaliana of a synchronous system of floral induction by one long day. *Plant J.*, 9(6):947–952, 1996.
- B. Costine, M. Zhang, B. Pearson, and S. S. Nadakuduti. Impact of blue light on plant growth, flowering and accumulation of medicinal flavones in scutellaria baicalensis and s. lateriflora. *Horticulturae*, 8(12):1141, 2022.
- J. De Caluwé, Q. Xiao, C. Hermans, N. Verbruggen, J.-C. Leloup, and D. Gonze. A compact model for the complex plant circadian clock. *Front. Plant. Sci.*, 7:74, 2016.
- P. Derbyshire, M. C. McCann, and K. Roberts. Restricted cell elongation in arabidopsis hypocotyls is associated with a reduced average pectin esterification level. *BMC Plant Biol.*, 7(1):1–12, 2007.
- J. Eaves and S. Eaves. Comparing the profitability of a greenhouse to a vertical farm in quebec. *Can. J. Agric. Econ.*, 66(1):43–54, 2018.
- K. Eskins. Light-quality effects on arabidopsis development. red, blue and far-red regulation of flowering and morphology. *Physiol. Plant.*, 86(3):439–444, 1992.
- M. Esteki, S. A. Khajehoddin, A. Safaee, and Y. Li. Led systems applications and led driver topologies: a review. *IEEE Access*, 11:38324–38358, 2023.
- J. E. Faust and J. Logan. Daily light integral: a research review and high-resolution maps of the united states. *HortScience*, 53(9):1250–1257, 2018.
- J. E. Faust, V. Holcombe, N. C. Rajapakse, and D. R. Layne. The effect of daily light integral on bedding plant growth and flowering. *HortScience*, 40(3):645–649, 2005.
- K. M. Folta and K. S. Childers. Light as a growth regulator: Controlling plant biology with narrow-bandwidth solid-state lighting systems. *HortScience*, 43(7):1957–1964, 2008.
- B. Fraszczk. The effect of different doses of blue light on the biometric traits and photosynthesis of dill plants. *Not. Bot. Horti. Agrobi.*, 44(1):34–40, 2016.
- W. Fu, P. Li, and Y. Wu. Effects of different light intensities on chlorophyll fluorescence characteristics and yield in lettuce. *Sci. Hortic.*, 135:45–51, 2012.
- B. H. Gangadhar, R. K. Mishra, G. Pandian, and S. W. Park. Comparative study of color, pungency, and biochemical composition in chili pepper (capsicum annuum) under different light-emitting diode treatments. *HortScience*, 47(12):1729–1735, 2012.
- P. Gautam, M. T. Terfa, J. E. Olsen, and S. Torre. Red and blue light effects on morphology and flowering of petunia × hybrida. *Sci. Hortic.*, 184:171–178, 2015.
- S. Ghosh, A. Watson, O. E. Gonzalez-Navarro, R. H. Ramirez-Gonzalez, L. Yanes, M. Mendoza-Suárez, J. Simmonds, R. Wells, T. Rayner, P. Green, et al. Speed breeding in growth chambers and glasshouses for crop breeding and model plant research. *Nat. Protoc.*, 13(12):2944–2963, 2018.
- H. Guo, H. Yang, T. C. Mockler, and C. Lin. Regulation of flowering time by arabidopsis photoreceptors. *Science*, 279(5355):1360–1363, 1998.
- R. Hernández and C. Kubota. Growth and morphological response of cucumber seedlings to supplemental red and blue photon flux ratios under varied solar daily light integrals. *Sci Hortic.*, 173:92–99, 2014.
- K. Hidaka, K. Dan, H. Imamura, Y. Miyoshi, T. Takayama, K. Sameshima, M. Kitano, and M. Okimura. Effect of supplemental lighting from different light sources on growth and yield of strawberry. *Environ. Control. Biol.*, 51(1):41–47, 2013.
- M. Hoenecke, R. Bula, and T. Tibbitts. Importance of blue photon levels for lettuce seedlings grown under red-light-emitting diodes. *HortScience*, 27(5):427–430, 1992.
- S. Hogewoning, H. Maljaars, and J. Harbinson. The acclimation of photosynthesis in cucumber leaves to different ratios of red and blue light. *Photosynth. Res.*, 91(2-3):287–288, 2007.
- J. Huang, Y.-l. Xu, F.-m. Duan, X. Du, Q.-c. Yang, and Y.-j. Zheng. Improvement of the growth and nutritional quality of two-leaf-color pak choi by supplemental alternating red and blue light. *HortScience*, 56(2):118–125, 2021.
- F. Jahne, V. Hahn, T. Wurschum, and W. L. Leiser. Speed breeding short-day crops by led-controlled light schemes. *Theor. Appl. Genet.*, 133(8):2335–2342, 2020.

- C. Kami, S. Lorrain, P. Hornitschek, and C. Fankhauser. Light-regulated plant growth and development. *Curr. Top. Dev. Biol.*, 91:29–66, 2010.
- T. Karanisa, Y. Achour, A. Ouammi, and S. Sayadi. Smart greenhouses as the path towards precision agriculture in the food-energy and water nexus: Case study of qatar. *Environ. Syst. Decis.*, 42(4):521–546, 2022.
- S. Kumar, G. Chowdhary, V. Udutalappally, D. Das, and S. P. Mohanty. Gerop: Internet-of-leaf-things (iolt) for monitoring of the growth of crops in smart agriculture. In *2019 IEEE International Symposium on Smart Electronic Systems (iSES)(Formerly iNiS)*, pages 53–56. IEEE, 2019.
- Y. Kwack, K. K. Kim, H. Hwang, and C. Chun. Growth and quality of sprouts of six vegetables cultivated under different light intensity and quality. *Hortic. Environ. Biotechnol.*, 56:437–443, 2015.
- L. Lan and Y. K. Tan. Advanced building energy monitoring using wireless sensor integrated energyplus platform for personal climate control. In *2015 IEEE 11th International Conference on Power Electronics and Drive Systems*, pages 567–574. IEEE, 2015.
- J. Lanoue, J. Zheng, C. Little, A. Thibodeau, B. Grodzinski, and X. Hao. Alternating red and blue light-emitting diodes allows for injury-free tomato production with continuous lighting. *Front. Plant Sci.*, 10:1114, 2019.
- J.-G. Lee, S.-S. Oh, S.-H. Cha, Y.-A. Jang, S.-Y. Kim, Y.-C. Um, and S.-R. Cheong. Effects of red/blue light ratio and short-term light quality conversion on growth and anthocyanin contents of baby leaf lettuce. *J. Bio-Env. Con.*, 19(4):351–359, 2010.
- W.-S. Lee and S.-G. Kim. Development of rotational smart lighting control system for plant factory. *World Acad. Eng. Technol.*, 62:741–744, 2012.
- L. Luccioni, M. Krzymuski, M. Sanchez-Lamas, E. Karayekov, P. D. Cerdan, and J. J. Casal. Constans delays arabidopsis flowering under short days. *Plant J.*, 97(5):923–932, 2019.
- N. Masoud and D. Murnick. High efficiency, fluorescent excimer lamps, an alternative to cfls and white light leds. *J. Light Vis. Environ.*, 37(4):171–175, 2013.
- R. Matsuda, K. Ohashi-Kaneko, K. Fujiwara, and K. Kurata. Analysis of the relationship between blue-light photon flux density and the photosynthetic properties of spinach (*spinacia oleracea* l.) leaves with regard to the acclimation of photosynthesis to growth irradiance. *Soil Sci. Plant Nutr.*, 53(4):459–465, 2007.
- C. R. McClung. Beyond arabidopsis: the circadian clock in non-model plant species. *Semin. Cell Dev. Biol.*, 24(5):430–436, 2013.
- T. Mockler, H. Yang, X. Yu, D. Parikh, Y. C. Cheng, S. Dolan, and C. Lin. Regulation of photoperiodic flowering by arabidopsis photoreceptors. *Proc. Natl. Acad. Sci. U.S.A.*, 100(4):2140–2145, 2003.
- E. S. Mohamed, A. Belal, S. K. Abd-Elmabod, M. A. El-Shirbeny, A. Gad, and M. B. Zahran. Smart farming for improving agricultural management. *Egypt. J. Remote. Sens. Space Sci.*, 24(3):971–981, 2021.
- S. Mosharafian, S. Afzali, G. M. Weaver, M. van Iersel, and J. M. Velni. Optimal lighting control in greenhouse by incorporating sunlight prediction. *Comput. Electron. Agric.*, 188:106300, 2021.
- S. Muneer, E. J. Kim, J. S. Park, and J. H. Lee. Influence of green, red and blue light emitting diodes on multiprotein complex proteins and photosynthetic activity under different light intensities in lettuce leaves (*lactuca sativa* l.). *Int. J. Mol. Sci.*, 15(3):4657–4670, 2014.
- D. C. J. Neo, M. M. X. Ong, Y. Y. Lee, E. J. Teo, Q. Ong, H. Tanoto, J. Xu, K. S. Ong, and V. Suresh. Shaping and tuning lighting conditions in controlled environment agriculture: A review. *ACS Agric. Sci. Technol.*, 2(1):3–16, 2022.
- N. Ohtake, M. Ishikura, H. Suzuki, W. Yamori, and E. Goto. Continuous irradiation with alternating red and blue light enhances plant growth while keeping nutritional quality in lettuce. *HortScience*, 53(12):1804–1809, 2018.
- E. Olvera-Gonzalez, D. Alaniz-Lumbreras, R. Ivanov-Tsonchev, J. Villa-Hernández, C. Olvera-Olvera, E. González-Ramírez, M. Araiza-Esquivel, V. Torres-Argüelles, and V. Castaño. Intelligent lighting system for plant growth and development. *Comput. Electron. Agric.*, 92:48–53, 2013.
- S. A. O’Shaughnessy, M. Kim, S. Lee, Y. Kim, H. Kim, and J. Shekailo. Towards smart farming solutions in the us and south korea: A comparison of the current status. *Geogr. Environ. Sustain*, 2(4):312–327, 2021.
- R. Paradiso, C. Arena, Y. Roupheal, L. d’Aquino, K. Makris, P. Vitaglione, and S. De Pascale. Growth, photosynthetic activity and tuber quality of two potato cultivars in controlled environment as affected by light source. *Plant Biosyst.*, 153(5):725–735, 2019.
- M. L. Pay, J. Christensen, F. He, L. Roden, H. Ahmed, and M. Foo. An extended plant circadian clock model for characterising flowering time under different light quality conditions. In *2022 22nd International Conference on Control, Automation and Systems (ICCAS)*, pages 1848–1853. IEEE, 2022a.
- M. L. Pay, D. W. Kim, D. E. Somers, J. K. Kim, and M. Foo. Modelling of plant circadian clock for characterizing hypocotyl growth under different light quality conditions. *in silico Plants*, 4(1):diac001, 2022b.
- J. Pereira, A. M. Mouazen, M. Foo, and H. Ahmed. A framework of artificial light management for optimal plant development for smart greenhouse application. *PLoS One*, 16(12):e0261281, 2021.
- C. Piovene, F. Orsini, S. Bosi, R. Sanoubar, V. Bregola, G. Dinelli, and G. Gianquinto. Optimal red: blue ratio in led lighting for nutraceutical indoor horticulture. *Sci Hortic.*, 193:202–208, 2015.
- J. D. Salazar, T. Saithong, P. E. Brown, J. Foreman, J. C. Locke, K. J. Halliday, I. A. Carré, D. A. Rand, and A. J. Millar. Prediction of photoperiodic regulators from quantitative gene circuit models. *Cell*, 139(6):1170–1179, 2009.
- D. D. Seaton, R. W. Smith, Y. H. Song, D. R. MacGregor, K. Stewart, G. Steel, J. Foreman, S. Penfield, T. Imaizumi, A. J. Millar, et al. Linked circadian outputs control elongation growth and flowering in response to photoperiod and temperature. *Mol. Syst. Biol.*, 11(1):776, 2015.
- R. Sellaro, U. Hoecker, M. Yanovsky, J. Chory, and J. J. Casal. Synergism of red and blue light in the control of arabidopsis gene expression and development. *Curr. Biol.*, 19(14):1216–1220, 2009.
- Y. H. Song, S. Ito, and T. Imaizumi. Similarities in the circadian clock and photoperiodism in plants. *Curr. Opin. Plant Biol.*, 13(5):594–603, 2010.
- K. Spaninks, J. Van Lieshout, W. Van Ieperen, and R. Offringa. Regulation of early plant development by red and blue light: A comparative analysis between arabidopsis thaliana and solanum lycopersicum. *Front. Plant Sci.*, 11:599982, 2020.

- G. W. Stutte, S. Edney, and T. Skerritt. Photoregulation of bioprotectant content of red leaf lettuce with light-emitting diodes. *HortScience*, 44(1):79–82, 2009.
- T. Takase, M. Nakazawa, A. Ishikawa, K. Manabe, and M. Matsui. Dfl2, a new member of the arabidopsis gh3 gene family, is involved in red light-specific hypocotyl elongation. *Plant Cell Physiol.*, 44(10):1071–1080, 2003.
- United Nation. The future of food and agriculture: Trends and challenges. <https://www.fao.org/policy-support/tools-and-publications/resources-details/en/c/472484>, 2017. Accessed: 4 Oct 2022.
- F. Vandenbussche, J.-P. Verbelen, and D. Van Der Straeten. Of light and length: regulation of hypocotyl growth in Arabidopsis. *Bioessays*, 27(3):275–284, 2005.
- V. Vaštakaitė, A. Viršilė, A. Brazaitytė, G. Samuolienė, J. Jankauskienė, R. Sirtautas, A. Novičkovas, L. Dabašinskas, S. Sakalauskienė, J. Miliauskienė, et al. The effect of blue light dosage on growth and antioxidant properties of microgreens. *Sodinink. Daržinink.*, 34:25–35, 2015.
- A. Viršilė, J. Miliauskienė, P. J. Haimi, K. Laužikė, and G. Samuolienė. The comparison of constant and dynamic red and blue light irradiation effects on red and green leaf lettuce. *Agronomy*, 10(11):1802, 2020.
- A. Walter, R. Finger, R. Huber, and N. Buchmann. Smart farming is key to developing sustainable agriculture. *Proc. Natl. Acad. Sci. U.S.A.*, 114(24):6148–6150, 2017. doi: 10.1073/pnas.1707462114. URL <https://www.pnas.org/doi/abs/10.1073/pnas.1707462114>.
- H. Wang, M. Gu, J. Cui, K. Shi, Y. Zhou, and J. Yu. Effects of light quality on co2 assimilation, chlorophyll-fluorescence quenching, expression of calvin cycle genes and carbohydrate accumulation in cucumis sativus. *J. Photochem. Photobiol. B.*, 96(1):30–37, 2009.
- A. Watson, S. Ghosh, M. J. Williams, W. S. Cuddy, J. Simmonds, M.-D. Rey, M. Asyraf Md Hatta, A. Hinchliffe, A. Steed, D. Reynolds, et al. Speed breeding is a powerful tool to accelerate crop research and breeding. *Nat. Plants*, 4(1):23–29, 2018.
- Z. Yan, D. He, G. Niu, Q. Zhou, and Y. Qu. Growth, nutritional quality, and energy use efficiency of hydroponic lettuce as influenced by daily light integrals exposed to white versus white plus red light-emitting diodes. *HortScience*, 54(10):1737–1744, 2019.
- H. Yoshida, S. Hikosaka, E. Goto, H. Takasuna, and T. Kudou. Effects of light quality and light period on flowering of everbearing strawberry in a closed plant production system. In *VII International Symposium on Light in Horticultural Systems 956*, pages 107–112, 2012.
- X. Zhang, D. He, G. Niu, Z. Yan, and J. Song. Effects of environment lighting on the growth, photosynthesis, and quality of hydroponic lettuce in a plant factory. *Int. J. Agric. Biol.*, 11(2):33–40, 2018.
- T. Zou, C. Huang, P. Wu, L. Ge, and Y. Xu. Optimization of artificial light for spinach growth in plant factory based on orthogonal test. *Plants*, 9(4):490, 2020.

Table 1. Comparison between simulated hypocotyl length and flowering time with experimental results. Note: ‘HL’ and ‘FT’ denote hypocotyl length and flowering time, respectively.

Plant type	Light conditions	Phenotypes	Experimental	Simulated	Reference
<i>Brassica rapa</i>	16L8D, Red 275 $\mu\text{mol}/\text{m}^2\text{s}^{-1}$, Blue 25 $\mu\text{mol}/\text{m}^2\text{s}^{-1}$	HL (8 days)	3.37-3.91 mm	0.408 mm	(Vaštakaitė et al., 2015)
	16L8D, Red 250 $\mu\text{mol}/\text{m}^2\text{s}^{-1}$, Blue 50 $\mu\text{mol}/\text{m}^2\text{s}^{-1}$	HL (8 days)	3.54-4.08 mm	0.400 mm	
	16L8D, Red 225 $\mu\text{mol}/\text{m}^2\text{s}^{-1}$, Blue 75 $\mu\text{mol}/\text{m}^2\text{s}^{-1}$	HL (8 days)	2.88-3.32 mm	0.398 mm	
	16L8D, Red 200 $\mu\text{mol}/\text{m}^2\text{s}^{-1}$, Blue 100 $\mu\text{mol}/\text{m}^2\text{s}^{-1}$	HL (8 days)	2.74-3.14 mm	0.395 mm	
<i>Anethum graveolens</i>	16L8D, Red 107.8 $\mu\text{mol}/\text{m}^2\text{s}^{-1}$, Blue 9.82 $\mu\text{mol}/\text{m}^2\text{s}^{-1}$	HL (14 days)	3.0-3.7 mm	0.726 mm	(Fraszczk, 2016)
	16L8D, Red 107.8 $\mu\text{mol}/\text{m}^2\text{s}^{-1}$, Blue 23.2 $\mu\text{mol}/\text{m}^2\text{s}^{-1}$	HL (14 days)	3.1-3.4 mm	0.710 mm	
	16L8D, Red 107.8 $\mu\text{mol}/\text{m}^2\text{s}^{-1}$, Blue 33.6 $\mu\text{mol}/\text{m}^2\text{s}^{-1}$	HL (14 days)	3.2-3.4 mm	0.700 mm	
	16L8D, Red 107.8 $\mu\text{mol}/\text{m}^2\text{s}^{-1}$, Blue 39.7 $\mu\text{mol}/\text{m}^2\text{s}^{-1}$	HL (14 days)	2.9-3.1 mm	0.696 mm	
<i>Petunia × hybrida</i>	14L10D, Red 150 $\mu\text{mol}/\text{m}^2\text{s}^{-1}$	FT (35 days)	27.3-27.8 days	49.0 days	(Gautam et al., 2015)
	14L10D, Blue 150 $\mu\text{mol}/\text{m}^2\text{s}^{-1}$	FT (35 days)	25.8-26.5 days	23.7 days	
	18L6D, Red 150 $\mu\text{mol}/\text{m}^2\text{s}^{-1}$	FT (35 days)	29.8-30.3 days	43.9 days	
	18L6D, Blue 150 $\mu\text{mol}/\text{m}^2\text{s}^{-1}$	FT (35 days)	29.6-29.8 days	22.8 days	
<i>Arabidopsis thaliana</i>	14L10D Blue 164 $\mu\text{mol}/\text{m}^2\text{s}^{-1}$	FT (10 days)	20-22 days	23.3 days	(Eskins, 1992)
	14L10D Red 87 $\mu\text{mol}/\text{m}^2\text{s}^{-1}$	FT (10 days)	50-55 days	56.2 days	

Table 2. Compilation of the best improvement of DTF and hypocotyl length from nominal values for each of the monochromatic light conditions (i.e., B1 and R1 and their reverse).

Light condition	Photoperiod	Light intensity ($\mu\text{mol}/\text{m}^2\text{s}^{-1}$)	DTF (days)	Hypocotyl length (mm)
B1	12B12D	B: 900	20	1.81
Reverse-B1	8D16B	B: 900	19	1.15
R1	14R10D	R: 1200	31	1.08
Reverse-R1	8D16R	R: 1200	30	1.22

Accepted Manuscript

Table 3. Compilation of the best improvement of DTF and hypocotyl length from nominal values from the B2 to B5 light conditions and their reverse.

Light condition	Photoperiod	Light ratio and intensity ($\mu\text{mol}/\text{m}^2\text{s}^{-1}$)	DTF (days)	Hypocotyl length (mm)
B2	14L10D (8B6R10D)	B:R 12:1 R: 83.33 B: 1000	19	1.00
Reverse-B2	8D16L (8D12B4R)	R:B 1:1 R: 1200 B: 1200	19	1.17
B3	14L10D (8D12B4R)	R:B 1:1 R: 900 B: 900	20	1.00
Reverse-B3	8D16L (8D14B2R)	R:B 1:1 R: 1200 B: 1200	19	1.17
B4	12L12D (2B10M12D)	R:B 1:1 R: 1100 B: 1100	20	1.75
Reverse-B4	8D16L (8D2B14M)	B:R 12:1 R: 75 B: 900	19	1.14
B5	12L12D (2B10M12D)	R:B 1:1 R: 1100 B: 1100	20	1.75
Reverse-B5	8D16L (8D2B14M)	R:B 1:1 R: 700 B: 700	19	1.14

Table 4. Compilation of the best improvement of DTF and hypocotyl length from nominal values from the R2 to R5 and their reverse.

Light condition	Photoperiod	Light ratio and intensity ($\mu\text{mol}/\text{m}^2\text{s}^{-1}$)	DTF (days)	Hypocotyl length (mm)
R2	14L10D (2R12B10D)	R:B 7:1 R: 1100 B: 157.14	25	1.00
Reverse-R2	8D16L (8D2R14B)	R:B 1:1 R: 900 B: 900	20	1.15
R3	14L10D (10R4B10D)	B:R 1:1 R: 1100 B: 1100	26	1.00
Reverse-R3	8D16L (8D2R14B)	B:R 2:1 R: 450 B: 900	20	1.15
R4	14L10D (8R6M10D)	R:B 7:1 R: 900 B: 128.57	28	1.00
Reverse-R4	8D16L (8D2R14M)	B:R 1:1 R: 1100 B: 1100	19	1.14
R5	14L10D (12R2M10D)	B:R 1:1 R: 800 B: 800	30	1.00
Reverse-R5	8D16L (8D2R14M)	B:R 1:1 R: 1100 B: 1100	19	1.14

Table 5. Compilation of the best improvement of DTF and hypocotyl length from nominal values from the M1 to M4 light conditions and their reverse.

Light condition	Photoperiod	Light ratio and intensity ($\mu\text{mol}/\text{m}^2\text{s}^{-1}$)	DTF (days)	Hypocotyl length (mm)
M1	14L10D (4M10B10D)	B:R 1:1 R: 100 B: 100	24	1.00
Reverse-M1	8D16L (8D12M4B)	B:R 1:1 R: 1100 B: 1100	18	1.14
M2	14L10D (4M10B10D)	R:B 8:1 R: 800 B: 100	22	1.00
Reverse-M2	8D16L (8D12M4B)	B:R 1:1 R: 1100 B: 1100	18	1.14
M3	14L10D (6M8R10D)	B:R 1:1 R: 900 B: 900	19	1.00
Reverse-M3	8D16L (8D12M4R)	B:R 1:1 R: 1200 B: 1200	18	1.17
M4	14L10D (8M6R10D)	B:R 12:1 R: 66.67 B: 800	19	1.00
Reverse-M4	8D16L (8D12M4R)	B:R 1:1 R: 1200 B: 1200	18	1.17

Table 6. Parameter value used for power consumption calculation.

Parameters	Values
C (Blue)	11.9
C (Red)	9.9
C (White)	68.2
η (Blue)	26 lm/W
η (Red)	50 lm/W
η (White)	70 lm/W

Accepted Manuscript

Table 7. Comparison of power consumption for three different light conditions.

Light conditions	Power consumption
Reverse-B1 (8D16B)	
$I_{blue} = 900 \mu\text{mol}/\text{m}^2\text{s}^{-1}$	200.36
M4 (8M6R10D)	
$I_{blue} = 800 \mu\text{mol}/\text{m}^2\text{s}^{-1}$	
$I_{red} = 66.7 \mu\text{mol}/\text{m}^2\text{s}^{-1}$	94.67
White (18L6D)	
$I_{blue} = 900 \mu\text{mol}/\text{m}^2\text{s}^{-1}$	109.20

Accepted Manuscript




Fig. 1.




Fig. 2.

Accepted Manuscript




Fig. 3.




Fig. 4.